

# Geochemical characteristics and tectonic interpretation of garnet mica schists of Patharkhola area in Kumaun Lesser Himalaya, Uttarakhand Himalaya, India

<sup>1</sup>HARITABH RANA, <sup>1</sup>HAREL THOMAS and <sup>1</sup>RISHABH BATRI


<sup>1</sup>Department of Applied Geology, School of Engineering & Technology, Dr. Harisingh Gour Vishwavidyalaya, Sagar (A Central University) 470003 – INDIA

Corresponding Author: harelthomas@gmail.com; hthomas@dhsgsu.edu.in

**Abstract:** The Patharkhola area is exposed in the core of the southern limb of the Dudhatoli syncline in the Kumaun Lesser Himalaya. Garnet mica schists of the Patharkhola occur in the folded outcrop pattern being located in between the limbs. They are dominantly composed of garnet, biotite, muscovite, chlorite, plagioclase and a subordinate amount of quartz. Garnet, both xenoblastic as well as idioblastic, is wrapped round by phyllosilicates minerals. These investigated rocks exhibits slightly LREE depleted patterns ( $\text{La}_{\text{N}}/\text{Sm}_{\text{N}} = 0.7$ ,  $\text{La}_{\text{N}}/\text{Yb}_{\text{N}} = 0.64$ ) with flat HREE patterns ( $\text{Sm}_{\text{N}}/\text{Yb}_{\text{N}} = 0.89$ ). Field evidence along with geochemical characteristics suggests that the Patharkhola schists are peraluminous in nature, showing sedimentary sources having silt to silty clay protolith. They were formed in active continental margins in arc-type setting where the sediments were received both from arc and continental crust.

**Key words:** Garnet, Schist, Geochemistry, Tectonic settings, Lesser Himalaya

## Graphical abstract




## Highlights

- The garnet mica schists from Patharkhola, Kumaun Lesser Himalaya forms a part of Dudhatoli Almora group.
- The geochemical data clearly demonstrates that the Patharkhola schists are characterized by their high alkali content and silica-saturated nature exhibiting slightly LREE depleted patterns ( $\text{LaN}/\text{SmN} = 0.7$ ,  $\text{LaN}/\text{YbN} = 0.64$ ) with flat HREE patterns ( $\text{SmN}/\text{YbN} = 0.89$ ).
- Shale dominated protolith representing nearby source sediments in arc type environment is inferred for the Patharkhola schists.

## 1 Introduction

The Himalaya, an arc-shaped mountain belt covering whole boundary of northern India is a type example of intercontinental collision between Indian and Asian plates around 55 Ma ago (Mukherjee, 2015; Yin, 2006). In the past, the structure, stratigraphy and tectonics of the Kumaun Lesser Himalaya have been described by Joshi & Tiwari (2009), Joshi et al. (2017) and Rana & Thomas (2018). In Kumaun region, the Lesser Himalayan sequence is delineated by Main Central Thrust (MCT) from the Higher Himalayan in the north and by Main Boundary Thrust (MBT) from the Siwaliks in south (Thakur et al., 2010). The rocks of the Dudhatoli group (Rana & Thomas, 2018) are exposed in the Patharkhola area. The “Inner Schistose Series” or the “Metamorphic and Crystalline Nappe” Tectonic Zone” of Lesser Himalaya forms a distinct structural unit which has witnessed multiple deformation


and polyphase metamorphism. The Dudhatoli Crystallines extend from Garhwal in WNW to Kumaun in ESE. Kumar & Agarwal (1975) included Mandhali, Chandpur and Naghat Formations and Dudhatoli-Almora Crystallines under a newly constituted group namely Dudhatoli group where the Dudhatoli-Almora Crystallines are the topmost horizon. Rocks of the Dudhatoli group have been considered to be of Precambrian age (Kumar & Agarwal, 1975). In Kumaun Himalaya, the Muniari gneisses of Almora Group were dated as  $1830 \pm 200$  Ma. old (Bhanot et al., 1977). Islam et al. (2005) proposed Proterozoic granitoids of Lesser Himalaya grouped into older clusters of 2200–1800 Ma and younger clusters of 1400–1200 Ma. Rana & Thomas (2023) studied the thermal and structural character of garnets from garnet mica schists of Patharkhola which is found to be of magnesian rich showing decomposition at higher temperatures. The metamorphic terrain exhibits multiple deformation patterns



**Fig. 1a.** Generalized geological map of Himalaya, Modified after Yin (2006).

and polyphase metamorphism (Joshi & Tiwari, 2009) which is well exposed in the central part of Almora nappe.

In Lesser Kumaun Himalaya, extensive work has been carried out on the pelitic gneisses and granites, rather less on the low-grade metamorphic i.e. schists and phyllites (Phukon et al., 2018; Das et al., 2019). Rana et al. (2023) studied the phyllites of Patharkhola and opined that the phyllites have high alumina content with enrichment of trace elements formed in active continental margins. The area around Patharkhola, exposes rocks of Lesser Kumaun Himalaya, lies between longitude 79°09'E to 79°17'56"E and latitude 29°47'42"N to 29°56'69"N with an approximate area of around 120 square kilometres (Fig. 1a). The rock types exposed in the area are gneisses, schists and phyllites (Fig. 1b). The main aim of the present paper is to provide geochemical character and petrogenesis of schists.



**Fig. 1b.** Geological map of the study area (Modified after Thomas & Thomas, 1992).

## 2 Macroscopic properties

In the area under investigation, schists are highly fragile and powdery in nature. Depending upon the biotite, muscovite and/or chlorite content, they exhibit a variation in colour from pinkish brown, dark brown, light grey to greenish green, (Fig. 2a, c). Increased percentage of quartz and feldspar imparts compaction. The schists of the area are characterized by the presence of garnet which ranges in size from 0.1 cm to 0.2 cm in hand


specimens (Fig. 2a). On weathering, pink garnet crystals impart a reddish brown tinge to the rock and when removed, leave behind small pits of reddish-brown colour. These are foliated medium to coarse-grained rocks where schistosity is defined by the parallel alignment of phyllosilicates showing folding and crenulations due to deformation.

### 3 Petrography

Rock samples have been collected from the entire area of the exposure of garnet mica schist terrain. Samples showing well-developed internal structures have been studied in detail and discussed here. Schists of the area is dominantly composed of garnet, biotite, muscovite, chlorite, plagioclase and subordinate amount of quartz. Two distinct varieties of garnet have been identified. Garnet-I occurs as pre-kinematic porphyroblasts, shattered and stretched parallel to the foliation, (Fig. 2c), containing ilmenite and quartz as inclusions and are highly fractured. Garnet has both xenoblastic as well as idioblastic textures suggesting the reaction:



Garnet-II occurs as rolled garnet wrapped round by flaky minerals and imparts a closed eye structure, (Fig. 2d). Two variants of Biotite recognised as Biotite-I, (Fig. 2b, d), as coarse lepidoblastic in intimate association with Muscovite-I defining the foliation while Biotite-II occurs as cross-cutting relationship with Biotite-I. Chlorite-I occurs as lepidoblasts, light greenish in colour showing pleochroism from light green to green in colour, defining the schistosity plane in close association with Biotite-I and Muscovite-I. Chlorite-II occurs as very similar to Chlorite-I, occurring as lepidoblasts cross cutting the schistosity plane varying from sub-parallel to low angles. Tiny shreds of plagioclase occur in the schists lying parallel to the schistosity plane in close association with quartz. At a few places, it is usually altered to sericite as a common alteration product. Apatite occurs as inclusion in feldspar and muscovite.



**Fig. 2.** (a, b) Field photograph showing pitted garnet; (c) Photomicrograph of stretched garnet parallel to foliation (d) Rolled garnet wrapped by flaky minerals (under plane polarised light).

#### 4 Geochemistry and petrogenesis

The results of the major oxides, trace elements along with rare earth elements (REE) are tabulated below in Table 1. Rock samples were crushed, and then pulverized using an agate carbide ring grinder. Major oxides and selected trace element concentrations present in the rock were measured on powder pellets by X-ray Fluorescence Spectrometer (XRF; Siemens SRS-3000) in the Wadia Institute of Himalayan Geology (WIHG), Dehradun. The samples were analysed in the same institute for their REE and some trace elements by Inductively Coupled Plasma-Mass Spectrometer (ICP-MS; PerkinElmer SCIEX ELAN DRC-e) using the open system rock digestion method. Rock powders were first thoroughly dissolved in HF and HNO<sub>3</sub> in Teflon crucibles and heated over a hot plate for three hours to prepare solutions from which trace elements and REE abundances were determined. Analytical precision


for major elements is well within  $\pm 2 - 3\%$  and  $\pm 5 - 6\%$  for trace elements. Accuracy of rare earth elements ranges from 2 to 12 % and precision varies from 1 to 8 %.

The SiO<sub>2</sub> and Al<sub>2</sub>O<sub>3</sub> content varies from 64.74 to 75.78 % and from 10.53 to 19.88 %, respectively. The average Na<sub>2</sub>O and K<sub>2</sub>O content in the schists is 2.59 % and 2.67 %, respectively. The average K<sub>2</sub>O/Na<sub>2</sub>O ratio of these rocks is 1.06; whereas TiO<sub>2</sub> content of the rocks varies from 0.61 to 0.8 %, as per Herron (1988), Fig. 3 (a) plot log(Fe<sub>2</sub>O<sub>3</sub>/K<sub>2</sub>O) versus log(SiO<sub>2</sub>/Al<sub>2</sub>O<sub>3</sub>) plot for classification of terrigenous sediments and shale, all studied samples of garnet mica schists fall within the shale to wacke-shale junction field specified for silty to silty clay protolith. This is due to the high Fe<sub>2</sub>O<sub>3</sub>/K<sub>2</sub>O ratio with less SiO<sub>2</sub>/Al<sub>2</sub>O<sub>3</sub> ratio representing that the schists have been derived from shale-dominated environment.

The A/CNK values of schists ranging from 1.8 to 2.8

support its characterization as a strongly peraluminous, relatively potassic-rich source. The plot of K<sub>2</sub>O/Al<sub>2</sub>O<sub>3</sub> versus Na<sub>2</sub>O/Al<sub>2</sub>O<sub>3</sub>, (Fig. 3b) has clearly differentiated the sedimentary from igneous rocks, Garrels and Mackenzie (1971). It is evident that all samples fall within the field specified for sedimentary rock. The plot between SiO<sub>2</sub> and other oxides, (Fig. 3c) shows good correlation, which indicates coherent behaviour of elements during different processes. SiO<sub>2</sub> versus Al<sub>2</sub>O<sub>3</sub>, CaO, K<sub>2</sub>O, MnO and TiO<sub>2</sub> have shown positive regression while others have poor regression values. The representation of good regression values of alumina and potash with silica represents the dominance of peraluminous nature and is substantially supported by the good regression of silica with Rb, La, Y and Ce. The behaviour of trace elements in the schists has also been studied with the help of variation diagrams presented in (Fig. 3d). Trace elements show good correlation with respect to SiO<sub>2</sub>.

During metamorphism, the REE are very little fractionated and also immobile during sedimentary processes. It is also considered that during metamorphism the REE remains unaffected up to the upper amphibolite facies regional metamorphism, (Taylor & McLennan, 1985). On the other hand,



**Fig. 3.** (a) log (SiO<sub>2</sub>/Al<sub>2</sub>O<sub>3</sub>) versus log (Fe<sub>2</sub>O<sub>3</sub>/K<sub>2</sub>O) after Herron (1988) for classification of terrigenous sediments and shales; (b) K<sub>2</sub>O/Al<sub>2</sub>O<sub>3</sub> versus Na<sub>2</sub>O/Al<sub>2</sub>O<sub>3</sub> after Garrels & Mackenzie (1971).




Fig. 3. (c) Silica versus major oxides for garnet mica schists.




Fig. 3. (d) Silica versus trace elements.

REEs have been shown to be mobile to some extent during metamorphism (McLennan, 1982).

The REE content of a metamorphic rock directly mimics that of the protolith. This assumption has formed the basis for a great number of studies in which the evolution of the protolith was examined in detail without specifically testing for the immobility of the REEs (Grauch, 1989). In this context, the issue of REE migration during metamorphism

in close proximity to the PAAS. The higher LREE/HREE ratio indicates a high degree of fractionation during the metamorphic stage as no such chemical heterogeneity has been observed in the protolith. The steep LREE and gentle HREE with negative europium anomaly are depicted by the Patharkhola schists. The similarity in the values of schists with PAAS is interpreted to be stemming from the chemical homogeneity of the protolith which has changed a very little during the metamorphism (Likhanov, 2008). These aforementioned features are typical of post Archean clay shales which are caused by the occurrence of erosional products of upper continental crust material (an average of 23 Australian shales of Post Archean age), (Taylor & McLennan, 1985) which imparts the presence of erosional products of sedimentary rocks in




Fig. 3. (e) REE normalised plot (Taylor & McLennan, 1985).

has been long debated between proponents of isochemical metamorphism and those of metasomatism. The only certainty is that REEs are mobile in certain circumstances (Vocke et al., 1987) and immobile in others (Rolland et al., 2003). Despite some progress in this direction, the nature of these circumstances has rarely been determined. Further research on a typical lithologically and chemically distinct types of sample suites may help our understanding of REE contents in metamorphic rocks, because they may contain a record of REE mobility or immobility (Grauch, 1989).

The analyzed REE data for the schists were normalized to Post Archean Australian Shales (PAAS), normalized values are from Taylor & McLennan (1985).

They exhibits slightly LREE depleted patterns ( $\text{LaN}/\text{SmN} = 0.7$ ,  $\text{LaN}/\text{YbN} = 0.64$ ) with flat HREE patterns ( $\text{SmN}/\text{YbN} = 0.89$ ). Concentration of lithophile elements (Rb, Cs, Ba and Sr) along with HFSE (Y, Nb and Zr) have shown values





Fig. 4. (a) Th-Sc-Zr/10 triangular plot (Bhatia, 1983); (b)  $\text{SiO}_2$  versus  $\text{K}_2\text{O}/\text{Al}_2\text{O}_3$  (Bhatia, 1983).

the detritus accompanied by the decrease in the Eu content during the sedimentation process of residual plagioclase (Taylor & McLennan, 1985).

## 5 Tectonic implications

Studies have traditionally shown that geochemistry plays a crucial role as sensitive indicator in determining the provenance of sedimentary and metasedimentary rocks and also to constrain the tectonic setting in which they were deposited (e.g., Bhatia, 1983; Bhatia & Crook, 1986; Roser & Korsch, 1986; Madukwe et al., 2015; Grzelj et al., 2017). Trace elements such as Co, Sc, Ni, Zr, Th, La and others are used for tectonic environment discrimination due to their fractionation and low mobility in sedimentary environments. Bhatia, (1983) proposed ternary plots of Sc-Th-Zr/10 and Sc-La-Th to ascertain the tectonic settings. The process of collisional tectonics and deformation is formed during the mechanism of orogeny or plate convergence leading to the formation of continental arc or active continental margin settings. These depositional environments forming in these regions are usually underlain by thick and elevated continental crust, (Bhatia & Crook, 1986). Geochemical composition of schists plotted in these ternary plots, falls in the active continental settings (Fig. 4a). Conversely, the plot of  $\text{SiO}_2$  versus  $\text{K}_2\text{O}/\text{Na}_2\text{O}$ , shows that schists of Patharkhola have been formed in the active continental margin (Fig. 4b).

## 6 Conclusion

The garnet mica schists of the Patharkhola area forms a part of Dudhatoli Almora group in Kumaun Lesser Himalaya. Petrographically, the garnet mica schist is dominantly composed of garnet, biotite, muscovite, chlorite, plagioclase and a subordinate amount of quartz. It is observed that the prekinematic garnet wrapped round first generation biotite and muscovite. Second generation Biotite-II and Muscovite-II occur as cross-cutting relation with the Biotite-I and Muscovite-I.

The geochemical data reflects that Patharkhola schists are alkali rich with silica-saturated nature. These investigated schist exhibits slightly LREE depleted patterns ( $\text{LaN}/\text{SmN} = 0.7$ ,  $\text{LaN}/\text{YbN} = 0.64$ ) with flat HREE patterns ( $\text{SmN}/\text{YbN} = 0.89$ ). The  $\log(\text{Fe}_2\text{O}_3/\text{K}_2\text{O})$  versus  $\log(\text{SiO}_2/\text{Al}_2\text{O}_3)$  plot shows that the protolith of schists is mainly shale-dominated, representing nearby source sediments formed in arc-type environment. Their geochemical signature further suggests that the schists of Patharkhola have formed in active continental margins. Thus we considered that the Lesser Himalayan rocks have formed in arc-type active continental setting where the sediments were received from both from the arc and continental crust.

## Acknowledgements

The authors thank the Head, Department of Applied Geology, Doctor Harisingh Gour Vishwavidyalaya, Sagar (M.P.) and the Department of Science and Technology, New Delhi, India, for providing facilities as including PURSE-Phase-II for conducting present research work. Thanks to Igor Petrik from Slovak Academy of Sciences and one anonymous reviewer for critical review for the improvement of manuscript.

## References

- AUDEN, J. B., 1937: Structure of the Himalaya in Garhwal. *Records Geol. Surv. of India*, 71, 407–433.
- BALI, R. & AGARWAL, K. K., 1999: Microstructures of mylonites in the Almora Crystalline Zone, Kumaun Lesser Himalaya. *Gondwana Res. Group Mem.*, 6, 111–116.
- BARKER, F., 1979: Trondhjemite: definition, environment and hypotheses of origin. In: *Developments in petrology*, Elsevier, 6, 1–12.
- BHANOT, V. B., SINGH, V. P., KANSAL, A. K. & THAKUR, V. C., 1977: Early Proterozoic Rb-Sr whole-rock age for central crystalline gneiss of Higher Himalaya, Kumaun. *Geol. Soc. India*, 18, 2, 90–91.
- BHARGAVA, O. N., FRANK, W. & BERTLE, R., 2011: Late Cambrian deformation in the Lesser Himalaya. *J. Asian Earth Sci.*, 40, 1, 201–212.
- BHATIA, M. R., 1983: Plate tectonics and geochemical composition of sandstones. *J. Geol.*, 91, 6, 611–627.
- BHATIA, M. R. & CROOK, K. A., 1986: Trace element characteristics of graywackes and tectonic setting discrimination of sedimentary basins. *Contr. Mineral. Petrology*, 92, 2, 181–193.
- BLEVIN, P., 2003: Metallogenesis of granitic rocks. In The Ishihara Symposium. *Granites Assoc. Metallogen.*, 14, 5–8.
- CHAPPELL, B. W., 1974: Two contrasting granite types. *Pacific Geol.*, 8, 173–174.
- CULLERS, R. L., CHAUDHURI, S., ARNOLD, B., LEE, M. & WOLF, W. J., 1975: REE distributions in clay minerals and in the clay-sized fraction of the lower Permian Havensville and Eskridge shales of Kansas and Oklahoma. *Geochim. cosmochim. Acta*, 39, 1691–1703.
- DE LA ROCHE, H. D., LETERRIER, J. T., GRANDCLAUDE, P. & MARCHAL, M., 1980: A classification of volcanic and plutonic rocks using R1R2-diagram and major-element analyses – its relationships with current nomenclature. *Chem. Geol.*, 29, 1–4, 183–210.
- FLEET, A. J., 1984: Aqueous and sedimentary geochemistry of the rare earth elements. In: *Developments in geochemistry*, Elsevier, 2, 343–373.
- FROST, B. R., BARNES, C. G., COLLINS, W. J., ARCUS, R. J., ELLIS, D. J. & FROST, C. D., 2001: A geochemical classification for granitic rocks. *J. Petrology*, 42, 11, 2033–2048.
- GAIROLA, V. K. & JOSHI, M., 1978: Structure of a part of Dudhatoli Almora crystalline thrust sheet around Thalisain, district Pauri Garhwal – Structure d'une partie de la nappe charriée cristalline de Dudhatoli-Almora dans la région de Thalisain, district de Pauri Garhwal Himal. *Geol.*, 8, 1, 379–398.

- GRIZELJ, A., PEH, Z., TIBLJAŠ, D., KOVACIĆ, M. & KUREČIĆ, T., 2017: Mineralogical and geochemical characteristics of Miocene pelitic sedimentary rocks from the south-western part of the Pannonian Basin System (Croatia): Implications for provenance studies. *Geosci. Frontiers*, 8, 1, 65–80.
- HUMPHRIS, S. E., 1984: The mobility of the rare earth elements in the crust. In: *Developments in Geochemistry*, Elsevier, 2, 317–342.
- IRVINE, T. N. & BARAGAR, W. R. A., 1971: A guide to the chemical classification of the common volcanic rocks. *Canad. J. Earth Sci.*, 8, 5, 523–548.
- ISLAM, R., AHMAD, T. & KHANNA, P. P., 2005: An overview on the granitoids of the NW Himalaya. *Himalayan Geol.*, 26, 1, 49–60.
- JOSHI, M. & TIWARI, A. N., 2007: Folded metamorphic reaction isograds in the Almora Nappe, Kumaun Lesser Himalaya: Field evidence and tectonic implications. *Neu. Jb. Geol. Paläont., Abh.*, 215–225.
- JOSHI, M. & TIWARI, A. N., 2009: Structural events and metamorphic consequences in Almora Nappe, during Himalayan collision tectonics. *J. Asian Earth Sci.*, 34, 3, 326–335.
- JOSHI, M. & TIWARI, A. N., 2004: Quartz C-axes and metastable phases in the metamorphic rocks of Almora Nappe: evidence of pre-Himalayan signatures. *Current Sci.*, 87, 995–998.
- JOSHI, G., AGARWAL, A., AGARWAL, K. K., SRIVASTAVA, S. & VALDIVIA, L. A., 2017: Microstructures and strain variation: Evidence of multiple splays in the North Almora Thrust Zone, Kumaun Lesser Himalaya, Uttarakhand, India. *Tectonophysics*, 694, 239–248.
- JOSHI, M., KUMAR, A., GHOSH, P., DAS, B. P. & DEVI, P. M., 2019: North Almora Fault: a crucial missing link in the strike slip tectonics of western Himalaya. *J. Asian Earth Sci.*, 172, 249–263.
- KARIMPOUR, M. H., 1983: Application of trace elements and isotopes for discriminating between porphyry molybdenum, copper, and tin systems and the implications for predicting the grade. *Glob. Tecton. Metallog.*, 29–36.
- MACKENZIE, F. T. & GARRELS, R. M., 1971: Evolution of sedimentary rocks. New York, Norton.
- MADUKWE, H. Y., OBASI, R. A., FAKOLADE, O. R. & BASSEY, C. E., 2015: Provenance, tectonic setting and source-area weathering of the coastal plain sediments, South West, Nigeria. *Sci. Res. J.*, 3, II, 20–31.
- MCLENNAN, S. M., 1982: On the geochemical evolution of sedimentary rocks. *Chem. Geol.*, 37, 3–4, 35–350.
- MEHDI, S. H., KUMAR, G. & PRAKASH, G., 1972: Tectonic evolution of eastern Kumaun Himalaya: a new approach. *Himalayan Geol.*, 2, 481–501.
- MOLNAR, P. & TAPPONNIER, P., 1975: Cenozoic Tectonics of Asia: Effects of a Continental Collision: Features of recent continental tectonics in Asia can be interpreted as results of the India-Eurasia collision. *Science*, 189, 4201, 419–426.
- MUECKE, G. K., PRIDE, C. & SARKAR, P., 1979: Rare-earth element geochemistry of regional metamorphic rocks. *Physics Chemistry Earth*, 11, 449–464.
- MUKHERJEE, S., 2005: Channel flow, ductile extrusion and exhumation of lower-mid crust in continental collision zones. *Current Sci.*, 89, 3, 435–436.
- MUKHERJEE, S., 2007: Geodynamics, deformation and mathematical analysis of metamorphic belts of the NW Himalaya. Unpublished Ph. D. thesis. Indian Institute of Technology Roorkee, 1–267.
- MUKHERJEE, S., 2013: Channel flow extrusion model to constrain dynamic viscosity and Prandtl number of the Higher Himalayan Shear Zone. *Int. J. Earth Sci.*, 102, 1811–1835.
- MUKHERJEE, S., CAROSI, R., VAN DER BEEK, P., MUKHERJEE, B. K. & ROBINSON, D. M., 2015: Tectonics of the Himalaya: an introduction. *Spec. Publ.*, 412, 1, 1–3.
- MUKHERJEE, S., PUNEKAR, J. N., MAHADANI, T. & MUKHERJEE, R., 2015: Intrafolial folds: review and examples from the western Indian Higher Himalaya. Ductile shear zones: From micro-to macro-scales, 182–205.
- NAKATA, T., 1989: Active faults of the Himalaya of India and Nepal. *Geol. Soc. Amer., Spec. Pap.*, 232, 1, 243–264.
- O'CONNOR, J. T., 1965: A classification for quartz-rich igneous rocks based on feldspar ratios. *US Geol. Surv., Profess. Pap.*, 525B, B79–B84.
- PEARCE, J. A., HARRIS, N. B. & TINDLE, A. G., 1984: Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *J. Petrology*, 25, 4, 956–983.
- RANA, H. & THOMAS, H., 2018: Geology of the Patharkhola area, Almora District, Uttarakhand (India): with special reference to the lithology and field relation. *Bull. Depart. Geol.*, 1–6. DOI: <https://doi.org/10.3126/bdg.v20i0.20716>.
- RANA, H., THOMAS, H., SONI, A. & SHUKLA, S., 2023: Petrochemistry of Phyllites from Patharkhola, Lesser Kumaun Himalaya with Reference to Tectonic Implications. *Earth Planet.*, 2, 2, 1–9. DOI: <https://doi.org/10.36956/eps.v2i2.842>.
- RANA, H. & THOMAS, H., 2023: Structural, Micro-Structural and Thermal Characterizations of Natural Garnet of Regions of Patharkhola from the State of Uttarakhand of India. *Bull. Pure Appl. Sci. Geol.*, 1.
- RAO, Y. B., CHETTY, T. R. K., JANARDHAN, A. S. & GOPALAN, K., 1996: Sm-Nd and Rb-Sr ages and PT history of the Archean Sittampundi and Bhavani layered meta-anorthosite complexes in Cauvery shear zone, South India: evidence for Neoproterozoic reworking of Archean crust. *Contr. Mineral. Petrology*, 125, 2–3, 237.
- ROSER, B. P. & KORSCH, R. J., 1986: Determination of tectonic setting of sandstone-mudstone suites using SiO<sub>2</sub> content and K<sub>2</sub>O/Na<sub>2</sub>O ratio. *J. Geol.*, 94, 5, 635–650.
- ROSSI, J. N., TOSELLI, A. J., BASEI, M. A., SIAL, A. N. & BAEZ, M., 2011: Geochemical indicators of metalliferous fertility in the Carboniferous San Blas pluton, Sierra de Velasco, Argentina. *Geol. Soc. London, Spec. Publ.*, 350, 1, 175–186.
- SHAND, S. J., 1943: Classic A/CNK vs A/NK Plot for Discriminating Metaluminous, Peraluminous and Peralkaline Compositions. New York, Hafner Publishing.
- THAKUR, V. C., 2004: Active tectonics of Himalayan frontal thrust and seismic hazard to Ganga Plain. *Current Sci.*, 1554–1560.
- THAKUR, V. C., JAYANGONDAPERUMAL, R. & MALIK, M. A., 2010: Redefining Medlicott – Wadia's main boundary fault from Jhelum to Yamuna: An active fault strand of the main boundary thrust in northwest Himalaya. *Tectonophysics*, 489, 1–4, 29–42.

- THOMAS, T. & THOMAS, H., 1992: Fold flattening and strain studies in a part of Almora Crystalline Zone, around Tamadhaun Kumaun Himalaya. *Indian Mining and Engineering Jour.*, 5–7.
- TAYLOR, S. R. & MCLENNAN, S. M., The Continental Crust: its Composition and Evolution. *Oxford, Blackwell Scientific Publication*, pp. 312.
- VALDIYA, K. S., 1980: Stratigraphic scheme of the sedimentary units of the Kumaun Lesser Himalaya. In: Valdiya, K. S., Bhatia, S. B. (eds.): Stratigraphy and Correlation of the Lesser Himalayan Formation. *Delhi, Hindustan Pub. Corp.*, 7–48.
- VALDIYA, K., 1980b: Geology of Kumaun lesser Himalaya. *W. I. H. G.*, 291–295.
- VALDIYA, K. S., 1988: Tectonics and evolution of the central sector of the Himalaya. *Phil. Trans. Roy. Soc. London. Ser. A, Mathem. Phys. Sci.*, 326, 1589, 151–175.
- WEBB, A. A. G., YIN, A., HARRISON, T. M., CÉLÉRIER, J., GEHRELS, G. E., MANNING, C. E. & GROVE, M., 2011: Cenozoic tectonic history of the Himachal Himalaya (northwestern India) and its constraints on the formation mechanism of the Himalayan orogen. *Geosphere*, 7, 4, 1013–1061.
- WEBB, A. A. G., 2013: Preliminary balanced palinspastic reconstruction of Cenozoic deformation across the Himachal Himalaya (northwestern India). *Geosphere*, 9, 3, 572–587.
- YIN, A., 2006: Cenozoic tectonic evolution of the Himalayan orogen as constrained by along-strike variation of structural geometry, exhumation history, and foreland sedimentation. *EarthSci. Rev.*, 76, 1–2, 1–131.

## Geochemické charakteristiky a tektonická interpretácia granatických svorov oblasti Patharkhola v Kumaunských Malých Himalájach, Uttarakhandske Himaláje, India

Pohorie Himaláje tvorí oblúkový horský pás pokrývajúci celú hraničnú zónu severnej Indie. Je typovým príkladom zrážky medzi indickou a áziskou litosférickou platňou v období pred približne 55 miliónmi rokmi (Mukherjee, 2015; Yin, 2006). Štruktúre, stratigrafii a tektonike územia Kumaun v Malých Himalájach sa venujú viaceré práce (Joshi a Tiwari, 2009; Joshi et al., 2017; Rana a Thomas, 2018). V oblasti Kumaun je sekvencia Malých Himalájí oddelená hlavnou centrálnou prešmykovou zónou (*Main Central Thrust*; MCT) od Vyšších Himalájí na severe. Hlavný hraničný prešmyk (*Main Boundary Thrust*; MBT) oddeluje túto sekvenciu od oblasti Siwalikov na juhu (Thakur et al., 2010). Horninové súbory skupiny Dudhatoli (Rana a Thomas, 2018) vystupujú v oblasti Patharkhola.

Vnútorná bridličnatá séria, označovaná aj ako tektonická zóna príkrovu metamorfovaných a kryštalických hornín, tvorí v Malých Himalájach výraznú štruktúrnu jednotku s polyfázovou deformáciou a metamorfózou. Kryštalíkum masívu Dudhatoli vystupuje od lokalít Garhwal na ZSZ po Kumaun na VSV. Kumar a Agarwal (1975) zahrnuli Mandhali a formácie Chandpur a Nagthat, rovnako ako kryštalíkum Dudhatoli-Almora, do novovytvorenej skupiny Dudhatoli, v ktorej kryštalíkum Dudhatoli-Almora reprezentuje najvyšší horizont. Horniny skupiny Dudhatoli sa považovali za prekambrické (Kumar a Agarwal, 1975). V Kumaunských Himalájach boli ruly jednotky Munsiari skupiny Almora datované na  $1\ 830 \pm 200$  mil. r. (Bhanot et al., 1977). Islam et al. (2005) rozčlenili proterozoické veky granitoidov Malých Himalájí do dvoch skupín: 2 200 – 1 800 mil. r. a 1 400 – 1 200 mil. r. Rana a Thomas (2023) študovali termálny a štruktúrny charakter granátov z granatických

svorov v bridliciach z lokality Patharkhola. Zistilo sa, že sú bohaté na Mg, čo dokladá ich vyššieplotný rozklad.

Študovaný terén vykazuje polyfázovú deformáciu a metamorfózu (Joshi a Tiwari, 2009). Je to dobre pozorovateľné v centrálnej časti almorského príkrovu. V Kumaunských Malých Himalájach sa realizoval rozsiahly výskum peliticých rúl a granitov a v menšom rozsahu aj nízkostupňových bridlíc a fylitov (Phukan et al., 2018; Das et al., 2019). Rana et al. (2023) štúdiom patharkholských fylitov dospeli k záveru, že fylity s vysokým obsahom Al a obohatené stopovými prvkami vznikli na aktívnom kontinentálnom okraji.

Oblast' okolo lokality Patharkhola s vystupovaním hornín Malých Kumaunských Himalájí je ohraničená zemepisnou dĺžkou  $79^\circ 09' V$  až  $79^\circ 17' 56'' V$  a zemepisnou šírkou  $29^\circ 47' 42'' S$  až  $29^\circ 56' 69'' S$ . Má rozlohu približne  $120\ km^2$  (obr. 1a). Vystupujúcimi horninami sú ruly, bridlice a fylity (obr. 1b). Článok je zameraný na geochemický charakter a petrogenézu bridlíc.

Pri tektonickej interpretácii je dôležitý geochemický výskum sedimentárnych a metasedimentárnych hornín, ktorý dokáže citlivu preukázať ich pôvod a tektonické prostredie ich vzniku. Analyzované údaje o REE bridlíc boli normalizované podľa *Archean Australian Shales* (PAAS) s normalizovanými hodnotami podľa Taylora a Mclennana (1985).

Vykazujú mierne ochudobnené LREE ( $\text{LaN}/\text{SmN} = 0,7$ ,  $\text{LaN}/\text{YbN} = 0,64$ ) s plochým priebehom HREE ( $\text{SmN}/\text{YbN} = 0,89$ ). Koncentrácia litofilných prvkov (Rb, Cs, Ba a Sr) spolu s HFSE (Y, Nb a Zr) má hodnoty v tesnej blízkosti PAAS. Vyšší pomer LREE/HREE označuje vysoký stupeň frakcionácie počas metamorfnej

fázy, keďže takáto chemická heterogenita sa nezistila v protolite. Patharkholské bridlice vykazujú strmé LREE a mierne HREE s negatívnou európskou anomáliou. Podobnosť hodnôt bridlíc s PAAS je interpretovaná ako kontaminácia z primárneho prostredia, pretože protolit sa málo zmenil počas metamorfózy (Likhanov, 2008). Uvedené znaky sú typické pre postarchaickú hlinu bridlíc s výskytom erozívnych produktov z vrchnej kontinentálnej kôry (priemer z 23 austrálskych bridlíc postarchaického veku). Prítomnosť erozívnych produktov sedimentárnych hornín v detrite je sprevádzaný poklesom obsahu Eu počas procesu sedimentácie zvyškov plagioklas (Taylor a McLennan, 1985).

Štúdie mnohých autorov preukázali, že geochémia je citlivý indikátor určujúci provenienciu sedimentárnych a metasedimentárnych hornín a tektonické prostredie, v ktorom sa nachádzali (napr. Bhatia, 1983; Bhatia a Crook, 1986; Roser a Korsch, 1986; Madukwe et al., 2015; Grzelj et al., 2017). Na určenie tektonického prostredia sa využívajú predovšetkým stopové prvky ako Co, Sc, Ni, Zr, Th, La a ďalšie. Na určenie tektonického prostredia Bhatia (1983) navrhol ternárne diagramy Sc-Th-Zr/10 a Sc-La-Th. Prejavy kolíznej tektoniky a deformácie sú v rámci orogenézy produktom konvergencie litosférických platní vedúcej k vzniku kontinentálneho oblúka alebo prostredia aktívneho kontinentálneho okraja. Depozičné prostredia v takýchto regiónoch sú zvyčajne podostielané hrubou a vyklenutou kontinentálnou kôrou (Bhatia a Crook, 1986). Geochemické zloženie bridlíc zo-

brazených v ternárnych diagramoch spadá do prostredia kontinentálnej kolízie (obr. 4a). Naopak, diagram  $\text{SiO}_2$  vs.  $\text{K}_2\text{O}/\text{Na}_2\text{O}$  indikuje, že bridlice z Patharkholu sa vytvorili na aktívnom kontinentálnom okraji (obr. 4b).

V záverečnom zhrnutí je možné konštatovať, že granatické svory v oblasti Patharkhola, ktoré sú súčasťou skupiny Dudhatoli-Almora v Kumaunských Malých Himalájach, petrograficky pozostávajú z granátu, biotitu, muskovitu, chloritu, plagioklasu a druhoradého množstva kremeňa. Zistilo sa, že predkinematický granát je obalený prvou generáciou biotitu a muskovitu. Druhá generácia biotitu-II a muskovitu-II sa vygenerovala priečne v mikrofraktúrach biotitu-I a muskovitu-I. Geochemické údaje dokladajú, že svory sú bohaté na alkálie a nasýtený  $\text{SiO}_2$ . Vykazujú mierne ochudobnené LREE ( $\text{LaN/SmN} = 0,7$ ,  $\text{LaN/YbN} = 0,64$ ) s plochým priebehom HREE ( $\text{SmN/YbN} = 0,89$ ). Diagram  $\log(\text{Fe}_2\text{O}_3/\text{K}_2\text{O})$  vs.  $\log(\text{SiO}_2/\text{Al}_2\text{O}_3)$  dokladá, že protolitom svorov bola predovšetkým bridlica reprezentujúca zdrojové sedimenty z aktívneho kontinentálneho okraja. To indikuje, že horniny z Malých Himalájí sa vytvorili v prostredí kontinentálneho oblúka, kde detrit sedimentov pochádzal z tohto oblúka, ako aj z kontinentálnej kôry.

Doručené / Received: 26. 9. 2023  
Prijaté na publikovanie / Accepted: 21. 12. 2023