


48/1/2016
ISSN 1338-3523
ISSN 0369-2086

Mineralia Slovaca



Štátny geologický ústav Dionýza Štúra Bratislava



Predseda vydavateľskej rady – Chairman of Editorial Board

BRANISLAV ŽEC

Štátny geologický ústav Dionýza Štúra Bratislava

Vedecký redaktor – Scientific Editor

ZOLTÁN NÉMETH

Štátny geologický ústav Dionýza Štúra
Regionálne centrum Košice
Jesenského 8, 040 01 Košice
zoltan.nemeth@geology.sk

REDAKČNÁ RADA – EDITORIAL BOARD

IGOR BROSKA, Geologický ústav SAV Bratislava

KLEMENT FORDINÁL, Štátny geologický ústav D. Štúra Bratislava

VLASTA JÁNOVÁ, Ministerstvo životného prostredia SR Bratislava

MARTIN KOVÁČIK, Štátny geologický ústav D. Štúra Bratislava

PETER MALIK, Štátny geologický ústav D. Štúra Bratislava

JOZEF MICHALÍK, Geologický ústav SAV Bratislava

LUBOMÍR PETRO, Štátny geologický ústav D. Štúra Košice

DUŠAN PLAŠIENKA, Prírodovedecká fakulta UK Bratislava

MARIÁN PUTIŠ, Prírodovedecká fakulta UK Bratislava

IGOR SLANINKA, Štátny geologický ústav D. Štúra Bratislava

JÁN SOTÁK, Geologický ústav SAV Banská Bystrica

LADISLAV ŠIMON, Štátny geologický ústav D. Štúra Bratislava

PAVEL UHER, Prírodovedecká fakulta UK Bratislava

REDAKCIA – EDITORIAL STAFF

Vedúci oddelenia vydavateľstva ŠGÚDŠ a propagácie – Head of the Department of ŠGÚDŠ Publishers and Promotion

LADISLAV MARTINSKÝ

ladislav.martinsky@geology.sk

Jazykoví redaktori – Lingual editors

Janka Hrtusová – Zoltán Németh

janka.hrtusova@geology.sk

Sadzb a korektúry – Typeset and proofreading

Grafická úprava a technické spracovanie – DTP processing

Mária Dryjová

maria.dryjova@geology.sk

Slávka Žideková

slavka.zidekova@geology.sk

Mineralia Slovaca (Web ISSN 1338-3523, ISSN 0369-2086), EV 3534/09, vychádza dvakrát ročne. Vydavateľ a tlač: Štátny geologický ústav Dionýza Štúra, Mlynská dolina 1, 817 04 Bratislava, IČO 31 753 604. Dátum vydania čísla 48/1/2016: jún 2016. Predplatné v roku 2016 vrátane DPH, poštovného a balného pre jednotlivcov 16,50 €, pre členov SGS a geologických asociácií 15,40 €, pre organizácie v SR 22,00 €. Cena jednotlivého čísla je 5,50 € vrátane DPH. Časopis možno objednať v redakcii a v knižnici regionálneho centra v Košiciach.

Adresa redakcie: Štátny geologický ústav D. Štúra – RC Košice (Mineralia Slovaca), Jesenského 8, 040 01 Košice. Telefón: 055/625 00 46, 055/625 00 43; fax: 055/625 00 44, e-mail: mineralia.slovaca@geology.sk, e-mail knižnica: secretary.ke@geology.sk

Mineralia Slovaca (Web ISSN 1338-3523, ISSN 0369-2086) is published twice a year by the State Geological Institute of Dionýz Štúr Bratislava, Slovak Republic. The date of issuing of the number 48/1/2016: June 2016. Subscription for 2016 calendar year: 33.00 € (Europe), 45.00 (besides Europe), including VAT, postage and packing cost. Price of one number is 5.50 € including VAT. Claims for nonreceipt of any issue will be filled gratis. Order send to address of the Editorial office: Štátny geologický ústav D. Štúra – RC Košice (Library), Jesenského 8, SK-040 01 Košice, Slovak Republic. Phone: +421/55/625 00 46, +421/55/625 00 43; fax: +421/55/625 00 44, e-mail: mineralia.slovaca@geology.sk, library: secretary.ke@geology.sk

© Štátny geologický ústav Dionýza Štúra Bratislava

PŮVODNÉ ČLÁNKY – ORIGINAL PAPERS

Fričovský, B., Tometz, L., & Fendek, M.

Geothermometry techniques in reservoir temperature estimation and conceptual site models construction: Principles, methods and application for the Bešeňová elevation hydrogeothermal structure, Slovakia

Využitie geotermometrie pri odhade rezervoárovej teploty a konštrukcii koncepčných modelov:

Princípy, metódy a aplikácie v podmienkach hlbokého rezervoáru hydrogeotermálnej štruktúry bešeňovskej elevácie, Slovensko 1

Zlinská, A.

Terciárna mikrofauna z hlbokých vrtov v železovskej priehlbine (Dunajská panva)

Tertiary microfauna in deep wells of the Železovce depression (Danube Basin) 61

Cvečková, V., Fajčíková, K., Jurkovič, L., Stehlíková, B., & Rapant, S..

Chemické zloženie podzemnej vody a úmrtnosť na kardiovaskulárne ochorenia v Slovenskej republike

Chemical composition of groundwater and mortality for cardiovascular diseases

in the Slovak Republic.....83

OBÁLKA: Pohľad z vrchu Veľký Choč (1 611 m n. m.) na hydrogeotermálne pole Liptovskej kotliny (horný obrázok). Dobre viditeľné sú vodné nádrže Liptovská Mara a Bešeňová s rovnomennou obcou. Šípka znázorňuje stredovú polohu produkčnej oblasti hydrogeotermálnej štruktúry Bešeňovej. Približne v oblasti medzi oboma vodnými priehradami prechádza štruktúra po zlomových líniiach do susednej hydrogeotermálnej štruktúry depresie Liptovskej Mary. Na južnej, protihľanej strane kotliny vystupujú severné svahy hydrogeologického masívu Nízkyh Tatier, ktoré sa podieľajú na tvorbe infiltračnej oblasti termálnych vôd. Pohľad smerom na JV. Foto: Z. Németh.

Staršia snímka (obrázok dolu) termálneho parku Bešeňová (Gino Paradise Bešeňová). Park sa využíva celoročne. Termálna voda z hlbokého rezervoáru v strednotriasových karbonátoch krížňanského príkrovu je exploatovaná dvomi vrtmi, ZGL-1 a FGTB-1, na zabezpečenie celoročnej prevádzky aj pri extrémne nízkej vonkajšej teplote. Tepelný výkon oboch vrtov je dostačujúci na bezpečné vykurovanie obytných priestorov a atrakcií na lokalite. Foto: J. Olejník.

Schéma tektonického rozčlenenia hlbokého rezervoáru v strednotriasových karbonátoch krížňanského príkrovu s vyznačenými zlomovými líniami. Šípky zobrazujú priestorovú orientáciu prúdenia podzemnej, respektíve termálnej vody a zároveň variabilitu prísunu tepla do rezervoárovej polohy na jeho báze. Práve nerovnomerný prísun tepla z podložia sa považuje za jednu z možných príčin vzniku izolovaných konvekčných buniek v rezervoári, ktoré vo výraznej miere ovplyvňujú termodynamické parametre a prúdenie termálnej vody. Problematikou sa zaoberá úvodný článok tohto čísla. Autor: B. Fričovský.

COVER: A view on a hydrogeothermal field of the Liptov Basin from the Veľký Choč hill (1 611 m a.s.l.). A dominant feature is a dam of the Liptovská Mara (left) and Bešeňová (right). An arrow points to a centre of productive part of the hydrogeothermal structure of the Bešeňová. At between both dams the structure terminates along fault systems with a neighboring Liptovská Mara depression hydrogeothermal structure. A hydrogeological Massif of the Nízke Tatry Mts. exposes to the south, contributing on formation of recharge zone for the structure. The view to the SE. Photo: Z. Németh.

A picturesque view on the thermal park Bešeňová (Gino Paradise Bešeňová) operated during an entire year. Thermal water produced from the deep reservoir associated with Mid Triassic carbonates of the Krížna Nappe system is exploited by two wells, the ZGL-1 and FGTB-1, securing a continuous operation even at extreme outdoor temperatures. A thermal output is high enough to provide an individual space heating of the resort and heating the pools at a site. Photo: J. Olejník.

A scheme of tectonic dissection of the deep reservoir of the Bešeňová elevation hosted within Mid Triassic carbonates, with delineated major tectonic lines. Arrows visualize a concept of groundwater and thermal water filtration and unbalanced heat influx from a base respectively. The reservoir base unbalanced overheating is believed as one of agents triggering formation of insulated convective cells in the reservoir, with substantial impact on thermal waters movement and thermodynamic properties. The problem is emphasized in the initial article of this volume. Author: B. Fričovský

Geothermometry techniques in reservoir temperature estimation and conceptual site models construction: Principles, methods and application for the Bešeňová elevation hydrogeothermal structure, Slovakia

BRANISLAV FRIČOVSKÝ¹, LADISLAV TOMETZ² and MARIÁN FENDEK³

¹Department of Hydrogeology and Geothermal Energy, State Geological Institute of Dionýz Štúr, Mlynská dolina 1, SK-817 04 Bratislava, branislav.fricovsky@geology.sk


²Institute of Geosciences, Faculty of Mining, Ecology, Process Control and Geotechnologies, Technical University of Košice, Park Komenského 15, SK-042 00 Košice, ladislav.tometz@tuke.sk

³Department of Hydrogeology, Faculty of Natural Sciences, Comenius University in Bratislava, Mlynská dolina, SK-842 15 Bratislava; fendek@fns.uniba.sk

(Received 15. 2. 2016, accepted for publishing 19. 4. 2016)

Abstract The Bešeňová elevation hydrogeothermal structure has been analysed applying the ion-solute (silica, cation) and multicomponent equilibrium geothermometry techniques. Instead of a simple model of infiltration, transition and discharge, the multiple mixing stages have been proven, whether referring to the infiltration zone, transition – accumulation area, or discharge zone, where most of a focus has been oriented to the Vyšné Sliače area. Only multicomponent equilibrium and silica geothermometry based on chalcedony conductive cooling models appears adequate in the reservoir temperature estimation. Application of the cation geothermometers suffers of the deep reservoir thermal waters immaturity, driven by the separate convection cells due to unconform aquifer base overheating. In the Vyšné Sliače area, warm springs are related to both reservoir zones by lateral leaking (driven by density, hydraulic and pressure gradient), later diluted at a contact with the shallow and cold groundwater. Qualitative use of geothermometers proves its adequacy in construction of conceptual site models for low enthalpy geothermal systems. Quantitative application, however, remains limited in essence.

Key words: Bešeňová elevation, conceptual site models, multicomponent equilibrium geothermometry, solute geothermometry, thermodynamics



Highlights

- Complex geothermometry applied for a first in conditions of the W. Carpathians in CSM construction
- Separate convection cells form within the deep reservoir
- Immaturity of thermal waters due to convection and carbonate environment
- Minor (SW) discharge zone in the Sliače area
- Dry CO₂ fluxes in the Sliače area due degassing during upwelling

FOREWORD

In the past decades, geothermometry gained conclusive credits worldwide since first of them have been developed (e.g. Fournier and Truesdell, 1971), whether for reservoir temperature assessment, reservoir dynamics identification (mixing, boiling, dilution, leakage, etc.), field development perspectives, geothermal history matching and reservoir production response monitoring, or conceptual site models (CSM) construction. Both aspects of geothermometry application, quantitative temperature estimation or qualitative

analysis, require comprehensive understanding of theory beyond, local thermodynamics and basic site concepts.

Essentially, geothermometers either formulate a degree of deep particular constituents concentration conservation over a reservoir fluid history, or determine a magnitude of equilibration of fluids to the dissolved mineral assemblage or wallrock.

A skepticism towards geothermometry in conditions of the Western Carpathian stems from misunderstood convention of its use for temperature estimation only, subsequently considering it undue, as being derived from high

enthalpy fields. The same disbelief, however, contravenes the principal philosophy of geothermometry application.

In contrast, this contribution explains principles and application conditions of geothermometry at tough background, with consequent demonstration of its qualitative and quantitative analysis in CSM construction for the Bešeňová elevation hydrogeothermal structure. We also provide an extended review on most important and well established publications, explaining this topic in details and in extension that our contribution could not cover.

THE BEŠEŇOVÁ ELEVATION STRUCTURE

The Bešeňová structure represents pre-Paleogene basement elevation (Maďar, 1997), trending N-S within the western part of the Liptov Basin (Fig. 1a), limited tectonically to its surroundings. Limitation to neighboring structures of the Liptovská Mara depression (E) and the Ivachnová depression (W) is formed by the N-S trending Vlachy – Lubel'a – Vrch Hája and Bešeňová fault respectively. Fault swarms of NW–SE to NE–SW trend terminate the structure at the Nízke Tatry Mts. on the south. To the north, the Choč fault provides a tectonic contact of the Bešeňová elevation structure with the Chočské vrchy Mts. (Remšík et al., 1998).

Pioneering research on local mineral waters assumed the springs nearby the town of Bešeňová as being associated with the deep and thermal water reservoirs due to the travertine domes formation, and connection of the groundwater basin (the Liptov Basin) with the massif of Nízke Tatry Mts. (Franko et al., 1974), later extending the conception to delineate both areas as the discharge and recharge zones (Franko, 1984). Meanwhile, temperature and density measurements of dry CO₂ fluxes in the Vyšné Sliače area assigned them a petrogenic origin (Zbořil et al., 1972). The analysis of springs and artesian wells at the southern margin of the basin has provided an argument to assume several infiltration channels at different depths towards the basin. Soon Zembjak et al. (1986) concluded that the infiltration and recharge propagation differentiation realizes at open fault systems at southern periphery of the basin, whilst local springs are not just of barrier, but the tectonic origin as well.

Geothermal prospection on thermal waters began in the late 1970s, installing a 450 m deep BEH-1 well. Thermal waters, found in the Choč Nappe (CHN) Mid Triassic carbonates and a secondary permeable top of the Križna Nappe (KNA) Jurassic–Mid Cretaceous variegated pelite limestones, have recorded the transient Ca-Mg-HCO₃-SO₄ type of waters at $T_{\text{btm}} \approx 35$ °C, unconform to the host rock environment, arising an impression of their origination in deeper, Križna Nappe (KNA) Mid Triassic carbonate complexes (Franko et al., 1979). Completion of the ZGL-1 Bešeňová in 1987 documented deep reservoir thermal waters with $T_{\text{wh}} \approx 61.5$ °C, compared to the formation

temperature of $T_{\text{btm}} \approx 68 - 71$ °C. Sampling revealed three geochemical facies at different positions: carbonatogene (Ca-Mg-HCO₃) to transient (Ca-Mg-SO₄-HCO₃) and sulphatogene (Ca-SO₄) geochemical types (Fendek et al., 1988). Later, the FBe-1 well (400 m) replacing the BEH-1 well after its failure, has produced the shallow reservoir waters of Ca-Mg-HCO₃ to Ca-Mg-HCO₃-SO₄ type waters at $T_{\text{wh}} \approx 25$ °C (Vandrová et al., 2009). Problems with ZGL-1 well called for a completion of the FGTB-1 Bešeňová well (1833 m) in 2011 that proved sulphatogene Ca-SO₄ to transient Ca-Mg-SO₄-HCO₃ character of deep reservoir waters at $T_{\text{btm}} = 69$ °C and $T_{\text{wh}} = 66$ °C (Vandrová et al., 2011). Hydraulic connection between both reservoirs and shallow reservoir extension to the Borové Formation (IWCP) has been concluded continuously (Remšík et al., 1998; Vandrová et al., 2009).

Since the beginning, thermal waters have been found as oversaturated with respect to calcite (e.g. Fendek et al., 1988). Geothermometry application resulted in marginal constations on immaturity to the Na/K system (Remšík et al., 1993), silica geothermometry inadequacy and uncertain validity of temperature – aqueous SO₄ concentration relation (Bodiš and Borosová, 1996). Only $\delta\text{O}_{\text{SO}_4} - \delta\text{O}_{\text{H}_2\text{O}}$ oxygen isotope geothermometry application (Remšík et al., 1998) yielded some success in temperature estimation ($T_{\text{est}} = 40 - 52$ °C).

Geology and tectonics

The Liptov Basin is a typical Tertiary intramountain depression of the Western Carpathians. Its recent arrangement into N–S elevated or depressed blocks has originated by the combination of (1) the pre-Paleogene relief segmentation and onset of the first karstification period (Mesozoic mass reduction; Činčura and Köhler, 1995); (2) Late Eocene–Oligocene synsedimentary tectonic activity regionally limiting deposition rate of the IWCP (Inner Western Carpathian Paleogene) sediments (Gross et al., 1979); (3) Neogene basin dissection and IWCP mass reduction along N–S, NW–SE and SW–NE faults (Němec and Bartková, 1987); and (4) Late Neogene–Early Quaternary rejuvenation of W–E faults inverting a relief to balance uplift tendencies of surrounding massifs (Jurewicz, 2005).

Review on geology

Vertical arrangement of the structure corresponds to its geotectonic evolution and recent tectonic position. *Devonian–Mid Carboniferous* magmatites and metamorphites form Tatricum Crystalline (TCR) bedrock beneath the *Early Triassic–Mid Cretaceous* para-autochthonous Tatricum Envelope Unit (TEU) and allochthonous Križna Nappe (KNA) system or *Mid Triassic* Choč Nappe (CHN) tectonic slag respectively; all carbonates domina-




Fig. 1. The Bešeňová elevation definition: Geographic position (a), simplified vertical profile (b), major faults delineation (c) and 3D tectonic scheme with deep reservoir body (Križna Nappe Mid Triassic carbonates) detail (d).

ted. Above, a *Mid Eocene–Latest Oligocene* succession of the Inner Western Carpathian Paleogene (IWCP) forms a siliciclastics dominated envelope (Gross et al., 1980), reduced in thickness centrewards the system (Gross et al., 1980). Atop, continental *Quaternary* sedimentary cover forms some meters thick cover (Remšík et al., 1998).

The geothermal system encompasses two reservoirs (Remšík et al., 2005). The *deep reservoir* associates with KNA Mid Triassic carbonates (limestones, dolomites, transient varieties), forming a solid body through an entire system, being segmented into several blocks in depths between 660–3 200 m and sealed by the Early Triassic (base) and Late Triassic (top) rich-in-evaporites siliciclastics (quartzites, pelite shales) dominated aquicludes. The *shallow reservoir* is recognized in the CHN Mid Triassic dolomites, hydraulically connected to conglomerates, breccia and detritic carbonates of the Borové Formation (IWCP). Atop, typical claystones-dominated (Huty Formation) and flysch-dominated (Zuberec Formation) horizons, both of IWCP group, form regional top insulator, while Jurassic–Mid Cretaceous KNA duplexed pelitic and organogene carbonates (Fig. 1b–d) represent a central aquitard (Fendek et al., 1988). Even the occurrence of thermal waters associated with the Mid Triassic TEU carbonates can not be excluded, though it has neither been proven yet.

Role of tectonics

Intersection of W–E and SW–NE faults at SE periphery allows the groundwater to enter the deep circulation regimes from the massif (Franko et al., 1979). Obviously, NW–SE and NE–SW faults towards the central part allow either further submerging of thermal waters already in the circulation, or hydraulic communication between CHN and KNA reservoirs (Fendek et al., 1988; Fričovský et al., 2015). The Choč–Ružbachy fault of trending W–E and located on the north, however, hydraulically separates thermal waters of the basin to waters documented in the Lúčky area (Fendek et al., 2013). To the south, contact of W–E, N–S and NW–SE faults respectively triggers upwelling of springs from moderate to deep circulation from the Nízke Tatry Mts. (e.g. Franko, 2002a; Remšík et al., 2005), or provide upwelling conditions for thermal waters from the basin (Fričovský and Tometz, 2013). It is fair then to assume those faults are open down to the KNA profile. If we accept some possibility on closed, TEU carbonates hosted structure, faults must be healed at a horizon of KNA Early Triassic, as any of mixing models gave a valuable indices on invasion of TEU thermal waters into the KNA carbonates (Fričovský and Tometz, 2013). Unlike, the Choč–Ružbachy fault zone is healed secondary, providing a lithological trap for hydraulic communication between structures of Bešeňová and Lúčky (Fendek et al., 2013; Franko, 2002b).

Review on geochemistry

The system has been concluded as open hydrogeological structure. Lateral inflow realizes along N–S faults from the Liptovská Mara depression, whilst runoff tendency is rather of intraformation character to the Ivachnová depression (Remšík et al., 1998). The infiltration zone is located towards the Nízke Tatry Mts. (Fendek et al., 1988) – Kľačianka valley area (Fričovský et al., 2015). Even there is a general agreement on discharge zone in the central part of the structure nearby Bešeňová, this paper discusses existence of second, minor, southwest located discharge zone of the Sliače area.

The BEH-1 and FBe-1 wells represent the shallow reservoir, along with ZGL-1 and FGTB-1 screening the deep aquifer. Both encompass the accumulation – transition zone. In the Vyšné Sliače area, waters are analysed by the Sliače springs and the VŠH-1 well. The recharge zone chemistry is approximated for shallow circulation by the Demänová springs and deep circulation by the wells in the Demänová valley area. Analyses are listed in Appendix C.

Review on facies analysis

Groundwaters are atmospheric in origin and petrogenic in chemistry. The shallow reservoir (SRTW) contains transient $\text{Ca}(\text{Mg})\text{-HCO}_3\text{-SO}_4$ to $\text{Ca-SO}_4\text{-HCO}_3$ and carbonatogene Ca-Mg-HCO_3 waters (Franko et al., 1979; Vandrová et al., 2009). Carbonatogene character is conformal to wallrock, defining this facies primary. Origin of sulphates is, however, expected for the KNA Early or Late Triassic formations (e.g. Fendek et al., 1988). These must then be secondary, resultant to deep reservoir waters evasion and mixing. Deep reservoir (DRTE) water samples at a bottom record sulphatogene (Ca-SO_4) and transient ($\text{Ca-SO}_4\text{-HCO}_3$) character, whilst at a top, carbonatogene Ca-Mg-HCO_3 to transient $\text{Ca-Mg-HCO}_3\text{-SO}_4$ facies were documented (e.g. Remšík et al., 1998). Variation in SO_4 content is due to vertical reservoir dynamics (Fričovský and Tometz, 2013). In central discharge zone, springs record mixed $\text{Ca-Na-HCO}_3\text{-SO}_4$ type, owing to Quaternary hosted groundwaters with the SRTW evasions. Waters of VSH-1 are of transient character. Mixed to hydrosilicatogene chemistry of springs (Franko et al., 1979) is, probably, due to the mixing of VSH-1 facies with the groundwater hosted in the Quaternary or Paleogene at shallow altitudes (Fričovský et al., 2015). Infiltrated waters from the N-slopes of the Nízke Tatry Mts. attain both, carbonatogene type for the shallow and transient type for deep transition.

Review on mixing models application

After Tshc diagram (Giggenbach, 1988), waters of accumulation zone are classified as immature and acid

to soda-springs (Fig. 2a), whilst both, discharge and recharge zone samples plot within the peripheral fluids only (Fričovský and Tometz, 2013). At Schoeller's plot (e.g. Marini, 2004), the thermal waters of deep reservoir are relatively stable in cation abundance, varying SO_4/HCO_3 ratio because of dissimilar contact with the sulphate source (Fričovský, 2014). There are no indices of adiabatic boiling (Fig. 2b) as curves do not change a shape significantly.

Some vertical dispersion is rather an implication of instable reservoir accommodation and, possibly, separate convection cells formation. Use of Gibb's plots (Gibbs, 1970) reveals a general lack in Na^+ for all facies. All waters may be classified as of continuous recharge insensitive to precipitation intensity (Fričovský and Tometz, 2013), but subjected to polystageous mixing because of facies similarities in between (Fig. 2c).




Fig. 2. Review on classification and mixing diagrams: The maturity Tshc plot (a), the Schoeller's plot for deep reservoir samples (b), and the Gibb's plot (c).

BACKGROUND

Understanding the nature and characteristics of the targeted system is key for its successful exploration, development and (sustainable) utilization. This is best achieved by conceptual model development (Axelsson, 2009), visualizing or interpreting relation between its particular components of physical and chemical provenience (Cummings,

2009). In other words, the conceptual model displays in-situ situation and relations generally.

Interpretation and validation of results, obtained by geothermometry application, necessitates as much knowledge on the processes the sample could underwent as possible. There is, thus, a need to understand its conceptual history, under which we understand every chemical, physical, dynamic and thermodynamic process, the fluid

or sample experienced since it entered a hydrogeothermal structure until it has been discharged or sampled. Improving the knowledge on conceptual model helps understanding the conceptual history and vice versa. The conceptual history must then be accepted as an inherent integrant of the CSM. Both must be considered and known, to some extension, before geothermometry application.

Geothermometers – classification and limitations

Geochemical composition of fluids projects saturation states for mineral assemblage(s) in solution (Browne, 1978), derived from a host rock environment and a conceptual history of a fluid. A chemical or isotope geothermometer is a model, based on temperature-dependent chemical reactions or isotope equilibrium fractionation of which reservoir equilibrium temperatures can be derived (Pang, 2001). These reactions are variably slow in re-equilibration at cooler temperatures and after effective phase separation, thus the equilibration temperature is believed as “printed” in a form of concentrations or ionic ratios (Karingithi, 2009).

Geothermometers consider individual constituents or their combination, of which three groups are distinguished conventionally (D’Amore and Arnórsson, 2000):

- solute geothermometers (silica and cation geothermometers)
- gas geothermometers (master gasses, noble gasses); and
- isotope geothermometers

Recent interest aims at thermodynamic equilibrium analysis, considering a fluid as a complex of solutes at various saturation states, introducing a multicomponent geothermometry (e.g. Spycher et al., 2011).

Yet the critical consideration in reservoir temperature estimation is a conservation of deep fluid chemical composition at a sampling point, assuming (Fournier et al., 1974):

- chemical reactions in a reservoir are temperature-dependent
- chemical and isotopic reactions do not change reservoir fluid chemistry significantly
- solids for equilibration reactions are sufficiently abundant in a host formation
- water–rock equilibration takes place at reservoir temperature
- the geofluid suffers little or no re-equilibration, mixing or dilution in upflow

Fluids meet all the criteria unlikely as they always lose heat by conductive cooling or boiling. Conductive cooling does not necessarily change a composition, but may change the saturation states of solutes, modifying the composition by precipitation or dissolution optionally. Adiabatic boiling (flashing) changes chemical composition immediately, subsequent to CO₂ evasion, variation in residual concentrations or pH (Marini, 2004). Isotopic

composition of flashed water depends upon whether the steam fraction remained with the boiling water until it separated in once, or there had been a continuous steam removal from a fluid, relying upon a number of flashing stages (Truesdell et al., 1977).

Amount of a heat lost during conductive cooling is proportional to vertical distance traveled and inversely proportional to the flow rate or discharge. To some extension, duration of conductive cooling results in some pH variation, which, in a consequence, buffers dissolution–precipitation processes (Arnórsson, 2000a). The heat lost by adiabatic boiling is proportional to boiling rate and propagation. Chemical composition of a vapor in its wet-steam or dry-steam form always experiences modification by condensation or air attainment (Marini, 2004).

The actual reservoir fluid composition is then representative at no-boiling, no-mixing conditions prior sampling, otherwise parental composition must be reconstructed. Regardless of conformity in results, geothermometry bears substantial information for CSM construction.

SILICA GEOTHERMOMETRY

The silica geothermometry reflects solubilities of its polymorphs (quartz, chalcedony, amorphous silica, LT/HT opal), being not too sensitive to hydrostatic pressure changes and salt additions below 300 °C (Fournier and Potter, 1982). Temperature is then the factor to control silica thermodynamics in fluids, forth the use of temperature – aqueous silica concentration correlations becomes valuable for the most of geothermal systems worldwide.

Silica solubility and aqueous SiO₂ control

Silica solubility

First studies on silica solubility (Fig. 4) were carried out by Morey et al. (1962), defining an essential reaction to control the process (Eq. 1):



for which the equilibrium constant (Eq. 2) is given (Fournier and Rowe, 1966):

$$K_{eq} = \frac{[\text{H}_4\text{SiO}_4]}{[\text{SiO}_{2(s)}][\text{H}_2\text{O}]^2} \quad (\text{Eq. 2})$$

which reduces to (Eq. 3) as activity of pure phases equals 1

$$K_{eq} = [\text{H}_4\text{SiO}_4] \quad (\text{Eq. 3})$$




Fig. 3. Basic conceptual histories of geofluids.

The silica solubility decreases dramatically with adiabatic or conductive cooling (Fig. 4). Silica may, however, precipitate before reaching surface or the wellhead, forth sampling concentrations can not be assumed representative in some occasions.

The quartz is the most stable silica polymorph at $T = 180\text{--}330\text{ }^{\circ}\text{C}$ (Walther and Helgeson, 1977; Fournier and Potter, 1982), controlling aqueous SiO_2 concentration in high enthalpy fields. At $T = 120\text{--}180\text{ }^{\circ}\text{C}$ (low enthalpy or cooled/boiled high enthalpy fluids), the chalcedony becomes metastable and dissolved SiO_2 depends on conceptual history of the water. Below $T = 120\text{ }^{\circ}\text{C}$ (most low enthalpy or dramatically boiled thermal waters) it is likely the solubility of chalcedony is in control (Fournier, 1989). Activity of other polymorphs is rather rare in geothermal systems, even they may express higher solubility (Fournier, 1977).

Silica regime

If the fluid boils before sampling, the $\text{SiO}_{2\text{aq}}$ concentration increases in the residual liquid proportionally to the amount of evaded steam, overestimating calculations (Fournier, 1989). This is the reason the chalcedony (Arnórsson et al., 1983) and quartz (Fournier, 1973) geothermometers were modified to correct for aqueous silica increase (see app. A).

Onset of silica precipitation is observed in conjunction to boiling, causing $\text{SiO}_{2\text{aq}}$ concentrations at sampling points lower than those in parental reservoir that counteracts effect of relative silica concentration increase. Analogously, geothermometers, which do or do not involve boiling correction, may yield good results

(D'Amore and Arnórsson, 2000). According to that, both should be used simultaneously and considered as probably providing a fairly reasonable range of temperature estimates.

An effect of silica solubility increase with pH is pronounced at above $\text{pH} = 7.5$ and $T > 175\text{ }^{\circ}\text{C}$. Correction upon silica enrichment is necessary if a solution became alkaline prior dissolving additional silica in a response to increase in pH (Fournier, 1981) as a consequence of boiling. Then, the aqueous SiO_2 is possible to originate from hydrolytic dissociation of alkali silicates, instead of hydrolysis of wallrock silica polymorphs (Fournier, 1977).

High enthalpy fluids record most likely $\text{pH} < 7.5$, being resultant to buffering the hydrogen ions by hydrolytic reactions involving silicates (Fournier, 1977). At lower temperatures ($T_{\text{res}} < 100\text{ }^{\circ}\text{C}$), thermal waters may have high CO_2 content and low pH because of, for example, dissolution of calcite.

Dilution decreases the silica content consequent to the onset of silica precipitation due to cooling. Because of fast response to reservoir dynamics, most of the equilibrium re-attainment with the silica causes the estimate to give a temperature of re-equilibration, otherwise parental temperatures tend to be underestimated (Fournier, 1980a).

Quartz geothermometers work best for $T_{\text{res}} > 150\text{ }^{\circ}\text{C}$. For plutonic systems the usual validity range for chalcedony geothermometry is $T_{\text{res}} = 120\text{--}180\text{ }^{\circ}\text{C}$, whilst the lower limit drops in carbonate, basaltic and sedimentary reservoirs to $T_{\text{res}} = 60\text{--}180\text{ }^{\circ}\text{C}$. Amorphous silica and cristobalite geothermometry is trendless, depending on their control only.

The majority of silica geothermometers (see Appendix A) applies to solely cooling springs at sub-boiling temperatures or

borehole samples (Arnórsson et al., 1983; Fournier, 1973, 1977, 1991; Truesdell and Fournier, 1975), assuming reservoir equilibration conservation, thus vertical velocity of upwelling fluids is higher than silica re-equilibration and any dilution. These are labeled as “conductive cooling”. Silica geothermometers with correction for boiling “adiabatic boiling geothermometers” (Fournier, 1973; Arnórsson et al., 1983) compensate the steam loss and residual SiO_2 . Modifications obtain precious results for boiling springs of high discharge ($> 2 \text{ kg}\cdot\text{s}^{-1}$) or samples that undergo reservoir boiling prior sampling (Fournier, 1977).

Application: silica solubility analysis

One of accented arguments against silica geothermometry applicability in carbonate reservoirs is a “sufficient source” objection. Indeed, the silica is definitely not excessive in pure carbonates. Referring to local conditions, some minor silica sources must be accounted:

- siliciclastic IWCP formations in early downflow/late upflow stages
- pelitic Jurassic–Mid Cretaceous carbonates in early downflow/late upflow stages
- siliciclastics and impure detritic carbonates in late downflow/early upflow stages
- siliciclastic intercalations and silica impurities in reservoir carbonates
- siliciclastic and detritic carbonates hosted shallow reservoir

Analytical approach in controlling silica polymorph analysis is an alternative, if the primary or alteration mineralogy data are missing. This is grounded by sit-and-cluster principle along solubility functions (Fig. 4), correlated to temperature and reservoir homogeneity criteria.

With a great confidence, the chalcedony is a controlling polymorph on aqueous SiO_2 in local conditions. At relatively low wellhead ($T_{\text{wh}} = 51\text{--}62^\circ\text{C}$) and deep borehole ($T_{\text{res}} = 61\text{--}69^\circ\text{C}$) temperature differences, the SiO_2 oscillates considerably ($c\text{SiO}_2 = 35\text{--}80 \text{ mg}\cdot\text{l}^{-1}$) in deep reservoir, lacking any trend with the temperature. Under expected reservoir homogeneity where carbonates dominate over siliciclastic intercalations and thin sandstone beds, the variation records unstable dissociation of silicates (Fig. 4). Because it is not possible for quartz to take over a control in the shallow reservoir, clustering along a chalcedony and quartz metastability line at higher temperature is rather a record of two different facies. The metastability is rather attributed to silica shortage in the deeper parts of the system (BEH-1; $c\text{SiO}_2 = 10\text{--}17 \text{ mg}\cdot\text{l}^{-1}$, $T_{\text{wh}} = 32\text{--}34^\circ\text{C}$) due to its deposition after evasion from DRTE. The upper parts of the reservoir (FBe-1; $c\text{SiO}_2 = 27\text{--}30 \text{ mg}\cdot\text{l}^{-1}$, $T_{\text{wh}} = 25\text{--}27^\circ\text{C}$) do not suffer a lack in source as connected to the IWCP formations. By a contrast, samples of the discharge zone plot well onto chalcedony solubility line, definitely answering a controlling silica polymorph question.

Application: silica saturation analysis (deep reservoir)

The deep reservoir is found a key uncertainty in understanding the complex silica stability at a site. We inspected opal (low temperature), chalcedony (moderate temperature) and quartz (high temperature) to review SiO_2 thermodynamics within. Plotting saturation indices against extended temperature range (Fig. 5) eliminated opal of any influence on $\text{SiO}_{2\text{aq}}$ concentration in thermal fluids as it remains undersaturated within the entire interval.




Fig. 4. Silica solubility analysis with (a) and against (b) the inverse of temperature. Modified after Morey et al. (1962).

The chalcedony equilibrates at $T = 56\text{--}94\text{ }^{\circ}\text{C}$ (Fig. 5) completely, with only two samples not attaining saturation below $T = 90\text{ }^{\circ}\text{C}$. Most of samples intercepts $SI = 0$ at $T = 65\text{--}70\text{ }^{\circ}\text{C}$. Irregularity in equilibration dispersion is either a consequence of the deficiency in silica source or unstable contact with the silica source (and hydrolysis reactions) within a deep reservoir. Thus this may be considered an indirect evidence on vertical reservoir filtration.

On a face of saturation curves distribution, the quartz may be of influence onto silica concentration in the water. This is because 14 out of 20 samples record equilibration at $T = 80\text{--}100\text{ }^{\circ}\text{C}$. In fact, reservoir temperature of over $90\text{ }^{\circ}\text{C}$ is an optimistic assumption, reducing $SI = 0$ interception to 4 samples only.

A clear trend of the increase in SiO_2 with p-T, depth, residential time and distance from an infiltration zone is missing (Fig. 4). The lack accounts to (1) the reservoir (intraformation) dynamics (mixing, vertical filtration); (2) insufficient source of silica for hydrolysis; and (3) silica origin by the water-rock cation exchange reactions. The controlling polymorph in the shallow filtration regime within the recharge zone may be opal, proceeding towards chalcedony metastability with the increase in p-T conditions during submersion. Both, SiO_2 concentration and saturation variation is resultant either to vertical mass flux in reservoir and insufficient available silica. Reservoir chalcedony saturation (Fig. 5) turns it available for cation exchange reactions with the wallrock, with minor deposition only (if any). Thus the deep SRTW facies are deficient in silica, explaining their affinity to plot onto quartz solubility line. By a contrast, the VSH-1 well records no lack in silica, even it should be implicit under given geological, p-T and initial DRTE conditions, evidencing other silica source to balance its loss. The compensation must then be due to mixing with SiO_2 rich source (SRTW evasions, groundwater hosted in IWCP or Jr/Cr-2 succession), or proportional to decrease in filtration velocity of DRTE facies towards discharge zone.

Reservoir dynamics and processes

Silica – enthalpy mixing model (XSih)

The silica – enthalpy mixing/geoindicator model (Fournier and Truesdell, 1974) is developed to determine reservoir temperatures regardless of dilution and steam loss. Use of enthalpy instead of temperature takes an advantage in the heat (energy) conservation in a reactive system. Involved silica solubility function must correspond to the polymorph controlling $\text{SiO}_{2\text{aq}}$ concentration.

For geothermal waters at $T_{\text{res}} = 50\text{--}340\text{ }^{\circ}\text{C}$ (Fournier and Potter, 1982) proposed polynomial functions to determine the enthalpy H_L for liquid (Eq. 4) or vapor (Eq. 5) component, including liquid enthalpy for low temperature systems (Eq. 6) of $T_{\text{res}} < 50\text{ }^{\circ}\text{C}$. Unknown dissolved silica (cSi) may be derived with (Eq. 7) below fluid enthalpy of $H < 1\,670\text{ J}\cdot\text{kg}^{-1}$. Besides (Eq. 4 to Eq. 6) enthalpies can be read from steam tables or calculated (e.g. using REFPROP). Silica concentrations may be taken from analyses.

$$H_L = 418.84 + 10.286T - 0.05092T^2 + 2.6309 \cdot 10^{-4}T^3 - 6.9303 \cdot 10^{-7}T^4 + 7.4566 \cdot 10^{-10}T^5 - 1209.8T^{-1} + 11.99T^{-2} - 353.76\log T \quad (\text{Eq. 4})$$

$$H_L = 2\,035 - 5.0499T + 0.057399T^2 - 3.0426 \cdot 10^{-4}T^3 + 7.909510 \cdot 10^{-7}T^4 - 8.6968 \cdot 10^{-10}T^5 + 1\,342.4T^{-1} - 13.298T^{-2} + 396.29\log T \quad (\text{Eq. 5})$$

$$H_L = 4.1868 \cdot T \quad (\text{Eq. 6})$$

$$cSi = -3.5532 + 0.146H_L - 4.927 \cdot 10^{-4}H_L^2 + 1.2305 \cdot 10^{-6}H_L^3 - 4.9421 \cdot 10^{-10}H_L^4 \quad (\text{Eq. 7})$$

Procedure for non-boiling CSMs is relatively straight-forward. The point (A) represents a non-thermal, cold groundwater at its silica concentration and enthalpy, the fluid is expected to mix with. The point (B) is stated for the sampled thermal spring depart of (sub) boiling conditions. Straight line (A–B) drawn intercepts the silica (quartz or chalcedony) solubility curve at point (C). Vertical line onto X-axis gives the enthalpy of resultant member, forth read from steam tables at pure water vapor pressure. The same model gives an opportunity to check for possible reservoir boiling. Usual assumption expects the fluid to boil under atmospheric pressure that equals the enthalpy of $H = 419\text{ kJ}\cdot\text{kg}^{-1}$. From that, vertical line along Y-axis intercepts the mixing (A–C) line at point (D). Contouring the horizontal line towards maximum steam loss curve gives the initial (intraformation) enthalpy of hot-water component (E), projected onto X-axis.

Reasonable result represents a reservoir temperature of parental thermal water before it has been mixed, so the horizontal line onto Y-axis gives initial reservoir silica concentration (Fournier, 1980a). For either case (no-boiling and boiling model), overestimation of derived initial reservoir temperatures is a hint on boiling prior mixing or silica addition after mixing. If lines do not intercept, it may be a consequence of cooling without silica deposition. Derived initial temperatures below the expected range give a vital sign on silica deposition prior isenthalpic mixing or steam loss (Truesdell and Fournier, 1977).

Application: shallow reservoir and discharge zone silica-enthalpy analysis

In this model, we accepted archive data (Appendix C) for DRTE because of no evidence on boiling/mixing correction necessity. Meanwhile, (Eq. 6) and (Eq. 7) were used to determine thermodynamic silica concentrations and temperatures in other zones, assuming isenthalpic mixing without boiling.

For DRTE, we calculated the reservoir enthalpy by REFPROP for pure water phase at atmospheric pressure, obtaining $H_{\text{res}} = 336\text{--}355\text{ kJ}\cdot\text{kg}^{-1}$. For the SRTW, we pre-

ferred silica concentrations computed (Eq. 7) of $c\text{SiO}_2 = 11\text{--}14 \text{ mg}\cdot\text{l}^{-1}$ under a mean enthalpy of $H_{\text{res}} = 105\text{--}142 \text{ kJ}\cdot\text{kg}^{-1}$ to minimize effect of silica hydrolysis or deposition. Mixing line (Fig. 6) between both facies is given by their median enthalpy and concentration values. This yields the enthalpy of $H_{\text{res}} = 580 \text{ kJ}\cdot\text{kg}^{-1}$, equal to overestimated $T = 138^\circ\text{C}$. For the shallow reservoir (cold-water component), the temperature excision implies silica deposition in deeper parts of the horizon. Cooler subfacies against DRTE individually give the final temperature of $T = 116^\circ\text{C}$, thus the rate of silica imbalance decreases proportionally, as they compensate loss in SiO_2 by its hydrolysis from IWCP formations.

Similar principles were adapted to carry an analysis at the discharge zone facies. Samples of cold Sliače springs represent cold water component at sampled concentrations. VSH-1 represent a thermal water mixture, with concentrations ($c\text{SiO}_2 = 12\text{--}13 \text{ mg}\cdot\text{l}^{-1}$) and enthalpy ($H_{\text{res}} = 163\text{--}176 \text{ kJ}\cdot\text{kg}^{-1}$) substituted according to (Eq. 6) and (Eq. 7). After solubility line extrapolation, springs samples sit onto. Drawing a mixing line between springs and VSH-1 (Fig. 6) overestimates temperature ($T = 70^\circ\text{C}$) and enthalpy ($H = 295 \text{ kJ}\cdot\text{kg}^{-1}$). VSH-1 waters must be enriched with SiO_2 prior mixing with S-CGW. If mixing line is plotted between deep reservoir representative sample and VSH-1, the overestimate counts $H = 555 \text{ kJ}\cdot\text{kg}^{-1}$ (or $T = 133^\circ\text{C}$). The yield is excessive because VSH-1 samples evidently loose some silica compared to deep reservoir they are expected to be relative with (Fig. 6).

Silica – carbonate mixing model (XSiCo)

Geofluids worldwide recorded strong correlation between total carbonate (ΣCO_3) and temperature, with most




Fig. 6. Silica – enthalpy mixing model. Modified after Fournier and Truesdell (1974).

of it partitioned into CO_2 (Arnórsson et al., 1983). Amount of aqueous silica in a sample relates to solubility of its polymorphs at fixed temperatures (Fournier et al., 1974). Most of silica speciates into H_4SiO_4 . The relation between CO_2 as total carbon, Ct , and SiO_2 in aqueous solution is, however, not linear. A model curve is obtained by plotting relative CO_2 (Eq. 8) and SiO_2 (Eq. 9) concentration at fixed T in Kelvin (Arnórsson, 1985).

$$\text{SiO}_2 = -15.433 - \frac{151.6}{T} - (2.977 \cdot 10^{-6} T^2) + 5.464(\log T) \quad (\text{Eq. 8})$$

$$\text{CO}_2 = -1.09 - \frac{3894.55}{T} + 2.532(\log T) \quad (\text{Eq. 9})$$

Resultant concentrations are in logs of molality. Silica concentration may be optionally modified by the chalcedony geothermometers. Boiled waters plot above the line or tend to cross the curve. Non-boiling samples are usually positioned below the line. Unmixed fluids plot onto the curve or cluster close around. If fluid boils before sampling, the aqueous CO_2 level is depleted proportionally to produced vapor, whilst aqueous SiO_2 increases in residual fluid. Therefore, the CO_2/SiO_2 ratio drops. Dilution of the fluid without boiling retains CO_2 in a liquid phase causing CO_2/SiO_2 ratios high relative to parental conditions (Arnórsson, 1985). Overestimated temperatures are resultant to silica removal (deposition) after mixing due to a conductive cooling. Temperatures below reasonable range for parental (reservoir) fluid may be a sign of partial degassing after earlier dilution or mixing respectively (Nicholson, 2012).

Relation between CO_2 and SiO_2 aids solute geothermometry application. Besides interpretation of positions, the model is used as geindicator, extending a trend line from provenienced populations to intercept the curve at point A. Drawing a line onto X-axis gives CO_2 content in a parental fluid. The initial silica concentration is obtained by projecting A onto Y-axis. This is correct only for a case where the supportive hints on mixing or boiling are present, as well as for a case where quartz controls the aqueous silica in the water. Temperature is given by substitution of yielded parental SiO_2 concentration into quartz geothermometer (Eq. 10) by Fournier and Potter (1982). In other cases, this procedure should not be used to estimate reservoir temperature, or the plot must be modified for the chalcedony geothermometers. Inserted concentrations (c) are in $\text{mg}\cdot\text{kg}^{-1}$.

$$T(K) = -53.5 + 0.11236c - 0.5559 \cdot 10^{-4} c^2 + 0.1772 \cdot 10^{-7} c^3 + 88.39 \log c \quad (\text{Eq. 10})$$

Application: discharge zone silica – total carbonate analysis

We used (Eq. 8–9) to draw the cross equilibrium function. The conversion of initial ($\text{mg}\cdot\text{l}^{-1}$) into mass (ppm)

concentrations was carried in WATCH v.2.4 package that accounts for H_2CO_3 dissociation under given pH and temperature. The Sliache springs (S-CGW) are identified as cold water component ($cSiO_2 = 56\text{--}128$ ppm; $cCO_2 = 79\text{--}1\,830$ ppm). Thermal waters of VSH-1 well represent a mixture ($cSiO_2 = 44\text{--}53$ ppm; $cCO_2 = 230\text{--}867$ ppm). Samples of ZGL and FGTB-1 wells stand for thermal end-member ($cSiO_2 = 56\text{--}128$ ppm; $cCO_2 = 79\text{--}1\,830$ ppm). All facies were upscaled to representative values by median concentrations.

Facies plot in the non-boiling region, disproving any boiling in the structure. Deep reservoir samples organize along a logarithmic trend line (Fig. 7), similar to the equilibrium function by Arnórsson (1985), secluded deep within the non-boiling region because of the general lack in silica typical for carbonates. This evidences DRTE stability to dilution. Samples of VSH-1 must be distinguished into D-LTC (discharge zone – low total carbonate) and D-HTC (discharge zone – high total carbonate), as they plot discontinuously.

A mixing line is drawn between S-CGW and D-LTC, fairly striking DRTE median, implying some influential facial relation (Fig. 7). Extended mixing line intercepts the equilibrium curve at 521 ppm of SiO_2 . Later, we substituted the inferred concentration into chalcedony conductive geothermometers (see App. A), obtaining a temperature of $T = 190\text{--}205$ °C, which is definitely an overestimate. Apparently, D-LTC loses CO_2 and SiO_2 , compared to parental conditions by silica consumption into hydrolysis reactions and minor precipitation.

To explain the D-HTC population, we introduce a shallow reservoir (SRTW) facies ($cSiO_2 = 16\text{--}48$ ppm; $cCO_2 = 110\text{--}1\,090$ ppm), that is, however, not a geothermal end-member. A line connecting the S-CGW and D-HTC facies intercepts a region of median shallow reservoir, but does never cut the XSiCo model (Fig. 7). Then, if there




Fig. 7. Silica – total carbonate model. Equilibrium curve after Arnórsson et al. (1983).

is some facies relation between the shallow reservoir and thermal waters, the D-HTC may originate by mixing with DRTE (increase in CO_2 and SiO_2), prior dilution by S-CGW (D-HTC $\rightarrow$ D-LTC), which would drop CO_2 instantaneously because of cooling.

Silica – enthalpy (no boiling) fraction model

Sampling and analysing geothermal springs is, by far, one of the easiest and cost-effective approaches in the early stages of geothermal field exploration. Warm springs or two-phase springs represent solid indices on underground geothermal reservoir and possible water evasion suffering some mixing with the shallow groundwater (early case) or degassing (latter case). Samples experiencing either of these, however, yield erroneous results (not only) for silica geothermometers application, except of those designed to correct estimates for boiling.

The fraction model based on enthalpy (Eq. 11) and mass balance (Eq. 12), developed by Fournier et al. (1974), is capable to infer the endmember temperature and a rate of parental chemistry modification by a dilution rate:

$$H_{gw} \cdot X + H_{tw} \cdot (1-X) = H_{spg} \quad (\text{Eq. 11})$$

and

$$CS_{gw} \cdot X + CS_{tw} \cdot (1-X) = CS_{spg} \quad (\text{Eq. 12}),$$

where H is the enthalpy of components ($\text{kJ} \cdot \text{kg}^{-1}$), X is the cold water fraction in the spring, and CS stands for aqueous silica concentration ($\text{mg} \cdot \text{l}^{-1}$), with following subscripts: “gw” for (cold) groundwater, “tw” for modeled thermal water component and “spg” for springs.

The enthalpy remains conserved in the system, as the thermal spring heats the groundwater. Some silica deposition due to parental water cooling is, in some extension, balanced with the aqueous SiO_2 added by the groundwater. The model assumes a spring with parental thermal water component originated in homogeneous reservoir below or at sub-boiling conditions (Sheridan et al., 1980). The parental fluid wells up towards surface and is diluted by encountering shallow and cold groundwater, which fixes a steam phase in the liquid to form the warm or hot spring (Fournier et al., 1974).

Due to enthalpy conservation, the weighting factor for cold water component “ X_H ” is given by transcription of “ X ” in (Eq. 11) towards (Eq. 13), whilst silica weighting component “ X_{CS} ” (Eq. 14) is derived from “ X ” in (Eq. 12):

$$X_H = \frac{(H_{tw} - T_{spg})}{(H_{tw} - T_{gw})} \quad (\text{Eq. 13})$$

and

$$X_{cs} = \frac{(CS_{tw} - CS_{spg})}{(CS_{tw} - CS_{gw})} \quad (\text{Eq. 14})$$

where enthalpy is in $\text{cal}\cdot\text{g}^{-1}$, T is the temperature in $^{\circ}\text{C}$ and CS is in $\text{mg}\cdot\text{l}^{-1}$.

A procedure is simple and straightforward. In a first, one must assume a reasonable temperature range for parental thermal water in natural, deep reservoir (origin) conditions. The X_H is calculated substituting variables into (Eq. 13) for every temperature, and plotted onto cross-diagram against temperature. Next step involves calculation of possible silica content (Sheridan et al., 1980) in unmixed and unboiled parental fluid per each designed temperature, here for a case a chalcedony controls the silica solubility in water (Eq. 15):

$$CS_{tw} = \frac{10^{\left[\frac{-1032}{T+273.15}\right] - 0.09}}{1665 \cdot 10^{-5}} \quad (\text{Eq. 15})$$

In following, the X_{CS} (Eq. 14) is determined for each silica content of parental fluid at each of fixed temperatures designed above, where the CS_{tw} is the only variable and CS_{spg} and CS_{gw} are constants. Next, the X_{CS} is plotted into the same graph as the X_H was in previous. Because of sensitivity to input data, a great certainty in representativity of a spring for thermal water is mandatory, requiring cross-correlation with other facial geochemical techniques.

Intersection projects a cold water fraction on the X and parental water temperature on the Y axis. Overestimate is then a hint on additional silica dissolution owing to a water-rock contact with glassy rocks or amorphous silica layers. If curves do not intercept, it is possible that the parental fluid lost steam before dilution (Fournier et al., 1974).

Application: deep reservoir fraction model analysis

Enthalpy of DRTE has been taken from REFPROP calculations, modified by a conversion factor of 0.239 to get a scale in $\text{cal}\cdot\text{g}^{-1}$. While the mixture (VSH-1) was identified at $T = 40^{\circ}\text{C}$ and $\text{cSiO}_2 = 30 \text{ mg}\cdot\text{l}^{-1}$, the cold water component attributes at $T = 9^{\circ}\text{C}$ and $\text{cSiO}_2 = 0.4 \text{ mg}\cdot\text{l}^{-1}$. The chalcedony model (Eq. 15) represents the DRTE component in the mass balance. The model was scaled for $T = 50\text{--}300^{\circ}\text{C}$ (Fig. 8).

Fraction functions (X_H , X_{CS}) intercept at $T = 130^{\circ}\text{C}$ and $X = 0.79$. The temperature is supposed to correspond to that of deep reservoir, geothermal end-member composition, however, there is a distinct overestimation. Under given undisturbed parental (DRTE) conditions, the high temperature is a result of SiO_2 additions after mixing. Another explanation is cooling of VSH-1 waters with S-CGW. Both explanations are, in fact, reasonable under local conditions. This is, however, in confidence to XSiH

and XSiCo plots. Moreover, temperature overrate makes the coldwater fraction estimate ($X = 79\%$) excessive.

Cross-check for post-mixing silica addition by use of amorphous silica solubility (Eq. 16) is recommended if resultant temperatures are overestimated. In this case, the (Eq. 16) replaces (Eq. 15) for DRTE silica upscaling at external variables and enthalpy balance constant.

$$\log CS = -\frac{731}{t + 273.15} + 4.52 \quad (\text{Eq. 16})$$




Fig. 8. Fraction model analysis.

At local conditions, the amorphous silica curve does not intercept the enthalpy function (Eq. 16). Therefore, silica addition is less probable after final mixing stage (see XSiCo model), and the temperature overestimate is rather resultant to cooling of VSH-1 wells subsequent to mixing with S-CGW and upflow.

Quantitative SiO_2 geothermometry application

Under known conception, the DRTE is expected to yield most precise results because of missing factors to modify reservoir geochemistry, although skewing by vertical reservoir filtration must be taken accounted. Chalcedony geothermometers (GSC1 – GSC4A) match best the expected deep reservoir temperature range (Fig. 9), varying the calculated temperature of $T_{\text{chalc}} = 51\text{--}98^{\circ}\text{C}$. This includes the chalcedony adiabatic boiling (Arnórsson et al., 1983) geothermometer too (GSC4A), which gives $T_{\text{chalc}} = 62\text{--}97^{\circ}\text{C}$. Yet this is a solid evidence of chalcedony control on aqueous SiO_2 concentrations in thermal waters.

The median of chalcedony conductive cooling equilibration $T_{\text{chalc-m}} = 74.8^{\circ}\text{C}$ increases to $T_{\text{chalc-m}} = 76.7^{\circ}\text{C}$ if adiabatic boiling model is included. Because of no boiling, this can be further neglected. Most of temperature estimates fall within the $T_{\text{chalc}} = 75\text{--}85^{\circ}\text{C}$ interval. In general, temperatures given by chalcedony geothermometers are

greater than those by solubility analysis (Fig. 5), with median $T_{\text{chalc}} = 67^\circ\text{C}$. In essence, the difference is because of variable sensitivity of geothermometry functions to local SiO_2 content, which is generally less than that at moderate to high volcanic fields, which they were derived from.

Variation in temperature estimates is, however, not only a function of relative sensitivity of geothermometry functions to aqueous silica level. We rather adjust the variation in estimate and absolute concentrations to unstable contact of thermal waters with a silica source, regardless it is because of its various incorporation in hydrolysis reactions or due to vertical intraformation filtration induced by unbalanced reservoir base overheating. Either way, results oppose a statement by Bodiš and Borosová (1996) on chalcedony geothermometry inadequacy. Instead of comparing results to the wellhead temperature, temperatures must be referenced to deep bottomhole (reservoir) conditions, as this is the environment where the equilibration is, due to principles of geothermometry, supposed to take place.

Both, the shallow reservoir and the discharge zone represent not an end-member but a mixture. In shallow reservoir, any silica geothermometry must simply account on two subfacies, different in the depth and silica content. The deeper subfacies (BEH-1) lacks silica, because of its deposition or consumption, after the parental, DRTE waters evade vertically the reservoir. The subfacies in shallower positions (FBe-1) are comparatively enriched with silica by its low grade (low temperature) dissociation from the IWCP and recharge. Because of numerical confidence (see Appendix D), there may be some re-equilibration with chalcedony within the formation.

For the discharge zone, inferred temperatures are lower than those expected at shallow parts of deep reservoir, where we set a parental fluid origination prior lateral leaking. Temperature estimates (Appendix D) may then point to temperature of chalcedony last equilibration during propagation towards discharge zone, or are simply skewed by multiple mixing episodes prior sampling. Because of mixture character, we would rather prefer the second scenario, and do not recommend to use quantitative silica geothermometry to either of these.




Fig. 9. Silica geothermometry results distribution: deep reservoir. See Appendix A for symbols and references.

CATION GEOTHERMOMETRY

Cation geothermometers refer to numerous ion-exchange reactions, mostly partitioning of alkalis, of various temperature dependent equilibrium constants (Fournier, 1989). This is because e.g. sulphates and carbonates adjust to changes in physical or chemical environment much faster, and so they equilibrate with the system rapidly, whilst Na^+ , K^+ or Li^{2+} silicates (i.e. tectosilicates, feldspars or micas) do so slowly even at high temperatures (Giggenbach, 1981). In other words, equilibrium preservation is more likely a case of alkali system rather than those of carbonates. Obtaining about the same results applying various cation geothermometers for a same sample at the same time is, thus, fairly rare (Fournier, 1989).

Complex mineral assemblages attain fluid-rock equilibrium rarely, as complete isochemical recrystallization of a primary rock matrix to a thermodynamically stable alteration assemblage is likely to take part in stagnant systems of an infinite age (Giggenbach, 1968; Ellis, 1970) only.

Obligatory “sufficient source” condition, especially for alkalis, fits for magmatic systems. Quantifying application of alkali geothermometers to carbonate systems often suffers from a lack of a source within reservoir and unstable water-source contact. If not for a temperature estimation, the Na-K-Mg and K-Mg-Ca thermodynamic subsystems are, at least, effective in analysis of reservoir maturity and reservoir dynamics at a site (Giggenbach, 1981).


Alteration and equilibration

A knowledge on controlling alteration assemblage is one of mandatories necessary to get a gain prior applying cation geothermometry. Most of alteration systems relate to Na-K and K-Mg alteration (thermodynamic) pairs at a base of alteration and partitioning of silicates.

Na/K geothermometry

The recalled Na/K geothermometers (Appendix B) are based on partitioning of Na^+ and K^+ between alumino-

silicates (feldspars) and water (Eq. 17), with temperature dependent equilibrium constant (Eq. 18) in control of the process (Fournier, 1979):





$$K_{eq} = \frac{[KAlSi_3O_8][Na^+]}{[NaAlSi_3O_8][K^+]} \quad (\text{Eq. 18})$$


The equilibrium constant (Eq. 18) assumes unity in reaction activities, that means reactants are pure end-members, thus activities of dissolved species are about equal to their molar concentrations (Truesdell, 1976). Then, for dilute solutions, the (Eq. 18) transforms to (Eq. 19), where Na^+ and K^+ are in desired units.

$$K_{eq} = \frac{Na}{K} \quad (\text{Eq. 19})$$

Also, for many geothermal systems, not only feldspars participate in cation exchange equilibrium reactions involving Na^+ and K^+ , but e.g. montmorillonites may do so (Eq. 20–21). Equilibrium constant for the reaction equals the Na/K ratio (Eq. 19), however, the aqueous Na/K ratio resultant to a cation exchange reaction at a given temperature controlled just by montmorillonites fairly differs to that controlled by alkali feldspars (Fournier, 1989).



After Fournier (1989b) albite may not be present at low enthalpies, and the Na/K exchange reaction is controlled by adularia – sodic clay – silica and hydrogen system (Eq. 22), equilibrated at (Eq. 23)



$$K_{eq} = \frac{[Na^+][H^+]^6}{[K^+]^7} \rightarrow \log K_{eq} = \log[Na/K] - 6\log[H] \quad (\text{Eq. 23})$$

Numerous other reactions contribute in geothermal systems at various temperatures, either involving water and equilibrium reactions with hydrogen or not. The variation in Na/K and suites controlling the Na/K ratios reasoned a range of empirically derived Na/K geothermometers (i.e. Truesdell, 1976; Tonani, 1980; Arnórsson et al., 1983; Fournier, 1979; Nieva and Nieva, 1987; Giggenbach, 1986; Giggenbach, 1988; Verma and San-

toyo, 1997; Simmons, 2013), thereupon precision of temperature estimation applying these depends on involved mineral assemblage and structural states of solids in local conditions (Fournier, 1980). Possibly, the Na/K equilibration may not be reached at $T < 100^\circ\text{C}$, as well as micas and zeolites may control the Na/K exchange, or the Na^+ and K^+ participation is rather a consequence of preferential cation leaching than exchange reactions (Fournier, 1979).

There is no universal Na/K geothermometer as one may give a correct estimation in one place and an erroneous temperature in another (Fournier, 1989), thus the use of Na/K geothermometry should be grounded by cautious mineral assemblage analysis.

The Na/K geothermometer re-equilibrates slowly in comparison to others at $T < 350^\circ\text{C}$, thereby it is less sensitive to dilution (dilutant is expected to have lower Na^+ and K^+ concentrations than the parental fluid) providing an ability to evaluate deeper parts of the high enthalpy systems (Fournier, 1980) by conservation of absolute concentrations at thermodynamic conditions the other models would fail in (Davraz, 2014). Thus it yields highest temperature estimates. Unfortunately, the highest temperature is not necessarily the correct temperature of equilibration, nor a reservoir temperature. Indeed, thermal waters may invade shallow reservoir zones prior upwelling and targeted aquifer may then be of lower temperature that the Na/K geothermometer may not recognize.

Cation geothermometers work best for reservoir fluids of $T_{res} > 150^\circ\text{C}$ (Nicholson, 2012). The necessary cation exchange equilibrium state is scarcely reached at $T_{res} < 100^\circ\text{C}$, and Na^+ and K^+ are rather resultant to Na^+ and K^+ leaching instead. Yet caution must be paid for carbonate reservoirs (rich in CO_2^- which transforms Na-silicates), and waters originated in organic sediments rich in NH_4^- (Fournier, 1980b).

Graphical tests for Na/K geothermometry application and interpretation are carried by conclusive cross charts (Fournier, 1980b; Marini, 2004) plotting a relative Na/K ratio over a reciprocal of absolute temperature, either to correlate the aqueous Na^+ and K^+ concentrations with a particular mineral assemblage and/or inspection of chemistry formation processes; or to correlate the ratios to particular equilibrations. Reservoir chemistry and dynamics are accessible for evaluation regarding position and spatial organization of samples within one or more systems (i.e. stratified reservoirs with mixing, conceptual fluid flow etc.).

Application: Na/K alteration system analysis

All facies suffer a lack of alkalis (Fig. 10), decreasing the Na/K ratio with temperature. Position of facies is far of any equilibration, whether for low (LT albite – microcline) or high (albite – adularia) enthalpy assemblage. Equilibrium pairs (Eq. 17–23) are not valid in local conditions. The inequilibrium results then in temperature overestimates (see Appendix D).





Fig. 10. Equilibrium ratios with Na-K system.

Inequilibration in deep reservoir is due to rock dissolution in control of Na/K ratio, suffering to insufficient source in carbonates and excess CO_2 which switches activities of Na-silicates to K-micas (Fig. 11). While rock dissolution controls alkali chemistry for DRTE originated subfacies in the shallow reservoir (slightly increasing Na/K due to drop in CO_2 by deposition of carbonates), subfacies recognized as recharged record preferential Na^+ and K^+ leaching from IWCP in a downflow regime. This is well correlated with records from many recharge waters (e.g. Marini). Preferential leaching inferred for VSH-1 samples is rather due to enrichment of DRTE evasions with SRTW rich in Na^+ , later enriched with Na^+ added by dilution of a mixture with cold groundwater (S-CGW) or from a wallrock acquire.


K/Mg geothermometry

Dynamic systems rarely attain overall equilibrium because of much faster reaction velocity for a thermodynamic system involving Mg^{2+} than that for Na^+ and/or K^+ (Giggenbach, 1984). The velocity is buffered by conductive cooling of ascending thermal waters retaining high CO_2 and low pH. Drop in CO_2 and increase in pH under steam loss restricts reactivity of thermal waters in a consequence (D'Amore and Arnórsson, 2000). Except of carbonatogene facies, geothermal waters are usually low in Mg^{2+} thus even its little addition alters the concentration and changes equilibration immediately.

Empirical basis (Giggenbach et al., 1983) assumes equilibration of the fluid with the K-feldspar (adularia), K-mica (illite and muscovite), Mg-bearing silicate (chlorite, clinocllore) and chalcedony assemblage (Eq. 24–25) described at equilibrium activity (Eq. 26–27):



respectively:



with equilibrium constant for Eq. 24 is:




Fig. 11. Na-K system analysis: equilibration processes.


$$K_{eq} = \left[\frac{[\text{Mg}^{2+}]}{[\text{Na}^+]^2} \right] \left[\frac{[\text{albite}]^2 [\text{K-feldspar}]^{0.8}}{[\text{muscovite}]^{0.8} [\text{clinocllore}]^{0.2}} \right] \quad (\text{Eq. 26})$$

meanwhile the latter equilibration reaction follows:

$$K_{eq} = \left[\frac{[\text{Mg}^{2+}]}{[\text{K}^+]^2} \right] \left[\frac{[\text{K-feldspar}]^{2.5}}{[\text{muscovite}]^{0.5} [\text{clinocllore}]^{0.2}} \right] \quad (\text{Eq. 27})$$

These reactions are typical for magmatic environment or can stand sufficiently impure carbonates. Progressive water-rock interaction towards equilibrium to the Na/K system increases K/Mg ratio rapidly as the Mg^{2+} incorporates into magnesium alkalis, such as chlorite (Nicholson, 2012). Alkali alterations (Eq. 24–27) are questionable in low enthalpy sedimentary aquifers, including evaporates, siliciclastics or carbonates, as source of K^+ and Mg^{2+} is rather in dolomite or seawater brine leaching instead (Sonney and Vuataz, 2010).

Likely alteration for impure carbonate environment (Eq. 28) accounts for plagioclase/clay–calcite alteration assemblage (Giggenbach, 1981):



Graphical tests for Mg/Na and Mg/K equilibration are applied to inspect for overall equilibration to both thermodynamic subsystems (discussed below) and overall equilibration or chemistry formation.

Development of K-Mg geothermometer allowed its comparison to the silica thermodynamic system as both are relatively sensitive to fluid dynamics. Their comparable adaptation to new thermodynamic conditions (dilution, mixing, boiling) helps identifying causes of deviation in equilibrium for particular samples or populations. The chart plots equilibrium silica concentration (Eq. 29; Giggenbach et al., 1994) to the K-Mg equilibrium function (Eq. 30), that is, practically, the K-Mg geothermometer by (Giggenbach, 1988).

$$\log(c\text{SiO}_2) = 4.55 - \frac{1\,000}{T + 273.15} \quad (\text{Eq. 29})$$

where silica concentration is in $\text{mg}\cdot\text{kg}^{-1}$ and temperature is in $^{\circ}\text{C}$; and

$$T_{\text{KMg}} (^{\circ}\text{C}) = \frac{4\,410}{14 - \log\left(\frac{K^+}{\text{Mg}^{2+}}\right)} - 273.15 \quad (\text{Eq. 30})$$

where K^+ and Mg^{2+} concentrations are in $\text{mg}\cdot\text{kg}^{-1}$ and temperature is in $^{\circ}\text{C}$.

Possible inspection on increase in aqueous concentrations in fluids due to partial or maximum steam loss is grounded by enthalpy and mass balance (Giggenbach et al., 1994), which gives correction functions for silica (Eq. 31) and K-Mg geothermometer (Eq. 32):

$$\log(c\text{SiO}_2)_b = 5.12 - \frac{1\,215}{T + 273.15} \quad (\text{Eq. 31})$$

and

$$\log(cK^+ / c\text{Mg}^{2+})_b = 14.6 - \frac{4\,630}{T + 273.15} \quad (\text{Eq. 32})$$

where all concentrations are in $\text{mg}\cdot\text{kg}^{-1}$ and temperature is in $^{\circ}\text{C}$.

The K-Mg geothermometer (Appendix B) is the fastest to adapt for Mg^{2+} dynamics quickly responding to re-equilibration. Because of that, the K-Mg geothermometer gives lowest temperature estimates (Giggenbach, 1988), most probably that of last equilibration. For barely real no mixing conditions, the K-Mg geothermometer yields temperature at shallowest reservoir zones. Consistency between K-Mg and silica geothermometers but overestimates for other cation geothermometers is most probably due to short residency and partial equilibration of water-rock system with most reactive phases only (Fournier, 1989).

Application: K-Mg alteration system analysis

Position of facies scatters between the rock dissolution and preferential leaching region (Fig. 12). Facies of DRTE are high in Mg/K ratio, because of poor K^+ but Mg^{2+} rich reservoir environment. Excess Mg^{2+} decreases estimated temperature ($T_{\text{Mg/K}} \approx 42\text{--}55\text{ }^{\circ}\text{C}$) below T_{samp} , disqualifying equilibrium reactions in the microcline – muscovite system (Eq. 24–25).

The residential subfacies of the shallow reservoir dissolve K^+ during downflow off the IWCP horizon, implied by comparison to recharge samples and a region they plot within. Thermal waters derived of DRTE in deeper positions are still affine to parental conditions. Given ionic ratio infers $T_{\text{Mg/K}} \approx 38\text{--}42\text{ }^{\circ}\text{C}$, which may be a temperature of last equilibration to silica (chalcedony) and dolomite system along with talc activation along an upflow ($T_{\text{Mg/K}} > T_{\text{samp}}$). Yet the temperature must be underestimated because SRTW facies intakes Mg^{2+} lost by Mc-calcite precipitation and talc activity. Samples of the VSH-1 plot well along a muscovite stability line, equilibrated at $T_{\text{Mg/K}} \approx 59\text{--}63\text{ }^{\circ}\text{C}$. The temperature corresponds to upper (evasion) zones of the deep reservoir. Yet the stability accounts rather activation of talc and consumption of Mg^{2+} . The increased K^+ is preferentially a sign of its leaching controlled by IWCP hosted S-CGW and preservation, compared to SRTW (Fig. 12).

In fact, thermal waters attain equilibrium to illite and talc rather, varying dolomite and calcite saturation of during upflow/upwell regime. Then, equilibrium reactions given by K/Mg geothermometry (Eq. 24–28) are not valid at local conditions.

Application: Mg-SiO₂ alteration system analysis

We used the chalcedony (Arnórsson et al., 1983), conductive quartz (Eq. 29) and adiabatic quartz (Giggenbach et al., 1994; Eq. 31) models plotted against conductive cooling (Eq. 30) K-Mg geothermometer.

Because of high Mg/K ratio and variable SiO₂ absolute concentrations, DRTE facies tend to organize towards amorphous silica corner at $T_{\text{SiO}_2} \approx 50\text{--}90\text{ }^{\circ}\text{C}$ and $T_{\text{Mg/K}} \approx 30\text{--}70\text{ }^{\circ}\text{C}$. At relatively stable (low) K^+ concentration, the scattering in temperature estimate reflects various Mg^{2+} level, because of change in saturation states of dissolved dolomite and, most probably, because of different dolomite dissolution intensity with vertical filtration. A trend towards amorphous silica is a sign of short reservoir residency. Somewhat reliable temperature estimations give a sign of equilibration with chalcedony, thus Mg^{2+} can either be consumed to activate a talc formation variably.

The SRTW population does not equilibrate to Mg-SiO₂ pair as any component yields reliable results. Inconsistency in a plot is (Fig. 13) evidence of mixed origin. Both

subfacies are undersaturated to chalcedony, which either precipitates or is consumed in dissociation of alkalis or talc activation. The same may be concluded for K/Mg variation, as K^+ enrichment is of low scale only. Discharge facies record more reliable conditions to quartz ($T_{SiO_2} \approx 55^\circ C$) rather than chalcedony ($T_{SiO_2} \approx 30^\circ C$), yet overestimating the K/Mg temperature ($T_{Mg/K} \approx 60^\circ C$). The first is explained by only partial SiO_2 replacement by mixing. Again, high Mg^{2+} content gives more evidence on enrichment by dilution of a mixture with S-CGW, realizing in greater intensity than some K^+ leaching from IWCP or Jr/Cr2 horizon before sampling.




Fig. 12. K-Mg system chemistry forming processes.


Associated thermal waters, regardless of a facies position, are definitely not equilibrated with the Mg-SiO₂ system, for they can not be considered equilibrated to Na⁺, K⁺, and Mg-alkalis. Besides calcite, the inequilibrium is controlled by saturation of dolomite at deep reservoir conditions and variation in free Mg^{2+} and CO₂ controlling talc activity and Na-silicates modification to K-silicates.

Na-K-Ca geothermometer

The Na/K geothermometer gives poor precision for low temperature ($T_{res} < 150^\circ C$) geothermal waters rich in Ca²⁺ and aggressive CO₂. The amount of dissolved carbon dioxide in a geofluid is controlled by dissolution of calcite bearing carbonates or silicates, pH, T and CO₂ fugacity.

In these, the amount of Na⁺ and K⁺ is not controlled by alkalis saturation. Linearity in ionic ratios (Ca/Na, Ca/K, Na/K) over inverse of temperature at HT fields contrast to its scattering, observed in LT systems, has been referred as to Ca²⁺ entering silicate reactions (Eq. 33), thus influencing aqueous Na⁺ and K⁺, if equilibrated (Eq. 34) under the same assumptions as above (Fournier and Truesdell, 1973). As the absolute K⁺ concentration in water

is much less than that of Na⁺, the change in K⁺ is more evident than for the Na⁺ in response to change in Ca²⁺ (Fournier, 1989), so the aqueous K⁺ changes as so as the Ca/Na ratio satisfies (Eq. 34)



$$K_{eq} = \frac{Na^{(1-2y)} \cdot Ca^x}{K} \quad (Eq. 34)$$

where coefficients “y” and “x” represent a stoichiometry of particular reactions.







Fig. 13. K-Mg system chemistry forming processes.

Three Na-K-Ca system reaction configurations (Eq. 35–37) are usually observed, with unifying equilibration constant (Eq. 38)



$$\log K_{eq} = \log \frac{Na}{K} + \beta \log \frac{\sqrt{Ca}}{Na} \quad (Eq. 38)$$

where “solid” denotes the silica or aluminosilicate respectively and “beta – β” depends on stoichiometry of particular reactions (Fournier and Truesdell, 1973).

To satisfy (Eq. 38), not only the excess silica must be present. Aluminum conservation in solids is rather accepted as

neglecting Al^{3+} contents in geothermal waters compared to other constituents, causing equality in deposition and dissociation of hydroxyle minerals so they cancel out of the equation (Fournier and Truesdell, 1973).

For the low temperature ($T_{\text{res}} < 100\text{ }^{\circ}\text{C}$) systems, the $\beta = 4/3$ for the net reaction (Eq. 39), whilst for the high temperature waters the net reaction of base exchange transforms to (Eq. 40) and the stoichiometry derived parameter turns $\beta = 1/3$ (Fournier, 1989), except for waters of $T_{\text{res}} < 100\text{ }^{\circ}\text{C}$ where $\log \sqrt{\text{Ca}/\text{Na}} < 0$ for which $\beta = 1/3$ (Fournier and Truesdell, 1973).

$$K^+ + \left(\frac{1}{3}\right)Na^+ + Ca_{\frac{2}{3}} - \text{solid} = \frac{2}{3}Ca^{2+} + K_{\frac{1}{3}}Na_{\frac{1}{3}} - \text{solid} \quad (\text{Eq. 39})$$

$$K^+ + Na_{\frac{2}{3}}Ca_{\frac{1}{6}} - \text{solid} = \left(\frac{1}{6}\right)Ca^{2+} + \left(\frac{2}{3}\right)Na^+ + K - \text{solid} \quad (\text{Eq. 40})$$

Both models account micas, silicates or clays to fix the aqueous Na-K-Ca system (Shikanozo, 1976). The Na-K-Ca geothermometer (see Appendix B) fits well for thermal waters of various temperatures rich in Ca^{2+} , fixing disregarding activities of Ca-silicates (Fournier and Truesdell, 1973).

Carbonate reservoirs usually lack any relation between actual temperature and cation chemistry of associated waters (Giordano and Swanberg, 1978). Most accurate temperature estimations are obtained for $T_{\text{res}} > 200\text{ }^{\circ}\text{C}$. Erratic results are frequent for low temperature waters of $T_{\text{res}} < 75\text{ }^{\circ}\text{C}$ and fluids with partial pressure of $\text{CO}_2 > 10^{-4}\text{ atm}$ (Pačes, 1975).

Cooler environments (evidenced by petrifying springs) are high in pCO_2 so the pH is controlled by carbonate equilibria rather than by exchange reactions of silicates (Fournier, 1989), thereupon the Na-K-Ca geothermometer scores only if calcite was not deposited after reservoir evasion. In fact, most of thermal waters in carbonate reservoirs are oversaturated to calcium carbonates, influencing their precipitation tendency. Thus the Na-K-Ca geothermometer overestimates reservoir temperatures (Fournier and Truesdell, 1973).

Either dilute (Na)- HCO_3 dominated waters rich in CO_2 are sensitive to Na-K-Ca equilibration (Eq. 39–40). In these, plagioclase likely converts to clay (kaolinite), modifying aqueous pH. Such a process involves interactions with additional Mg^{2+} or Al^{3+} . Typical underestimates are gained so for waters low in Ca^{2+} content and slow evasion velocities, within which a fluid–mineral system is able to react with a wallrock and re-equilibrate. The CO_2 rich springs, if boiled or cooled continuously, yield erratically high estimates owing to intense CaCO_3 deposition and slow Na/K reactions. Mixing has a little effect upon Na/K ratios and precision onto Na-K-Ca geothermometer if the difference in Ca^{2+} concentration is small. If the calcium rich water is diluted, the contact changes the final calcium content for fair (Fournier and Truesdell, 1973). Still, effect

onto Na-K-Ca temperature prediction is negligible unless the parental thermal water component is reduced to less than 30 % (Fournier and Truesdell, 1974).

Mg-correction for Na-K-Ca geothermometer

The Na-K-Ca geothermometer routinely fails in temperature estimation for waters rich in Mg^{2+} . Elevated aqueous Mg^{2+} is expected to originate by dilution, easy uptake of Mg^{2+} out of the wallrock (e.g. chlorite, montmorillonite, dolomite, Mg-calcite) and, for coastal systems, by invasion of a seawater into reservoir (Fournier and Potter, 1978).

In fact, the more the sample deviates from the Na-K-Ca equilibration and purity conditions, the less precise is the $T_{\text{Na-K-Ca}}$ estimate, and the more is the water excess in Mg^{2+} . The magnesium proportion in equivalents is defined by (Eq. 41):

$$R = \frac{100 \text{ Mg}}{(\text{Mg} + \text{Ca} + \text{K})} \quad (\text{Eq. 41})$$

The empirical correction assumes thermal waters of $R > 50$ equilibrated at formation temperatures regardless of $T_{\text{Na-K-Ca}}$ estimate, or assumes their complex unequilibration (Fournier and Potter, 1978). Variation in Mg^{2+} and deviation in functional continuity pronounced mathematical corrections for samples at $5 < R < 50$ (Eq. 42) and $R < 5$ (Eq. 43) to determine the temperature of magnesium correction Δt_{Mg} (Fournier and Potter, 1979).

$$\Delta t_{\text{Mg}}(^{\circ}\text{C}) = 10.66 - 4.7415 \log(R) + 325.87 [\log(R)]^2 - 1.032 \frac{10^5 [\log(R)]^2}{T} \quad (\text{Eq. 42})$$

$$\Delta t_{\text{Mg}}(^{\circ}\text{C}) = 1.03 + 59.97 [\log(R)] + 145.05 [\log(R)]^2 - 36711 \frac{[\log(R)]^2}{T} - 1.67 \frac{10^7 [\log(R)]}{T^2} \quad (\text{Eq. 43})$$

where T is the $T_{\text{Na-K-Ca}}$ in Kelvins (Fournier and Potter, 1979).

The graphical technique (Fig. 14) is presentation of (Eq. 42–43). The X axis represents the $T_{\text{Na-K-Ca}}$ calculated temperature in degrees of Celsius, whilst the Y axis plots desired Δt_{Mg} . Solid curvatures represent the calculated proportion (R) of Mg^{2+} in the sample (Eq. 41). The correction temperature is obtained drawing a vertical line from the X axis at calculated $T_{\text{Na-K-Ca}}$ temperature to intercept the appropriate R function, with later projection of horizontal line onto the Y axis. Each correction must, however, match (Eq. 44):

$$T_{\text{Na-K-Ca}(C)} = T_{\text{Na-K-Ca}} - \Delta t_{\text{Mg}} \quad (\text{Eq. 44})$$



where temperatures are in $^{\circ}\text{C}$ (Fournier, 1980b).

The empirical correction does not account on structural characteristics of reactants and complexing of dissolved solids,

thus it is not based on particular activity coefficients or reaction stoichiometry (Fournier and Potter, 1979). Overestimation preservation after correction is a solid hint on secondary enrichment in Mg^{2+} (usually by dilution).

Lithium geothermometry review

Lithium geothermometers, i.e. Na/Li and Mg/Li (see Appendix B) are the least used in analyses. This is because: (a) there is no immediate geochemical reaction that grounds the relationship between Na^+/Li^+ or Mg^{2+}/Li^+ ; (b) any explicit equilibrium alteration between Na^+ and Li^+ has been proven; (c) both geothermometers appear sensitive to host rock, thus application may be fairly limited by initial lithium concentrations. Best results for both are obtained in sedimentary reservoirs rich in magmatic clasts (Nicholson, 2012). The general relation of Li^{2+} involvement (Eq. 45) rewritten to (Eq. 46) obviously realizes in deeper parts of geothermal systems (Marini, 2004), still, there is a question of contribution of lithium pure end-member silicates (Sanjuan et al., 2003).



Li^{2+} is less reactive compared to Mg^{2+} , Ca^{2+} , Na^+ or K^+ , thus comparison of Na/Li and Mg/Li ratios between springs and borehole samples may give a solid note on upflow velocity (D'Amore and Arnórsson, 2000).

Application: Summary on alkali alteration

The recharges originate by infiltration at various altitudes in the Kľačianka valley area. Because of primary dissolution of carbonates, CO_2 concentration increases with sink and temperature. At this conditions, LT albite

dissolved in a downflow off the IWCP and the Tr3–Cr2 formation transforms to K-micas, i.e. muscovite, even hypothetical, activates just before recharges enter deep reservoir. The oscillation in Ca/Mg, Mg/Na and Mg/SiO₂ recorded for DRTE supports assumption on vertical reservoir filtration within deep reservoir, varying CO₂ level by saturation regime of carbonates and, possibly, LT talc. Upflow regime consumes CO₂ and increases pH stabilizing Na-silicates partially by reduction of activities (not necessarily oversaturating) of illite and muscovite (Figs. 15 and 24), whilst onset of talc activity. Yet chalcedony turns oversaturated (Fig. 5), thus available for LT initial and weak metamorphosis of impure reservoir dolomites (at least at a scale of thermodynamic activities).

The rate of CO₂ consumption increases during evasion towards the shallow reservoir as calcite and Mg-calcite precipitate partially, simultaneous with preferential leaching of Na⁺ from Tr3–Cr2 formations, that takes up more aqueous SiO₂ from solution before inflow. Shallow recharges attain higher K/Na ratio by dissolution of dolomite impurities and hydrolysis of IWCP formations. The SRTW facies is then formed by mixing of both, destroying partial equilibration. The rate of inequilibration is somewhat lower towards the Sliache discharge zone. The Na/K increases simultaneously due to preferential Na⁺ leaching during transition, answering to drop in CO₂ by preferential calcite deposition. Still, the K⁺ is higher compared to DRTE. Alkali source is then not only in rock dissolution, but mixing with SRTW facies. Increase in Mg²⁺ instead of its consumption by Mg-calcite deposition and talc activation is a sign of secondary enrichment by dilution with S-CGW.

Definitely, numerical application of Na-K-Ca geothermometer (Appendix D) disqualifies any equilibration with the Na-K-Ca thermodynamic subsystem. Let all the facies originate in carbonates and form a chemistry by leaching of carbonates in major. Any reactions towards equilibrium




Fig. 14. Mg-correction derivation for $T_{Na-K-Ca}$ at $5 < R < 50$ (left) and $R < 5$ (right).

with Ca-silicates are continuously destructed by excess CO_2 .

Hydrochemical and thermodynamic maturity

In geochemistry of thermal waters, the maturity is understood as a quality referring to reservoir residency analysis and flow history (Giggenbach and Goguel, 1989); or equilibration to Na-K-Mg and Na-K-Ca subsystems (Giggenbach, 1981).

Equilibration to Na-K-Mg-Ca thermodynamic system and CO_2 is objective for cation geothermometry results reliability. This is because cation geothermometers are based on ionic ratios or stoichiometry derivation resultant to partitioning of alkalis (Fournier, 1989), with less focus on other (such as carbonate) systems. Inequilibration causes temperature estimation failure. In other words, immaturity prompts for precise interpretation of given results and allows detailed look on conceptual history and model of sampled waters.

Na-K-Mg subsystem

The Na-K-Mg subsystem composes of three thermodynamic pairs different in equilibration departure velocity and sensitivity to external dynamics (Fournier, 1982). The Na-K pair conserves primary equilibration well, whereas the K-Mg system re-equilibrates, for example by cooling, fast. The Na-Mg pair is of a robust response to any dynamics, pronounced in absence of Na-silicates to control the Na-Mg relation at LT conditions. Facies relation to all is usually inspected with the self-policing Na-K-Mg geoinicator (Giggenbach, 1988).

Use of tri-linear plots with absolute values upscaled and/or normalized to relative governs simultaneous correlation. To facilitate practical application, the concentration is scaled for $\text{mg}\cdot\text{kg}^{-1}$, whilst axes are normalized to accommodate geothermal waters of whatever geothermal system. Straight lines between Na/1000 proportion and $\text{Mg}^{0.5}$ apex represent equal ionic ratios of Na-K geothermometry (Giggenbach, 1988), either due to controlling assemblage or reservoir type and temperature (Giggenbach, 1986). Lines towards the Na/1000 apex at relative K/100 axis show constant K-Mg $^{0.5}$ ratios of K-Mg geothermometer (Giggenbach, 1988).

Intersection of isolines defines complex equilibration temperature to the Na-K-Mg system, termed the full-equilibrium line, above that thermal waters are classified “mature” or “equilibrated” and satisfactory for cation geothermometry application. Position of the curve depends on a choice of the Na-K geothermometer. The curve below represents arbitrary given line representing a dissolution line for “average crustal rock” at the Na-K-Mg system (Giggenbach, 1986). Samples positioned between are “partially equilibrated”, due to mixing or water-rock reactions prior sampling, thus cation geothermometers must be




Fig. 15. Na-K-Mg-(SiO_2) alteration assemblage saturation.

used with a caution in order to assess formation temperatures. A cross-check using another Na-K geothermometer may prove or disapprove that partial equilibration (D'Amore and Arnórsson, 2000). Samples plotted below the average crustal rock dissolution line are “immature” and so “inequilibrated”, and, in most cases, use of Na-K geothermometry for temperature estimation is ambiguous, whilst results of K-Mg geothermometry must be validated by more correlation (Fournier, 1989).

Derivation of the full-equilibrium line is the maturity index (Eq. 47) for geothermal waters regardless of origin, as being a linear combination of Na-K (Giggenbach, 1988) and K-Mg (Giggenbach, 1988) geothermometer:

$$MI = 0.315 L_{km} - L_{kn} = 0.315 \left[\log \left(\frac{c^2 K^+}{c M g^{2+}} \right) \right] \left[\log \left(\frac{c K^+}{c N a^+} \right) \right] \quad (\text{Eq. 47})$$

where concentrations (c) are in $\text{mg} \cdot \text{kg}^{-1}$ and 0.315 is the van't Hoff's slope for linear combination of both geothermometers.

Mature waters yield the $MI > 2$, henceafter use of cation geothermometry is recommended for waters with a maturity index score of above 2 or close around. For mature, partially equilibrated, and rarely inequilibrated parental thermal waters, the corresponding springs plot departed towards the $\text{Mg}^{0.5}$ apex due to acquisition of Mg^{2+} by cooling, mixing or interaction with the wallrock (Marini, 2004). Parental fluids plotted closer to the $\text{Mg}^{0.5}$ corner than a spring or dilutant are signs of silicates or carbonates deposition prior sampling in the borehole, or incorrect steam separation at the wellhead. Meanwhile, the diagram is sensitive to reservoir environment, especially for carbonates, or admixture of low temperature waters (intense thermal – cold mixing) as both, carbonatogene reservoir waters or dilutants are routinely rich in Mg^{2+} and HCO_3^- . These tend to cluster solely in the $\text{Mg}^{0.5}$ corner.

The larger is the distance of the sample from the full equilibrium line, the less reliability gains application of cation geothermometry (Giggenbach, 1988), or, the more caution and cross-validation of partial geothermometry results must be accounted whether for temperature assessment or conceptual relations interpretation.

Application: maturity index analysis

Obviously, associated facies, regardless of position are definitely immature (Fig. 16). Thermal waters of DRTE score $MI = -0.1 - 0.4$ on maturity analysis, classifying them distinctly immature. Samples of SRTW, however, record increase to $MI = 0.2-0.9$. Previously identified sub-facies (low-silica and high-silica) can not be distinguished according to MI. Samples of recharge zone yield $MI = 0.49-0.56$ for shallow and $MI = 0.47-0.54$ for deep approximated circulation regime. Highest score is obtained

for VSH-1 samples, reaching up to $MI = 0.8-0.99$. Springs (S-CGW) of the Sliače area record $MI = 0.51-0.57$.

The maturity index increases with rNa^+ and cNa^+ . This is most pronounced for VSH-1 samples, where maturity is inverse to Mg^{2+} . Enrichment in Na^+ realizes due to mixing of DRTE (Na^+ poor) with SRTW (Na^+ moderate), being followed by the second stage mixing (dilution) with S-CGW (Na^+ rich) prior sampling from the borehole.

Higher Na^+ level is not just because of its slower response to cooling. The S-CGW waters are expected to form and accumulate in siliciclastics. While the dilution balances Mg^{2+} lost in first stage, it adds Na^+ . The relation of rNa^+ and cNa^+ or rMg^{2+} and cMg^{2+} to the MI vanishes within DRTE. The low score relates to Na^+ deficiency and to oversaturation with carbonates. Hence, the Na^+ level is controlled by aggressive CO_2 content in fluids.

Application: Na-K-Mg geoindicator analysis

The structure is immature to the Na-K-Mg subsystem (Fig. 16). To draw the Na-K-Mg diagram we used Na/K and K/Mg geothermometers by Giggenbach (1988). All samples plot as immature (Fig. 17), proving inequilibrium to alkalis, thus ionic ratios overestimate temperatures. All populations score $T_{\text{Na/K}} > 350^\circ\text{C}$. While for K/Mg ratios the DRTE samples exceed a reservoir temperature by a factor of 1–2 ($T_{\text{K-Mg}} \approx 80-140^\circ\text{C}$), samples of VSH-1 cluster along a $T_{\text{K-Mg}} \approx 120-125^\circ\text{C}$, which does not correspond to the reservoir environment.




Fig. 16. Maturity index distribution versus sodium and magnesium concentrations.

Rock dissolution and mineral precipitation regime controls chemistry of all facies. This is, at the same time, controlled by rock dissolution or mineral precipitation. Compared to reservoir facies, samples of the Sliache springs depart towards cooling springs region, evidencing their S-CGW association. Samples of the VSH-1 organize off the cooling springs region from the DRTE facies. The position decreases a relative Mg^{2+} proportion compared to DRTE, combining losses of carbonates, dilution and alkali intake (Fig. 15). Increase in maturity of VSH-1 reflects Na^+ enrichment over stable K^+ level. Similar proportion in Mg^{2+} between SRTW and VSH-1 infers other magnesium source.

An idea of DRTE and SRTW mixing in the upflow zone (decrease in Mg^{2+} , increase in K^+ and Na^+) recalled a first mixing stage preceding dilution of a mixture with S-CGW facies, assumed a second mixing stage (Na^+ increase, Mg^{2+} decrease, K^+ stability) may be satisfied, if the first realizes in Jurassic – Mid Cretaceous succession (Na^+/K^+ source), while dilution takes part in the Mid Triassic dolomites or the Borové Fm. (K^+ source).





Fig. 17. Giggenbach maturity diagram (geothermometer). Modified after giggenbach (1986).

K-Mg-Ca subsystem

A limitation of Na-K-Ca geothermometer is its sensitivity to pCO_2 (Pačes, 1975) not too surprising in extensively magma-steam heated HT or carbonate reservoirs. In the HT equilibrated systems, the hydrolysis of calcium-alumino-silicates towards calcite production (Eq. 48) fixes the pCO_2 (Giggenbach, 1984) at given reservoir formation temperature (Eq. 49):




where the Ca-Al-silicate is isochemically recrystallised stable Ca-mineral, and:

$$\log(fCO_2) = \log(PCO_2) = 0.0168 \cdot t - 3.78 \quad (\text{Eq. 49})$$

where t is temperature of formation in °C.

At initial conditions, the CO_2 fugacity is below that of (Eq. 49), so calcite can not precipitate. CO_2 equilibrium activity drops rapidly with drop in T , turning the fluid reactive with a CO_2 system (Ellis, 1970), recapturing equilibrium by calcite deposition (Eq. 50–51).



$$K_{eq} = \left[\frac{[K^+]^2}{[Ca^{2+}] + pCO_2} \right] \left[\frac{[muscovite][CaCO_3][SiO_2]^6}{[microcline]^3} \right] \quad (\text{Eq. 51})$$

In LT carbonate systems, the carbon dioxide originates by precipitation of calcite and generation of available CO_2 (Eq. 52), or by equilibration to carbonates and production of excess CO_2 by water-rock reaction at continuous contact with a source (Simmons and Christenson, 1993).



Under new equilibration (Eq. 48–50), involving the $CaCO_3$ deposition in a solid form, a correlation (Eq. 53) between concentration of major constituents for stated reactions (Ca^{2+} and K^+) and fugacity is given (Giggenbach, 1986):

$$\log \left[\frac{[c^2K^+]}{[cCa^{2+}]} \right] = \log(PCO_2) + 3 \rightarrow \log(PCO_2) = L_{kc} - 3 \quad (\text{Eq. 53})$$

Presented L_{kc} system adapts to changes in CO_2 or concentrations of constituents fast. While concentrations for major components are a function of variations in temperature, especially cooling, the (Eq. 53) does not account for a CO_2 dependence neither on formation, nor on the equilibration temperature. Correlation of fugacity, geochemical composition and temperature is possible only if combining the L_{kc} system with a geothermometer likely adapting to reservoir dynamics, that is, the K-Mg (L_{km}) geothermometer (Giggenbach, 1988).

The K-Mg-Ca ($K-Ca-p_{CO_2}$) diagram plots samples with L_{kc} to the L_{km} . A solid black line represents a full equilibrium line for CO_2 partial pressures for a full equilibrium assemblage (Eq. 49). If water equilibrates with the K-Mg-Ca and Na-K-Mg system,

samples plot above the line and give a maturity index (Eq. 45) of $MI > 2$ considering them mature or equilibrated (Giggenbach, 1988). These waters are of CO_2 content too low to induce the rock alteration due to (Eq. 48). Low concentration may be a consequence of initial carbon dioxide attainment from the CO_2 deficient host-rocks prior sampling or dilution by magmatic waters with meteoric, or, either by a loss of CO_2 by boiling. A region between the full equilibrium curve and curve for calcite deposition fugacity (Eq. 53) involves samples able to deposit $CaCO_3$ during the upwell. Overall equilibrium to the K-Mg-Ca system is limited by the calcite deposition rate (Giggenbach, 1986). Samples below the dashed line are immature, with chemistry rather as a function of water-rock interaction than water-mineral equilibrium and alteration. Departure from the minimum calcite deposition line implies upflow velocity too fast for calcite alteration onset (Joseph et al., 2013).

Application: K-Mg-Ca geoindicator analysis

The theory expects immature samples to plot below the full equilibrium line. Actual records of all facies within a calcite formation and mature region (Fig. 18) may first be explained by extremely low K^+ concentrations, compared to Ca^{2+} , presumable for carbonate reservoirs.




Fig. 18. K-Mg-Ca geoindicator analysis. Model modified after Giggenbach (1988).

DRTE samples plot along an equilibrium line due to low K^2/Mg and K^2/Ca ratios at TK-Mg $\approx 45\text{--}60^\circ\text{C}$, which is somewhat below the reservoir interval, instantly due to inequilibrium with the K-Mg-Ca subsystem. Assumed LT equilibration with calcite (Eq. 52) is not valid as $pCO_2 \approx 2.6 \cdot 10^{-4} \text{--} 3.2 \cdot 10^{-3}$ bar is too few to induce calcite precipitation. Calcite formation by hydrolysis of aluminosilicates (Eq. 46) yields better counts ($pCO_2 \approx 1.2 \cdot 10^{-3} \text{--} 2.3 \cdot 10^{-3}$

bar), allowing calcite deposition, which may be because of calcite saturation prior deep reservoir inlet. As there are no indices on boiling for calcite (Fig. 19), L_{kc} and L_{km} variation defining the fCO_2 results from vertical reservoir filtration at continuous contact with a wallrock.

In other words, the unit of calcite deposited hypothetically due to pressure change along a vertical upflow filtration is replaced by a unit of calcite dissolved during a vertical downflow in separate convection cells.




Fig. 19. Deep reservoir analysis: calcite saturation.

Temperature overestimate for the shallow reservoir, TK-Mg $\approx 45\text{--}65^\circ\text{C}$, evidences unequilibrium of SRTW facies to the K-Mg-Ca subsystem (Fig. 18), as the K^2/Mg ratio is too low. If we accept a genesis of the facies by mixing of recharges with thermal waters evaded from the deep reservoir, the excess Mg^{2+} reflects continuous dolomite dissolution within CHN dolomites and fresh intake, balancing the Mg^{2+} lost off the DRTE.




Fig. 20. Deep reservoir analysis: dolomite saturation.

Presumably, Mg-calcite precipitates (Fig. 20) after evasion from the deep reservoir. Calcite addition produces excess CO_2 and decreases an apparent pCO_2 . Under low temperatures, the K^+ increases slower than Mg^{2+} , as K^+ leaching (simultaneous to alteration of Na-silicates to K-micas) is less intense with drop in CO_2 .

In absolute concentrations, K^+ increases in VSH-1 samples, compared to DRTE, whilst keeping similarity to SRTW, either at lower Ca^{2+} . K^2/Ca and K^2/Mg ratios increase probably at a magnitude of calcite and dolomite deposition during DRTE and SRTW leaking towards discharge zone, conserving K^+ prior mixing and later enrichment of a mixture with K^+ and Mg^{2+} from dilution with S-CGW. After projection onto CO_2 fugacity line, the $T_{\text{K-Mg}} \approx 50^\circ\text{C}$, which evidences drop in aggressive CO_2 by some calcite and Mg-calcite deposition and ceasing of Na-silicates modification towards K-micas, calling the (Eq. 52) apparently valid.

Na-K-Mg-Ca geoinicator

The Na-K-Mg-Ca geoinicator (Giggenbach, 1988) combines Na-K-Mg and K-Mg-Ca thermodynamic subsystems, comparing relative proportions of Mg over Ca (carbonates) and Na to the K (alkali silicates). The solid line represents here a full equilibrium to both thermodynamic subsystems (Fig. 21). Samples plotted below the curve are already equilibrated without a tendency to dissolve host-rocks spontaneously (Wishart, 2015). Instead, the geochemistry of such samples is controlled by mineral-water equilibrium.

Average seawater (Na-Cl brine) region locates within high Mg^{2+} /low K^+ corner, whilst high Mg^{2+} /high K^+ corner represents a rock dissolution region. Here, samples plot outermost to the equilibrium, thus are immature and geochemical composition is controlled by a water-rock interaction (Giggenbach, 1995), i.e. dissolution well n, hydrometasomatism or hydrolysis. Affinity to subregions representing average crustal rock composition (Taylor, 1964) is controlled by relative ratios of the samples. Transition of samples towards equilibrium line in a direction of X axis indicates a possible seawater influence on their final geochemistry. A diagonal transition trend is an indice of mixing or dilution (Giggenbach, 1988).

Ionic ratios in a crustal rock region represent those needed to attain a full equilibrium to a specific rock. Drawing a vertical line onto X axis conserves Na/K ratio and can account on expected L_{km} temperature. A horizontal line projected onto Y axis gives an estimation by L_{km} (Giggenbach, 1988). Interpretation must, however, take a caution onto both values as they can not gain same values in an essence. This plot is then, rather, used to outline reservoir or conceptual relations.

Application: Complex maturity analysis

Organization of all facies along and nearby the rock dissolution region is distinct evidence on immaturity and unequilibration to both subsystems as they plot outermost to the full equilibrium line (Fig. 21), recording three facies trends meanwhile.

Deep reservoir samples plot within or to the right of rock dissolution region. The equilibrium attainment realizes along an Y axis, evidencing the equilibration due to saturation of waters with carbonates. A drop in Mg/Ca ratio is observed along a DRTE/SRTW trend (1), implying Mg-calcite precipitation rate above that of “pure” calcite, or its acquire from Jr-Cr2 horizon. Apparent increase in Na/K towards the shallow reservoir gives some evidence on Na-silicates modification deceleration with drop in CO_2 and possible enrichment within.

Some equilibration to the wallrock is possible for residential SRTW subfacies, destructed by a mixing with DRTE originated subfacies instantly. At cooler downflow temperatures (2), limestone is leached easier than dolomite, which dissociation starts after the reservoir accommodation. Increase in alkalis in downflow is inverse to filtration velocity decline in the Borové Formation that is a source of Na^+ . The K^+ originates by dissolution of silicates and impurities in dolomite, with questionable contribution of Na-silicates modification.

VSH-1 facies record somehow a most evident trend towards partial equilibration, off the rock dissolution region. In fact, only weak relation may be found between SRTW, DRTE and VSH-1 facies (3) as a sign of poly-




Fig. 21. Facial analysis: Na-K-Mg-Ca geoinicator. Modified after Giggenbach (1988).

stageous mixing. Some increase in Na^+ , compared to parental conditions is either because of fair conservation of alkalis from SRTW and dilution with S-CGW hosted in IWCP (second mixing stage), however, effect of minor Na^+ preferential leaching in Jr-Cr2 horizon can not be excluded.

CO_2 induced modification of dissolved alkali silicates must be terminated already due to its consumption in deposited carbonates and activation of Mg-silicates, such as talc.

Quantitative cation geothermometry application

Na/K geothermometers (GN1–GN11) yield temperatures of over 100 °C for each zone, consequent to inequilibrium with Na/K system, excess CO_2 and deficient source. Variation by a magnitude of orders is also general for Na/Li geothermometers (GLN1–GLN5), resultant to combination of lack in either Na^+ and Li^{2+} . Temperature overestimates by a factor of 2–5 result from Na/Ca (GNC), K/Ca (GKC) and Na-K-Ca (GNKC) application, consequent to oversaturation of fluids with calcite and excess CO_2 in the system available, immediately restraining any equilibration to given thermodynamic pairs (Appendix D).

K-Mg geothermometry

Deep reservoir waters are oversaturated with dolomite under permanent contact producing excess Mg^{2+} and CO_2^- . The K^+ is consumed in activation of K-tectosilicates due to Na-silicates modification under aggressive CO_2 (cf. Giggenbach, 1986). Both factors cause high Mg/K ratio, underestimating DRTE temperatures by a factor of about 2 (Fig. 22). Apparent match of GKM3 (Giggenbach, 1988) at $T_{\text{Mg/K}} \approx 30\text{--}41$ °C is observed for the SRTW. Equilibration with Mg/K-silicates is not possible, rather the result is underestimated with respect to DRTE evaded subfacies by balancing Mg^{2+} lost during Mg-calcite deposition with its leaching from dolomites and enrichment from recharges, more intense than K^+ intake from the IWCP formations. Indeed, the K-Mg-SiO₂ relation aids assumption on fluent rewashing of the zone and residency too short for any equilibrium attainment. Depending on a depth of sampling, GKM3 scores reasonable estimates for the SRTW as long as $\text{K/Mg}^{0.5} = 0.5\text{--}14$. GKM3 yields $T_{\text{Mg/K}} \approx 31\text{--}33$ °C at VSH-1 (Fig. 22). Following qualitative applications, the temperature is definitely underestimated, as $T_{\text{Mg/K}} \approx T_{\text{sam}}$. Inflow temperature of waters from both reservoirs prior mixing must be, at least, of $T = 45$ °C or $T = 40$ °C respectively. We would not recommend K/Mg geothermometry quantitative application in local conditions.

Li-Mg geothermometry

A difference in lithium concentrations from early years (1987–1992) of deep reservoir production ($\text{cLi}^{2+} \approx$




Fig. 22. Review on cation geothermometry quantitative application. See Appendix B for references and symbols.

1.4–2.5 mg·kg⁻¹), compared to steady lithium lowers that has been monitored at ZGL-1 or FGTB-1, henceforth ($c\text{Li}^{2+} \approx 0.2\text{--}0.3 \text{ mg}\cdot\text{kg}^{-1}$) varies Li/Mg estimations from $T_{\text{Li/MgO}} \approx 65\text{--}83 \text{ }^{\circ}\text{C}$ towards current $T_{\text{Li/MgO}} \approx 27\text{--}37 \text{ }^{\circ}\text{C}$. This is easily explained by the disproportion between low Li^{+} and Mg^{2+} , even magnesium is upscaled by a root of square. There is just no sign on Li-clays equilibration (Eq. 45–46).

Such a drop may then be because of initial marinogene residuals removal or due to increasing an overall deep filtration velocity within the structure, as not only Li^{+} , but Cl^{-} decreases in comparable scale (Fig. 23). Definitive rewashing of long-term and long-residential waters while building up the hydraulic gradient between recharge and accumulation zone towards pseudo-steady state would also drop the Li^{+} .

Stable Li^{+} concentrations are recorded in the shallow reservoir ($c\text{Li}^{+} \approx 0.3\text{--}0.6 \text{ mg}\cdot\text{kg}^{-1}$) during its production (1979–2015) yielding $T_{\text{Li/MgO}} \approx 39\text{--}52 \text{ }^{\circ}\text{C}$, with no relation to sampling depth. The lithium content comes off its conservation during DRTE upwelling or leaching from IWCP and Jr-Cr2 positions in a downflow regime of recharges. Either the way, Li^{+} appears conserved only, not entering any ion exchange reactions, questioning Li/Mg geothermometry application as no equilibrium is attained.

With given VSH-1 samples (see Appendix C) the score in temperature ($T_{\text{Li/Mg}} \approx 62\text{--}69 \text{ }^{\circ}\text{C}$) estimation at ($c\text{Li}^{2+} \approx 1.0\text{--}1.4 \text{ mg}\cdot\text{kg}^{-1}$) is rather a hint on additional Li^{2+} addition from a host rock while waters of deep and shallow reservoir laterally leak towards discharge zone. For this interpretation, we would prefer the influence of transition at a contact with sufficient source, such as IWCP, or with Jurassic–Mid Cretaceous succession.




Fig. 23. Time-domain variability in selected ion constituents at ZGL-1.

Too few correlations for lithium geothermometry exist, available for detailed look on lithium origin. The deep reservoir waters are of higher temperature, thus may dissolve Li^{+} intensively, thus do not need a long water-rock contact. Shallow reservoir

waters are of half a lower temperature, but remain constantly in potential lithium source rock. If both facies mix together, the resultant lithium is then proportional to rate of mixing. A process to control the lithium level is, most probably, a simple enrichment, as association of lithium clays in ion-exchange reactions is unlikely. As far as uncertainty onto lithium origin exists up to a recent state, use of lithium geothermometry in local conditions is rather not recommended.

Na-K-Ca-Mg corrected geothermometry

Thermal waters of the Bešeňová elevation, regardless of facies affinity are acid, referring to volatile CO_2 content, thus with calcite and dolomite precipitation in control on aqueous pH, explaining why most of cation geothermometers fail in quantitative application (Appendix D). Besides saturation of calcite, aragonite or dolomite (Figs. 19–20), structures hosted in carbonates are usually rewashed, fast in filtration and diluted by recharges or during upwell in discharge zones, thus rich in excess Mg^{2+} .

For the deep reservoir, the excess Mg^{2+} is obviously controlled by magnesium (dolomite, calcareous dolomites, talc) saturation cycle. Using the Mg-correction, the initial $T_{\text{NKC}} \approx 211\text{--}325 \text{ }^{\circ}\text{C}$ decreases to $T_{\text{NKCm}} \approx 27\text{--}95 \text{ }^{\circ}\text{C}$, out of that 24 % of samples (8/33) do not match the reservoir interval, with one overestimate only. There is no external Mg^{2+} source than dissolution of dolomites and LT alteration of impure dolomites. Correction estimation range implies rather variation in saturation of dolomite and talc thermodynamic activation.

Even after correction temperatures are overestimated, both in shallow reservoir ($T_{\text{NKCm}} \approx 61\text{--}79 \text{ }^{\circ}\text{C}$) and VSH-1 samples ($T_{\text{NKCm}} \approx 42\text{--}49 \text{ }^{\circ}\text{C}$). For both, the function can not be used in equilibrium conservation correlation to the origin of formation. The mismatch points to excess Mg^{2+} due to additional dolomite dissolution or mixing (dilution) with recharges, aiding previous balance of Mg-calcite deposited during lateral leaks or vertical evasion.

COMMENTS ON SOLUTE GEOTHERMOMETRY

Isotope and gaseous geothermometry

Use of solute geothermometry is grounded by conception of relative stable upflow during which a fluid does not undergo any mixing with cooler groundwater or is at sub-boiling conditions. Development of gas and isotope geothermometers responded to problem of inconsistency of many thermal springs with an obligatory conception in essence. As both are not a scope of our paper, we deal with either only in a brief review.

Thermal springs are not the only surface geothermal propagations. Many fields manifested by fumaroles do not involve springs, as the water table remains deep in grounds, thus aqueous geothermometers can not be applied (Giggenbach, 1991). The fact resulted in development of the gas geothermometers based on gas–gas equilibria, gas–mineral equilibria and mineral–residu-

al gas equilibria, related to CO_2 , H_2S , H_2 or CH_4 (e.g. Arnórsson and Gunnlaugsson, 1985; Arnórsson et al., 1998). Steam geothermometry is more difficult to handle, because: (a) gas concentrations are affected by water–steam ratio; (b) in fumaroles, the gas content of a steam is affected by mechanism of boiling and condensation, and; (c) gaseous components fluxes from external reservoir or magmatic source may be of influence overprinting initial mineral–gas equilibria (D’Amore and Arnórsson, 2000). Application of these is ambiguous for low enthalpy systems with steam phase separated artificially.

In natural systems, equilibrium achievement for isotope exchange reactions is temperature dependent (Ellis and Mahon, 1977). Isotope geothermometers are less sensitive to thermal water flow dynamics, as achievement on chemical equilibrium is obligatory prior isotope redistribution into equilibration (Lyon, 1974). Out of many proposed, most frequently used geothermometers for high enthalpy fluids are those based on $\text{CO}_2 - \text{CH}_4$, $\text{CH}_3\text{D} - \text{H}_2$ and $\text{HD} - \text{H}_2\text{O}$, $\text{CO}_2 - \text{H}_2\text{O}$ and $\text{SO}_4 - \text{H}_2\text{S}$, whilst for low enthalpy systems, applicable isotope geothermometers refer to $\text{SO}_4 - \text{H}_2\text{O}$ and $\text{CO}_2 - \text{H}_2\text{O}$ (D’Amore and Arnórsson, 2000).

Critique on solute (cation) geothermometry

Criticism accents violating principal aspects of formal chemistry in setting assumptions for grounding and application of cation geothermometers. In balancing equilibrium reactions (e.g. Eq. 17), solids on both sides are not the same and there can not be a free ion in equilibrium reaction, thus the equation can not be in equilibrium, and mixed minerals are not pure phases, so their activity coefficients can not equal one. Another impossibility is in existence of a fraction of atom in formula (e.g. Eq. 39). In practical aspects of computational solute geothermometry, adds imbalance in physical units within solute geothermometers equations, as there are numbers on a one, and variable with a unit on the other side (Verma, 2012).

Silica geothermometry has also been subjected to criticism of its basics. Experimental solubilities of silica decrease along a water-vapor saturation curve at temperatures above 300 °C, neglecting a vapor formed fraction. It is mandatory to account on total silica discharge, thus to include a silica in the vapor fraction, in construction of silica solubility basics. To know that, correct knowledge on reservoir temperature is crucial. Then a role of silica geothermometers limits to reservoir water-rock interactions and equilibrium states (Verma, 2000). Nor SiO_2 geothermometry models are balanced in physical units (Verma, 2012).

Isotope geothermometry techniques and understanding on isotope cycles in thermal waters is in a constant development. There is, however, a rising accent on their relation not only to temperature, but onto influence of reservoir pressure and salinity addition as well, which is not included in conventional isotope geothermometry calculations (Horita, 2005). At the same time, variation in isotope thermodynamic history due to mixing is frequently missed out.

MULTICOMPONENT EQUILIBRIUM GEOTHERMOMETRY (MCG)

Instead of solubility of one or a few minerals and semi-empirical correlations, the multicomponent geothermometry uses complete fluid analysis and a solid thermodynamic background at given assemblage. The complexity provides the multicomponent geothermometry with the advantage of a great precision, if applied correctly. Exactly, minimum assumptions grounding the MCG turn this method most reliable, whether in temperature estimation, saturation states analysis, equilibrium correlations and CSM construction.

Mineral equilibria

Saturation state

Chemistry of thermal waters is not only a print of co-existing dissolved phases (Arnórsson et al., 1982; Reed, 1982), but a record of its conceptual history either. One of consequences of the conceptual history is an equilibrium state (Eq. 53), at which the activity quotient and equilibrium constant (Eq. 54) are equal, so the saturation index $SI = 0$ (Reed and Spycher, 1984).

$$SI = \log \left(\frac{Q_k}{K_{eq}} \right) \quad (\text{Eq. 53})$$

where Q_k is the activity quotient for aqueous species formed by the dissolution of a mineral “k” and K_{eq} is the equilibrium constant for the dissolution reaction of a mineral “k”:

$$K_{eq} = \frac{\prod_{i,k} a_{i,k}^{v_{i,k}}}{a_k}, \text{ and } Q_k = \prod_{i,k} a_{i,k}^{v_{i,k}} \quad (\text{Eq. 54})$$

where “ $a_{i,k}$ ” – activity coefficient, “ $v_{i,k}$ ” – stoichiometric coefficient for species “i” in equilibrium mass action expression for phase “k”, “ a_k ” – activity coefficient of „k-th“ phase.

If the phase is undersaturated ($SI < 0$), the thermal water tries to approach its equilibrium by dissolving/leaching activation from a wallrock or onset of equilibration ion-exchange reactions, or, in other words, the phase and its components remain in solution. At oversaturation ($SI > 0$), the water lets the phase to contribute on equilibration reactions or deposits, the phase after loosing a contact with a source rock or if p-T conditions change dramatically (e.g. onset of cooling, boiling).

A frequent technique in analysis of saturation states for one or more minerals is plotting the SI over an optional range of T, which implicitly displays thermodynamic state of mineral phases of various composition, instead of using activity diagrams where only two compositional variables are plotted at fixed temperature (Reed and Spycher, 1984).

Approach

The recalled complexity is, at early exploration stages, of greatest disadvantage for the MCG, requiring not only at least some knowledge on associated mineral suite, but a solid concept on reservoir dynamics, i.e. generation of adiabatic boiling and processes of heating and cooling during conceptual history. For the MCG, only deep reservoir is evaluated, as this is identified the only zone not affected by any mixing or dilution.

In a sake of review, geochemical models (Fig. 2) together with geothermometry (e.g. Figs. 6 and 7) give a solid supposition on conductive heating/cooling controlling reservoir intake or runoff temperature variations regardless of direction.

Computation module

The MCG targets saturation states of associated mineral phases. Several notes onto the problem have been listed e.g. in Remšík et al. (1998), as well as Vandrová et al. (2011), aimed at actual, sampling temperature saturation of the deep and shallow reservoir. The MCG, however, requires analysis over desired temperature interval, which usually means between reservoir base and reservoir outlet, or, rather, geothermal infrastructure intake temperature, with possible extension to reference temperature assuming constant p-T conditions.

Under local conditions, we set the temperature interval for $T = 20\text{--}100\text{ }^{\circ}\text{C}$, referring to usual reference temperature and basal temperature extended by $\sim 10\%$ for better precision at the base. Saturation calculations (Eq. 53–54) were carried using Phreeqc (USGS) and WATCH/WAIN (ÍSOR) scripta, applying cooling springs, no boiling $\approx$ conductive cooling/conductive heating modules off the sampling temperature. At this setup, both modules are able to vary thermodynamic properties and SI or desired solids, not changing macrochemistry of analysed solution.

Selection of fluid samples

Whether for simple saturation indexes (SI) or advanced multicomponent geothermometry (TSI, RMED) analysis, we limited calculations to deep reservoir waters only. The reservoir has been selected because it represents the only zone within the structure not affected by any mixing or dilution. Obviously, the reservoir intake from the deep infiltration channels is expected to be of quasi-isochemical regime. To narrow the population, the modelling procedure involves only samples of complete chemical analysis in referenced literature (see Appendix C). We iterated then the selection towards samples which yielded relevant temperature estimate at chalcedony geothermometer (Fournier, 1977) simulation in WATCH. To tighten homogeneity of samples, the last population reduction selected samples taken at wellhead only through a sampling campaigns since 1988.

Complex equilibrium temperature (T_{CEQ})

Type curve method

Interpretational procedures using SI vs. T diagrams are based on convergence and shape analysis. In T_{CEQ} evaluation, equilibrated assemblage intercepts the SI zero level at a definite temperature, or, at least, at a narrow temperature interval reservoir boundary conditions (Pang, 1988). In fact, no more than 10–12 phases coexist in equilibrium (Arnórsson et al., 1982). More interception clusters imply inequilibrium or partial re-equilibration. Conceptual decision is rather a question of the interpreter. Equilibration to a relevant assemblage may be well seen by sharp maxima and irregular skewness of histograms (Reed and Spycher, 1984).

Boiling effects upon saturation states relate to an immediate change in pH and loss of CO_2 or H_2S respectively. For silicates, the $\text{SI} = 0$ intersection temperature increases as SiO_2 concentration in residual fluid is higher as of an initial (Tole, 1988). Mineral phases containing sulphide, aluminum and iron shift equilibration compared to the pre-boiling conditions as speciation of H_2S , Al^{3+} and $\text{Fe}^{2+,3+}$ changes with an increase of CO_2 . However, the loss of CO_2 definitely causes supersaturation with respect to carbonates (Reed and Spycher, 1983). Boiling plots irregular dispersion and shifting of SI curves generally (Neupane et al., 2014).

By mixing with cooler thermal water or mineralized groundwater, the convergence temperature for particular phases shifts irregularly, and there is usually neither equilibrium attained, nor convergence clusters may be identified. A contact with dilutant (Fig. 11c), however, shifts apparent equilibrium to negative or causes “re-equilibration”, or, simply, double interceptions for phases or complete assemblage (Tole, 1988).

Application: Selection of solids procedure

Numerous methods have been developed since the concept of equilibrium (saturation) analysis was introduced (e.g. Reed and Spycher, 1984). Besides computation and analysis principles, these methods vary in mineral assemblage selection/definition. Complex equilibrium analysis selects the assemblage by arbitrary.

Together 18 solids were selected to represent reservoir and conceptual model conditions, assigned into groups of master- primary (calcite, dolomite, chalcedony, gypsum, talc), alteration (albite, adularia, fluorite, kaolinite, illite, muscovite, chlorite) and hypothetical- minor phases (montmorillonites, gibbsite). Master and alteration phases control geochemistry of reservoir waters by their saturation states. Minor phases were introduced for additional control on stability for major components and pH. Neither

alteration, nor hypothetical phases must necessarily precipitate because of oversaturation. Their state of equilibrium is called to control ability of master components (OH^- , K^+ , H^+ , Na^+ , F^- , Al^{3+}) to enter wallrock dissolution or ion exchange reactions, with respect to CO_2 reservoir cycle.

Application: Complex equilibrium temperature analysis

Derivation of complex equilibrium temperature requires calculation of temperature at which the saturation state (Eq. 53) equals $\text{SI} = 0$ for each of selected phases (Tab. 1) per every sample. With this procedure, the overall temperature varies $T_{\text{CEQ}} = 56\text{--}79\text{ }^\circ\text{C}$, as median values of $T_{\text{SI}} = 0$. Because of skeweness of results, the representative temperature of complex equilibrium equals $T_{\text{CEQ}} = 66\text{ }^\circ\text{C}$. Out of 20 samples, only two were disqualified, as number of equilibrated phases within $T = 20\text{--}100\text{ }^\circ\text{C}$ has not reached 50 % (9/18). The $T_{\text{CEQ}} = 66\text{ }^\circ\text{C}$ corresponds to upper half of the deep reservoir.

Quantitative equilibration analysis may be referred to reservoir temperature ($T_{\text{res}} = 56\text{--}79\text{ }^\circ\text{C}$) or entire temperature ($T = 20\text{--}100\text{ }^\circ\text{C}$) interval. The first approach inspects reservoir conditions

only, whilst the latter allows evaluation of not only reservoir, but runoff conditions simultaneously, assuming no mixing and maintaining conductive heat loss as a controlling cooling process only.

Only 8 phases equilibrate at over 50 % frequency at T_{res} . Only chalcedony, Na- and K- montmorillonites equilibrate completely. Chalcedony saturation proves its control on silica solubility (Fig. 5). Equilibration of montmorillonites proves the adularia–albite system is not in control on Na/K stability (see Fig. 15). Adularia (mean $T_{\text{CEQ}} = 55\text{ }^\circ\text{C}$) equilibrates in 16 samples at ($T_{\text{CEQ}} = 40\text{--}65$), however, in 8 samples within the reservoir. Albite (mean $T_{\text{CEQ}} = 42\text{ }^\circ\text{C}$) intercepts $\text{SI} = 0$ in 9 samples only ($T_{\text{CEQ}} = 30\text{--}46$), not reaching equilibrium within reservoir. Anhydrite and fluorite never converge to $\text{SI} = 0$ at reservoir temperature as well. Inequilibration with anhydrite aids control of gypsum saturation on SO_4^{2-} content within deep reservoir. Gibbsite, kaolinite and chlorite (Tab. 2) equilibrate in over 15 samples. First two simulate available OH^- and Al^{3+} for alkali hydrolysis reactions. The Mg^{2+} available by chlorite saturation variation allows its re-attainment within LT metamorphism of detritic dolomites, activating talc, leaching Mg-calcite and producing additional CO_2 .

Tab. 1
Deep reservoir: review on CEQ of selected samples

Mineral phases	Wellhead samples codes																			
	z21/2/1987	z4/3/1987a	z4/3/1987b	z5/3/1987	z6/3/1987a	z6/3/1987b	z7/3/1987	z22/3/1987	z31/3/1987	z2/4/1987	z3/4/1987	z20/2/1992	z12/11/2007	z17/12/2008	z18/2/2008	z26/2/2009	z30/3/2011	z13/4/2011	f15/3/2011	f30/3/2011
Equilibrium temperature ($^\circ\text{C}$)																				
Adularia	50	49	61	U	51	59	U	55	U	40	U	58	48	55	53	53	56	54	65	63
Albite	37	U	U	U	U	U	U	43	U	U	U	43	U	31	42	42	46	43	U	U
Anhydrite	20	22	25	20	22	24	22	21	20	21	O	O	O	O	O	O	O	O	O	O
Calcite	O	O	20	45	O	O	62	O	25	O	O	O	O	O	O	O	O	O	O	O
Ca-mont.	71	78	O	O	88	91	O	80	86	84	84	74	76	80	71	73	74	74	85	86
Dolomite	O	31	82	82	46	45	U	O	45	32	29	O	O	O	O	O	O	O	O	20
Fluorite	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O
Am. pyrite	56	45	O	U	U	U	U	22	U	57	U	55	U	60	46	55	55	62	79	80
Gibbsite	53	62	69	67	65	65	70	58	64	64	62	48	58	58	51	53	50	53	58	61
Chalcedony	56	64	94	81	72	81	94	67	67	67	67	72	58	67	60	62	66	62	77	79
Chlorite	61	85	97	99	94	91	O	73	88	87	83	44	77	74	53	60	52	60	74	80
Illite	49	U	64	U	55	61	U	55	U	52	52	51	51	54	50	52	52	52	59	60
Kaolinite	72	83	O	O	88	90	O	79	87	84	84	71	77	79	71	73	72	73	81	84
K-mont.	62	67	90	81	75	79	99	69	73	71	71	63	65	69	61	64	64	64	75	72
Mg-mont.	73	80	O	O	89	92	O	81	88	85	85	75	77	81	72	75	75	75	86	87
Na-mont.	62	68	92	84	76	80	O	70	74	73	73	63	66	70	62	64	64	65	74	75
Talc	38	63	74	81	71	66	83	49	66	64	60	O	54	44	27	35	25	35	45	52
Muscovite	91	95	O	O	O	O	O	98	O	O	O	92	95	98	91	93	93	93	O	O

Explanations: O – oversaturated; U – undersaturated

Tab. 2
Deep reservoir: review on CEQ of selected samples

Mineral phase	Reservoir equilibration frequency		Main SI = 0 interval
	x = n	%	(°C)
Adularia	8	40	50–55
Albite	0	0	35–40
Anhydrite	0	0	20–25
Calcite	1	5	25–30
Ca-mont.	17	86	70–75
Dolomite	2	10	80–85
Fluorite	0	0	–
Am. pyrite	9	45	55–60
Gibbsite	15	75	55–60
Chalcedony	20	100	65–70
Chlorite	17	85	70–75
Illite	2	10	45–50
Kaolinite	18	90	80–85
K-mont.	20	100	60–65
Mg-mont.	17	85	70–75
Na-mont.	20	100	70–75
Talc	9	45	65–70
Muscovite	9	45	90–95

Application: Type curve analysis

The type curve analysis applies to inspect reservoir dynamics. In previous, we already plotted chalcedony (Fig. 5), calcite (Fig. 19), dolomite (Fig. 20), and adularia–albite–muscovite suite (Fig. 15). Primary phases record uniform trend in SI-curves throughout the analysed interval. Alteration minerals, however, plot some minor variation (Figs. 15 and 24) at $T < 65$ °C. Because most of these are alkalis, we interpret the observed break as due to unstable contact with reservoir Na^+ and K^+ source (at $T > 65$ °C), and variable CO_2 saturation, which boosts or limits Na-silicates conversion into K-micas or phyllosilicates, with some secondary effect on aqueous Al^{3+} during the onset of conductive cooling as the sample is simulated to leave the reservoir.

A unifying feature of all phases plotted against temperature is variation in $\text{SI} = 0$ convergence, much above tolerated $\delta T_{\text{SI}} = 0$ difference of 10 % (± 9 °C). This is some sign of intraformation mixing, however, in pseudo-isothermal and pseudo-isochemical regime. The only possible scheme to fit this postulate is intraformation mixing of waters in vertical filtration due to formation of isolated, induced-convection cells, with steady, accommodated waters within upper parts of the reservoir.

Multicomponent equilibrium

Since the MEG was introduced (Reed and Spycher, 1984) and modified (Pang and Reed, 1998; Palandri and Reed, 2001), the application of some conceptual approaches aimed either at T_{res} derivation (e.g. Tole et al., 1993; Neupane et al., 2014), or initial reservoir fluids and formation brines reconstruction with onward temperature predictability improvements (e.g. Peiffer et al., 2013; Peiffer et al., 2014; Spycher et al., 2014). Hundreds of minerals can be easily plotted onto the SI vs. T diagram, making it all but useful. Moreover, dozens of assemblages may be misleadingly analysed, if these are selected inappropriately. Crucial for multicomponent equilibrium geothermometry is to know the assemblage and the procedure searches for, thus, the assemblage that is in control of chemistry of water samples.

Phase rule

The Gibb's phase rule identifies a number of minerals that may equilibrate with the geothermal fluid by defining degrees of freedom (Eq. 55), controlling a real number of components ready to equilibrate (Eq. 56), if temperature and pressure is fixed in the system, e.g. in the case of water saturated with steam (Cooper et al., 2013):

$$F = C - P + 2 \quad (\text{eq. 55}),$$

where F is a number of independent variables, C is a number of components and P is a number of phases in the system, so the number of equilibrium minerals (M) becomes:

$$M = C - F \quad (\text{Eq. 56})$$

The phase rule gives an upper and theoretical limit for number of minerals that may coexist in equilibrium, even a real number of equilibrated minerals may be significantly less (Palmer et al., 2014), as this is due to combination of conceptual history of a fluid and p-T conditions.

In this paper, we have continuously accented multiple geothermometry application, however, temperature estimation resonates as the principal. Reservoir temperature may be derived by applying the TSI (Cooper et al., 2013) or CSI/RMED (Spycher et al., 2011; Spycher et al., 2014) methods.

Total saturation index (TSI)

Assemblage selection does not necessarily impede unreadability of SI vs. T plots if disarranged clustering or several re-equilibrations are displayed. The optimizing function for considered mineral assemblage is termed as




Fig. 24. K-Mg-(SiO₂) alteration assemblage saturation.

the Total Saturation Index – TSI (Cooper et al., 2013). In cited publication, the function is squared (Eq. 57) to search for loss in volatiles and reconstruction of amount of CO₂ degassed. However, if there is no boiling in the reservoir, alternative linear form (Eq. 58) can be used to search for absolute saturation states and yield hints on overall reservoir situation (Fričovský, 2014).

$$TSI = \sum_{i=1}^n (SI_i / wt_i)^2 = \sum_{i=1}^n \left[\left(\log \frac{Q_{k-i}}{K_{eq-i}} \right) / wt_i \right]^2 \quad (\text{Eq. 57})$$

$$TSI = \sum_{i=1}^n (SI / wt_i) = \sum_{i=1}^n \left[\left(\log \frac{Q_{k-i}}{K_{eq-i}} \right) / wt_i \right] \quad (\text{Eq. 58})$$

The (Eq. 57) is square-termed, thus always greater than zero. Temperature is estimated by plotting possible amounts of evaded CO₂ and searching for a function with one minima converging to zero TSI level. Then, the point at which the curve converges the TSI = 0 may be a solid hint on reservoir temperature (Palmer et al., 2014).

This allows to check for accuracy in degassing coefficient evaluation for SI calculations or, reversely, the optimized CO₂ loss may be revised by evaluation of degassing coefficient by trial-and-fail procedure while calculating SI of partial assemblage components. Initial CO₂ level estimation helps to reconstruct initial fluid composition prior sampling (if the fluid flashed before the action) or boiling respectively.

Use of linear function (Eq. 57) gives rough estimation of total equilibrium states in the reservoir, and, apparently, gains robust estimation of prevailing undersaturation or oversaturation. A hint on reservoir temperature may be read of the TSI = 0 line interception. Both functions equal for non-boiled, equilibrated waters.

TSI type curve analysis

Linear TSI functions (Eq. 58) are similar to those plotted as SI vs T, grouping number of phases together instead of plotting them individually. An assumption is the equilibration (no boiling, no mixing) of records TSI = 0 interception at narrow ($\delta T_{TSI} < 10\%$ of $T_{btm} - T_{top}$) temperature interval and stable shape.

Fluids mixed prior sampling record unstable interception because of various cooling and changed saturation states of particular phase. Vertical reservoir flows cause the re-equilibration due to instable contact with sources of controlling assemblage components, which is of relative effect on thermodynamic activity stability of the phases. Irregularity in shape and wide δT_{TSI} dispersion is a sign of dilution, responding to destruction of initial fluid composition. Reservoir boiling or (uncontrolled) borehole flashing gives TSI curves shifting in total saturation, maintaining one of saturation extremes or distributed irregularly. Yet

vertical resolution in TSI is a function of SI extremes of selected assemblage (Fričovský, 2014).

Quadratic function (Eq. 57) respects possible analytical and sampling errors disqualifying possibility of $TSI = 0$. The type curve analysis is used rather to evaluate a possible degassing effect as it should be a curve of closest approach to the $TSI = 0$ line and with one minima and more-or-less concave shape in a solid (e.g. Cooper et al., 2013). Because of squared term, the curve reflects SI by changing a shape or introducing several local extremes.

Upscaling and weighting factors

By definition, SI is a function of reaction stoichiometry (Smith et al., 2012). Both transcriptions defining the total saturation index (Eq. 57–58) are weighted by a factors (wt_i) ensuring all assemblage minerals are considered equally and a TSI (or TSI temperature) estimation is not skewed by stoichiometry (Neupane et al., 2014). Even numerous methods to upscale the total saturation index exist, such as thermodynamic activity, inverse of reservoir alteration frequency etc., the upscaling by thermodynamic components defined in (Cooper et al., 2013) entering reactions is the one used most frequently, still developing continuously (Cannon et al., 2014; Cooper et al., 2013; Neupane et al., 2014).

Application: Weighting factors and assemblage definition

We preferred the approach based on number of thermodynamic components. In that principle, the factor is derived from number of thermodynamic components for i -th mineral and a number of times each component enters dissolution or (equilibrium) ion exchange reaction (Tab. 3) of the i -th solid. This scales up the weighting factors as 1 mole of ion is added into solution (Cooper et al., 2013).

To ensure analysis coincidence, we kept the assemblage selected arbitrary and described in complex equilibrium analysis (Tab. 2). A reason for that is in a simple fact that the use of different assemblage would account different solids, skewing onward interpretations as referring to different equilibration or alteration conditions.

Application: TSI temperature calculation

Even there is no boiling, we used linear and quadratic total saturation index models for temperature estimation. TSI temperature is inferred by $TSI = 0$ interception for linear, and $TSI = 0$ convergence for quadratic approach.

Only two samples (Tab. 4) remained oversaturated through the entire interval ($T = 20$ – 100 °C), that disqualified them out of representative temperature assessment immediately. The rest has been determined to equilibrate

Tab. 3
Weighted factors calculation for normalized assemblage

Mineral	Thermodynamic components	wt_i
Albite	1* Al, 1* Na, 3* SiO ₂	5
Adularia	1* Al, 1* K, 3* SiO ₂	5
Anhydrite	1* Ca, 1* SO ₄	2
Calcite	1* Ca, 1* CO ₃	2
Ca-mont.	1* Ca, 2* Al, 2* Mg, 2* OH, 4* SiO ₂	11
Dolomite	1* Ca, 1* Mg, 2* CO ₃	4
Fluorite	1* Ca, 2* F	3
Gibbsite	1* Al, 3* OH	4
Chalcedony	1* SiO ₂	1
Chlorite	2* Al, 3* SiO ₂ , 8* OH, 5* Mg, 5* Fe	23
Illite	1* K, 2* Al, 3* SiO ₂ , 2* OH	8
Kaolinite	2* Al, 4* OH, 4* SiO ₂	8
K-mont.	1* K, 2* Al, 2* Mg, 2* OH, 4* SiO ₂	11
Mg-mont.	1* Ca, 1* Na, 2* Al, 2* Mg, 4* SiO ₂ , 2* OH	12
Muscovite	1* K, 3* Al, 3* SiO ₂	7
Na-mont.	1* Na, 2* Al, 2* Mg, 2* OH, 4* SiO ₂	11
Am. Pyr.	1* Fe, 2* H ₂ S	2
Talc	1* Mg, 2* OH, 4* SiO ₂	7

within $T_{TSI} = 72$ – 94 °C. Low coefficient of variation ($c_v = 0.682$) supports the median temperature $T_{TSI} = 79$ °C as representative, that corresponds to lower half of the reservoir (Fig. 25).




Fig. 25. TSI: linear approach.

Same two samples have been eliminated for squared T_{TSI} analysis (Fig. 26). The method is more sensitive to local SI minima and maxima extremes. In total, 18 samples converged closest to $TSI = 0$ at $T_{TSI} = 69\text{--}89\text{ }^{\circ}\text{C}$ (Tab. 4), with a median of $T_{TSI} = 74\text{ }^{\circ}\text{C}$. The difference between both is $\delta T_{TSI} = 5\text{ }^{\circ}\text{C}$, which is less than 10 % of reservoir temperature coverage interval, forth it is possible to refer the difference to extremes individual phases attain. Henceforth the temperature corresponds to middle/lower half part of the reservoir again.




Fig. 26. TSI: quadratic approach.

Application: TSI type curve analysis

Two groups of linear TSI are identified (Fig. 25). Six samples record linear, or quasi linear shape, intercepting the $TSI = 0$ line with $5\text{ }^{\circ}\text{C}$ dispersion ($T_{TSI} = 72\text{--}77\text{ }^{\circ}\text{C}$). The rest equilibrates at $T_{TSI} = 73\text{--}94\text{ }^{\circ}\text{C}$, however, records a break in TSI increase at $T_{TSI} = 35\text{--}50\text{ }^{\circ}\text{C}$, out of reservoir conditions. Yet no break is observed within the reservoir temperature interval. Moreover, three samples plot some re-equilibration potential at $T = 23, 30$ and $31\text{ }^{\circ}\text{C}$.

The reservoir temperature difference between a top and a base equals $\delta T_{res} = 40\text{ }^{\circ}\text{C}$. If we take the entire group, the $\delta T_{TSI} = 22\text{ }^{\circ}\text{C}$ ($> 50\%$), that is over tolerance. Uniform run of TSI functions contrast to high δT_{TSI} is indicative of reservoir immaturity. Either a minor change in concentration of major constituents disperses curves out of median TSI. This is proportional to instability in dissolution of macro-components, whether it may be due to quasi-isochemical mixing at reservoir intake (= reservoir is sensitive to re-charge dynamics) or due to intraformation mixing between residential and vertically filtrating waters (= reservoir is sensitive to rate of convection). Stability is indicative of no boiling environment.

The use of squared function determined a convex (linear TSI) and sinusoidal (linear with break below $50\text{ }^{\circ}\text{C}$) lines. The number of samples per group equals to distribution of linear analysis (Tab. 4). The break, which gives the sinusoidal shape realizes at $T_{TSI} < 50\text{ }^{\circ}\text{C}$, similar to linear

Tab. 4
Multicomponent equilibrium comparison

Approach	z21/2/1987	z4/3/1987a	z4/3/1987b	z5/3/1987	z6/3/1987a	z6/3/1987b	z7/3/1987	z22/3/1987	z31/3/1987	z2/4/1987	z3/4/1987	z20/2/1992	z12/11/2007	z17/12/2008	z18/2/2008	z26/2/2009	z30/3/2011	z13/4/2011	f15/3/2011	f30/3/2011
CEQ	56	65	74	–	72	79	–	67	70	65	71	63	65	68	56	61	60	62	75	75
TSI (L)	72	76	94	–	81	86	–	78	80	80	80	77	74	80	74	76	77	77	86	86
TSI (SQ)	70	71	89	–	81	89	–	73	79	80	80	70	70	75	69	70	69	70	80	80
RMED	80	50	30	–	60	80	70	80	–	80	–	30	50	80	50	60	60	–	80	80
n-RMED	2	7	7	–	4	6	1	6	–	2	–	2	8	7	6	10	10	–	13	9
L-TSI T	L	A	A	A	A	A	A	A	A	A	A	L	A	A	L	L	L	L	A	A
SQ-TSI T	V	S	S	S	S	S	S	S	S	S	S	V	S	S	V	V	V	V	S	S

Explanations: n-RMED – number of equilibrated phases; L-TSI T – linear type curve; SQ-TSI T – squared type curve; L – linear shape; A – concave (break) shape; S – sinusoidal shape; V – convex shape

analysis. The variation in convergence $\delta T_{\text{TSI}} = 20^\circ\text{C}$ narrows compared to linear analysis, however, still equals 50 % of reservoir temperature interval. We add the difference is just as sensibility of the method increases towards saturation extremes, meanwhile supporting previous interpretations.

Clustered saturation indexes approach (RMED)

Optimization procedure

The MEG analysis and reservoir temperature estimation is clearly a function of selected mineral assemblage, either set arbitrary, or by numerical optimization (Spycher et al., 2014). Saturation indexes of a wide assemblage are normalized by a weighting factor (e.g. Cooper et al., 2013) and calculated against desired temperature interval. First iteration eliminates the phases which SI does not converge to 0.05 in absolute over the analysed range. Then, a median of SI (RMED) is calculated at each temperature step, normalized by a factor of 1.2. The suite to calculate temperature of overall equilibrium is narrowed by ignoring those solids violating $SI \leq 1.2\text{RMED}$. This eliminates skewing of TSI calculation, finally the reservoir temperature (T_{RMED}) is inferred from that of minimum median of absolute SI, yielded by residual phases – suite (Spycher et al., 2014).

Application: Assemblage upscaling

Similar to previous methods (CEQ, TSI), we decided to analyse the primary-alteration-hypothetical assemblage selected before to ensure conceptual homogeneity for onward correlation and analysis. Saturation indexes are, analogously, weighted as according to Tab. 3. The assemblage has been eliminated with calcite and muscovite, both not reaching $-0.05 < SI < 0.05$ throughout the reservoir temperature interval. Later upscaling iterations have been conducted individually. The Ca- and Mg- montmorillonites were disqualified as most (Tab. 5), whilst chlorite, chalcedony and gibbsite were involved in each calculation.

Application: Upscaling implicatons

Elimination of calcite is more or less expected, as thermal waters simply must be oversaturated with calcite (Fig. 19) even before intake into the reservoir zone. The oversaturation to calcite, simultaneous to a permanent contact with a source rock is then one source of excess CO_2 in deep reservoir waters. As thermal waters are hosted in impure limestones and dolomites (and their transient forms) at low enthalpy conditions, the oversaturation with muscovite (Fig. 15), restraining its reaction activity, allows us to assume the LT alteration of siliceous (detritic) dolomites,

which does not involve the metamorphic ion-exchange reactions, but activates talc, produces Mg-calcite and increases CO_2 . Muscovite oversaturation consumes the SiO_2 and available Mg^{2+} into chlorite alteration. Yet chlorite and talc record confident equilibration. Their declining saturation tendency (Fig. 24) with decreasing temperature aids the process. With the use of this approach, the dolomite equilibrates well at $T < 50^\circ\text{C}$ (80 % of equilibration records of $SI_w \leq \text{RMED} \cdot 1.2$), which corresponds to initial conception and assumption on some Mg-calcite precipitation while fluid experiences lateral leaking or vertical evasion, maintaining high CO_2 level at reservoir outlet, restraining balanced Na/K alteration (Fričovský and Tometz, 2014). Indeed, even with upscaled saturation, albite and dissociated Na-silicates are rather consumed to form illite or maintain muscovite uptorn, at least upon reservoir outlet.

Tab. 5
RMED summary table

Phase	Eliminations	Preferential RMEDx1.2 record	Preferential T_{RMED} contribution
Adularia	3	50	50
Albite	6	40	30
Anhydrite	18	20	20
Calcite	20	–	–
Ca-mont.	13	80	80
Dolomite	6	40	50
Fluorite	5	90	80
Am. pyrite	4	50	60
Gibbsite	0	60	60
Chalcedony	0	70	80
Chlorite	0	80	80
Illite	1	50	50
Kaolinite	1	80	80
K-mont.	3	70	80
Mg-mont.	11	80	80
Na-mont.	2	70	80
Talc	1	60	60
Muscovite	20	–	–

The other reason for inequilibrium of albite–adularia system is relative good correlation of Na- and K- montmorillonites (or their activity coefficients), optionally controlling reservoir Na^+ and K^+ hydrolysis. Some loss in

CO₂, due to the precipitation of calcite and Mg-calcite, or because of consumption for talc activation, can not compensate its oversaturation referring to the Na/K system. Obviously, the rate of Na/K immaturity is a function of adularia-albite inequilibrium, proportional to excess CO₂ level, supporting an idea of immaturity relation to dissociation of carbonates, but to low grade alteration of detritic (or impure) dolomites as well. Carbonates are deficient in alkalis, forth the metamorphosis intensity depends on a water-rock contact with separate horizons containing siliciclastics. Instability of the water-source interaction relates then to variable vertical filtration. The deep reservoir is off any conditions for intrinsic convection. However, because of unconform base overheating, there are separate convection cells formed within, even not covering the entire effective profile. It is just obvious then that all reservoir reactions, whether do or do not producing CO₂ are a function of induced convection and so to the mass movement controlling final equilibration or re-equilibration. Further on, the rate of immaturity and inequilibrium in Na/K relates explicitly to the rate of reservoir convection extension (vertical thickness, lateral impact), duration (vertical filtration velocity), and intensity (bulk mass movement).

Application: RMED temperature (T_{RMED}) calculation

Because of “hands on” elimination, we carried computations in 10 °C intervals, forth the real extension of inferred temperature is of $T_{\text{RMED}} \pm 5$ °C coverage. In total, 16

samples equilibrated according to RMED approach. The inferred temperature varies in $T_{\text{RMED}} \approx 30 \pm 5$ °C – 80 ± 5 °C interval, with medium (representative) temperature of reservoir equilibration at $T_{\text{RMED}} = 65$ °C (the average $T_{\text{RMED}} = 64$ °C). This is fairly similar to temperature of complex equilibration ($T_{\text{CEQ}} = 66$ °C), but is still less than that of total saturation index ($T_{\text{TSI}} = 79$ °C). The T_{RMED} versus T_{CEQ} similarity is fairly coincidental. With RMED, both most samples equilibrate and most equilibration records are observed for $T_{\text{RMED}} = 80 \pm 5$ °C (Tab. 6), which is, however, close to the temperature inferred by TSI. The consistency here is due to idem upscaling (weighting factors) application (Tab. 4).

Ternary chloride mixing model

Use of thermodynamic systems based on two or three components has been adopted within the Na-K-Mg (Giggenbach, 1988) and K-Mg-Ca (Giggenbach, 1986; Giggenbach, 1988) geothermometers. Both do, however, target the equilibrium temperature of initial spring and fail to provide relevant results while analysing mixed samples. Such samples are, however, rather scarce.

The ternary mixing model geothermometer expects the sampled water is a solution of parental and shallow thermal or cold groundwater, which is of some relation to aqueous chloride (Cioni et al., 1992). Then, constituents involved in particular geothermometers are regressed to the chloride. If their concentrations are fixed by primary

Tab. 6
 T_{RMED} summary table

Temperature (°C)	Adularia	Albite	Anhydrite	Calcite	Ca-mont.	Dolomite	Fluorite	Am. pyrite	Gibbsite	Chalcedony	Chlorite	Illite	Kaolinite	K-mont.	Mg-mont.	Na-mont.	Talc	Muscovite	n-RMED	T_{RMED} contribution records
100	–	–	–	–	–	–	1	–	–	–	–	–	–	–	–	–	–	–	1	1
90	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	0	0
80	2	–	–	–	3	–	5	2	3	6	7	1	7	3	5	3	1	–	13	48
70	–	–	–	–	–	–	–	–	1	–	–	–	–	–	–	–	–	–	1	1
60	3	1	–	–	–	1	–	3	4	2	2	3	2	2	–	2	2	–	12	27
50	5	4	1	–	–	2	1	1	4	1	2	4	–	–	–	–	1	–	11	26
40	–	5	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	1	5
30	–	–	–	–	–	–	–	–	–	–	1	–	–	–	–	–	–	–	1	1
20	1	–	2	–	–	1	–	1	–	–	–	–	–	–	–	1	–	–	5	6

water–cold water mixing, their concentrations (cX) should fit a straight line and a simple linear regression should be obtained (Eq. 53).

$$cX = a \cdot cCl \pm b \quad (\text{Eq. 53})$$

where “a” and “b” are coefficients of linear regression, while concentrations are in various units.

Still, if there is not any linearity between aqueous concentrations and mixed water with dilutant, other forms may occur. This is for example a case of irregular and unbalanced mixing. Linearity destruction may be of supplementary proof on no-mixing relations between selected geothermal–coldwater pairs.

Regressed concentrations are inserted into geothermometry functions (Antroicchia et al., 1985; Marini, 2004). For these, at least necessary conditions of conceptual model, which is in control of particular geothermometer, must be met. Besides a silica geothermometer, various validated, and reasonable cation geothermometers may be included, such as that of Na/K (Eq. 55) and K-Mg (Eq. 56) as it is examined in (Chiodini et al., 1996; Marini, 2004):

$$T_{SiO_2} (^{\circ}C) = \frac{X}{Y - \log(a \cdot Cl \pm b)} - 273.15 \quad (\text{Eq. 54})$$

$$T_{Na/K} (^{\circ}C) = \frac{X}{Y \pm \log(a \cdot Cl \pm b) - \log(a \cdot Cl \pm b)} \quad (\text{Eq. 55})$$

$$T_{K/Mg} (^{\circ}C) = \frac{X}{Y - Z \cdot \log(a \cdot Cl \pm b) - W \cdot \log(a \cdot Cl \pm b)} \quad (\text{Eq. 56})$$

where “X”, “Y”, and “Z” are generalized constants for SiO_2 , Na/K and K-Mg geothermometers derived by different authors and Cl[−] concentrations are in attributable units.

In next step, all temperatures are calculated for increasing chloride and plotted against it (Marini, 2004). If the fluid is in overall equilibrium, all functions converge some point, from that drawing a horizontal line onto Y axis gives an estimate of parental (primary) thermal water component, whilst a vertical line onto X axis gives an initial, reservoir chloride concentration at overall equilibrium (Chiodini et al., 1996). Admittedly, the given temperature must be validated due to a conceptual model of a system. If ternary function does not give any interception point, it is a sign on disequilibrium. However, geothermometry functions may intercept at a temperature that is far of reasonable, which must be interpreted immediately. Most probably, discrepancy in temperature to that expected in the reservoir is of error in magnitude of fluids disequilibrium to one or more selected geothermometers.

A key advantage of a model is that functions of solute geothermometry may be, if validated, substituted by other geothermometers (Antroicchia et al., 1985), such as isotopes or

multicomponent equilibrium functions (Fričovský and Tometz, 2014), or involving a conservation enthalpy function (Cioni et al., 1992).

Application: Ternary combination approach

The selection of samples requires, at least, some record on affinity to parental reservoir fluid, besides the mandatory condition of mixing character. In local conditions, the parental endmember is considered the DRTE. Waters sampled at VSH-1 has been assigned as mixture according to geothermometry (this paper) and geochemical indicators (e.g. Fričovský and Tometz, 2013; Fričovský et al., 2015). Groundwaters, representing a dilutant, must be facially relative to the site. We used cold springs from southern margin of the Liptov Basin. The most important is, however, a controlling thermodynamic system selection, represented by most reliable geothermometers. To meet this condition, we used chalcedony geothermometer GSC3 (Fournier, 1991), in combination with GKM3 K²/Mg model by (Giggenbach, 1988), and GLM Li/Mg^{0.5} function (Kharaka and Mariner, 1989). Coupling of these refers to chalcedony control on dissolved SiO_2 in thermal waters, which try to equilibrate with K-Mg silicates by adapting the CO_2 cycle during lateral or vertical evasion from the reservoir, meanwhile conserving Li⁺ concentration restraining its reaction with clay minerals.

Application: Deep reservoir analysis TCLM (with VSH-1 samples)

The TCLM plot of VSH-1 samples over increase in chloride shows no ternary convergence, either because of secondary dilution or involvement of more than one thermal water component in the mixture (Fig. 27). The SiO_2 – Li/Mg pair intercepts at $T_{TCLM} \approx 70$ °C. Low K²/Mg ratio slips the GKM3 evolution with chloride down, converging with GSC3 at $T_{TCLM} \approx 30$ °C. The dispersion is rather due to high Mg^{2+} and some K⁺ deficiency. Supposed magnesium consumption by activation of talc, chlorite alteration and Mg-calcite precipitation must be then compensated by additional Mg^{2+} from dilution. The GSC3/GLM interception is rather a sign of lithium concentration conservation, with minor enrichment only (if any). Definitely, the lithium is not added into solution by the ion exchange reactions.

Let that be the conservation of Li⁺ from parental (DRTE) reservoir. The lithium clays are not able to form, as SiO_2 and Al^{3+} is consumed in Na-silicates transformation to K-micas (at least in terms of thermodynamic activities). Then, the Li⁺ addition should be by dilution (S-CGW facies form in siliciclastics rich IWCP) or mixing with SRTW, at permanent contacts with IWCP or Jurassic/Mid Cretaceous succession. Either way, lithium clays may be conserved, dissolved, but do not enter the ion exchange reactions.




Fig. 27. TCLM analysis for DRTE/VSH-1 facies relation.




Fig. 28. TCLM analysis for DRTE/SRTW facies relation.

Application: Deep reservoir analysis (with BEH-1 and FBe-1 samples)

To verify secondary origin of the shallow reservoir, assuming mixing of parental DRTE waters with recharged (residential) waters, we let the BEH-1 and FBe-1 waters cool with springs from the Kľačianka valley. Excess SiO_2 in combination with low K/Mg ratio restrain the GSC3/GKM3 to intercept (Fig. 28). If we refuse some amorphous silica solubility activation at lower temperatures, the additional SiO_2 source must come from sufficient source, which is rather the IWCP (Borové Fm.), or Jurassic–Mid Cretaceous succession at a base.

As we have already discussed above, other SiO_2 source is fair to assume in the IWCP formations, that distinguished both subfacies in the SRTW (SiO_2 deficient BEH-1 and SiO_2 sufficient FBe-1). Neither Mg^{2+} deposition due to dolomite precipitation, nor its consumption in talc acti-

vation balances the Mg^{2+} enrichment by possible mixing (dilution) within shallow reservoir, or its simple increase by the local dissolution of dolomite from the host rock (expecting DRTE waters to get undersaturated to dolomite). All processes restrain equilibrium attainment to the K/Mg . The Li^+ concentration within shallow reservoir is more or less constant in time, compared to the deep reservoir. Convergence of GSC3/GLM gives a temperature estimate of $T_{\text{TCLM}} \approx 50^\circ\text{C}$, which corresponds to deep reservoir evasion temperature and may be a sign of conservation of lithium during vertical evasion, as with decreasing temperature, the activity to dissociate lithium clays decreases rapidly.

SUMMARY – CONCEPTUAL SITE MODEL UPDATE

First CSM models were developed in the late 1970s to conclude research of mineral in early (Zbořil et al., 1972; Franko et al., 1974) and shallow thermal waters of the Liptov Basin in a latter stage (Franko et al., 1979; Zembjak et al., 1982). A consensus on open hydrogeological character identified central discharge zone by travertine accumulations analysis in the Bešeňová vicinity (Franko et al., 1974), assigning the dome formations to thermal waters stored in deep reservoirs below. Drilling campaigns (Franko et al., 1979; Fendek et al., 1988) revealed stratified character of the accumulation zone. Numerical modelling (Fendek et al., 1988) and spatial piezometry analysis (Remšík et al., 1993) speculated the thermal waters to intake deep in permeable structures from the east (Liptovská Mara Depression), and leak laterally towards west after transition and some residency in central parts of the system. The open character of the structure has been proven continuously in time by total discharge analysis (Remšík et al., 1998), pumping tests (Vandrová et al., 2011), lumped parameter modelling (Fendek et al., 2013), or reservoir response analysis (Fričovský et al., 2014).

Geochemical methods paid favour onto geochemical origin and mineralization type (Franko et al., 1979; Fendek et al., 1988; Bodiš and Borosová, 1996), not much clearing up a contention on primary recharge zone position between the Lúčky area or the northern slopes of the Nízke Tatry Mts. The latter has been evidenced by isotope elevation analysis (Franko, 2002a, b). Application of boiling and mixing models (Fričovský and Tometz, 2013) provided more support to the assumption, emplacing the recharge zone to the Kľačianka valley area, that is in confidence with hydraulic (Fendek et al., 1988; Fendek et al., 2013), stationary temperature (Fričovský, 2014), dynamic hydrogeothermal (Fričovský et al., 2014) and reservoir response (Fričovský et al., 2015) models.

Application of several geothermometry techniques presented an ample scale of models and hypotheses, which is a must to interpret complexly for a CSM construction. In past, geothermometry has not gained a solid tribute, relieving importance of saturation (equilibrium) analysis methods by a few constraints on its inadequacy in local, low temperature, carbonate environment. An initial con-

ceptual model of simple infiltration, downflow transition and accumulation, with later westward expelling should then be, modified, as both, macrochemistry (boiling and mixing models) and geothermometry, aided with thermodynamic microanalysis, give some evidences on several mixing episodes, the primary model does not account.

Infiltration zone model

It is a must to pronounce the infiltration, based on spatial analogy and lithological homogeneity in seeping conditions, as there is no well available. Some indices on geochemistry of waters in early transition stages may be derived of springs and shallow wells in the valley, however, early infiltration stages taking part in prior must be assumed by lithology in a major.

We use the Rudolf well, LM springs (warm) and adjacent cold springs (e.g. Bočný prameň, pod Bielym etc.) to derive a model of the infiltration zone (Fig. 29). Previous studies (e.g. Franko, 1984; Fričovský and Tometz, 2013) localized the infiltration zone to the Kľačianka valley. The Rudolf and LM springs are analogized to represent early infiltration/transition facies, recording transient Ca-Mg-HCO₃-SO₄ type. By lithology, the only possible source of sulphate acquirement is the Late Triassic, found in the hydrogeological massif of the Nízke Tatry Mts. Application of Mg-correction (Fournier and Potter, 1979) to the T_{NKC} (Fournier and Truesdell, 1973) yields overrated temperatures, implying excess Mg²⁺, that can not simply be adjusted to oversaturation with Mg-carbonates. This is, moreover, accented by positive T_{NKC} – pH correlation for springs pointing to various Mg²⁺ and Ca²⁺ source.

We prefer hypothesis based on two flow channels; a deeper one off the Nízke Tatry Mts. in deep and relatively slow circulation controlled by fissured and karst-fissured permeability, and the shallow one associated with fast downflows along a fissure swarms at the periphery of the basin. Both mix together at the periphery in TCM (thermal-cold-mixing) regime, satisfying excess Mg²⁺ objection and relatively stable Na/K ratio compared to cold springs by reciprocal leaching of Na/K silicates, not destructed by CO₂ saturation at this stage.

By the excess Mg²⁺, the mixing rate is over 50 %, which is, in fact, a partial dilution. The TCM is then objected to realize prior differentiation for downflow (transition-accumulation donation) and upflow (springs upwelling) regime. Indeed, there is a difference between cold and warm springs in pH. Then, the mixing must realize in CHN Mid Triassic carbonates or KNA Jr-Cr2 succession (satisfying stable Na/K ratio).

Transition zone model

The entire structure of the Bešeňová elevation is considered open in hydrogeological regime referring to both reservoir zones. To satisfy this objection, recharge must

be supplied in at least two effective profiles, separated spatially and horizontally. This calls the transition zone to split for a shallow and deep channels, obviously just after a short period of conjoint submersion off the infiltration zone.

Shallow transition/shallow recharge channel

Comparison of facies defined by wells and springs from the infiltration zone with the shallow reservoir (BEH-1, FBe-1): At an early stage, the transition realizes in weathered or fissured upper zone of the Jurassic–Mid Cretaceous KNA horizon, involving dissolution of limestones primary. Increase in Na/K reflects partial temperature build-up and decrease in filtration velocity prior effective inflow into shallow reservoir that allows preferential Na⁺ leaching out of Jr–Cr2 bedrock. Releasing of K⁺ from silicates in the IWCP is obviously less intense. A slow transition in silica metastability from the LT opal to chalcedony regime gives a supportive proof onto decrease in filtration velocity with depth and distance from the infiltration zone, maintaining a contact with silicates-dominated environment (Fig. 29). Still, there is no sign of equilibrium attainment with carbonates, which dissolution alternates spatially. While dissolution of limestones realizes in early shallow transition stage (increase in Ca/Mg ratio) within Jurassic–Mid Cretaceous succession, this is soon altered by dissolution of impure dolomites from the Borové Formation and the CHN Mid Triassic dolomites (decrease in Ca/Mg ratio).

Deep transition/deep recharge channel

A model of deep transition is interpreted by comparing infiltration zone and deep reservoir facies (FGTB-1, ZGL-1), analogously to previous. Obviously, Na-silicates are dissociated from the Jurassic–Mid Cretaceous horizon preferentially along with calcite. By decrease in absolute CO₂ content between infiltration and accumulation/reservoir production zone, along with a better fit of pCO₂ related to dissociation of Ca-silicates (hypothetical phase), derived of the K-Mg-Ca geoinicator (Fig. 18) saturation of submerging thermal waters with calcite must take part before effective inflow to the production zone (CO₂ is consumed in activity and saturation variation in the deep reservoir variation). Preferential (primary) saturation with respect to calcite during downflow projects into equilibration with an average crustal rock by acquiring Ca²⁺ on the Na-K-Mg-Ca plot (Fig. 21). Qualitative Na/K geothermometry analysis (Figs. 10 and 11) defines the Na⁺ and K⁺ addition of preferential rock leaching origin. Decrease in Na/K ratio with the depth and temperature reflects aggressive CO₂ accretion during submersion (or at least its conservation from the infiltration zone), supplied through a

contact with carbonates of variable impurity, progressively setting on a modification or restriction of easily leached albite into K-silicates.

An initial, transient type chemistry ($\text{Ca-Mg-HCO}_3\text{-SO}_4$) may be partially altered towards $\text{Ca-Mg-SO}_4\text{-HCO}_3$, acquiring some SO_4^{2-} at a contact with Late Triassic horizon and Carpathian Keuper shales and slates during down-flow, modifying peripheral character of recharges towards immature. Attaining, or conserving the transient character supports then an assumption of quasi-isochemical mixing with residential thermal waters at an effective inflow towards production zone, implied e.g. by high $\delta T_{\text{TSI}} = 0$ dispersion (Figs. 25 and 26).

A presented model is based on geometry of the structure and interpolation of records between both production zones and infiltration area. Probability of identified processes is than a matter of uncertainties of the CSM.

Deep reservoir (accumulation/production zone) model

CSM for the deep reservoir (Fig. 30), referred as to Mid Triassic carbonates of the Krížna Nappe, is constructed as based on results of geothermometry and essential mixing models applied to samples of the ZGL-1 and FGTB-1 well, with simultaneous comparison with the infiltration zone to respect a conceptual histories of samples.

The deep reservoir is all but homogeneous referring to its vertical profile (already documented on both production wells), hydrogeochemistry and intraformational processes. The effective inflow realizes most probably towards base. Because of similarity in a macroscale (deep reservoir basal facies and recharges are of $\text{Ca-Mg-HCO}_3\text{-SO}_4$ to Ca-Mg-

$\text{SO}_4\text{-HCO}_3$), the inflow is most probably of quasi iso-chemical TTM (thermal-thermal-mixing) regime. However, the velocity of submerging filtration controls initial intake geochemistry, that is definitely not constant. A sensitivity to recharges is then documented through a dispersion of $\delta T_{\text{TSI}} = 0 > 50\%$ of the reservoir range (Figs. 25 and 26). Apparent similarity in TDS content and geochemistry type plots as quasi-logarithmic distribution of samples on the silica – total carbonate model (Fig. 7), where the organization of the population follows the $\text{SiO}_2\text{ – CO}_3$ equilibrium function (by Arnórsson, 1985) trend, departed towards no-boiling region due to general silica lows.

Discussion on reservoir convection

In previous, the studies took the deep reservoir as a relatively homogeneous aquiferous body of “simple” tectonics bound vertical evasion towards surface, referring to the transition/accumulation zone. Yet already (Fričovský et al., 2014) presented an idea of minor convection as based on reservoir base overheating analysis, inferring formation of isolated convection cells of various spatial extension. Application of the reservoir geothermometry and thermodynamics (this paper) reveals, however, several more indices to combine with initial hypothesis.

A domestic convention on silica geothermometry inadequacy gainsays its principles in essence. Let the reservoir carbonates be impure (siliciclastics intercalations, beds, lenses, grains) and limited in spatial extension. A variation in absolute SiO_2 (Fig. 4) gives a fair evidence on instable contact of samples with its source (stable accommodation




Fig. 29. Infiltration and transition zone CSM. For its localization see Fig. 1.

in reservoir positions would result just in SiO_2 lows), projected consequently onto quantitative temperature estimations (Appendix D).

A most vivid evidence on some vertical mass movement in the reservoir is provided by a use of K-Mg-Ca geoinicator (Fig. 18) or geobarometer (Giggenbach, 1988). The model reveals the reservoir partial CO_2 off a range to induce a calcite precipitation, thus its saturation realizes already in downflow to the production zone. In details, variations in $p\text{CO}_2$ are equal to differences in the L_{kc} and L_{km} functions (Eq. 47). This is because reservoir Mg^{2+} varies (e.g. according to Mg- SiO_2 alteration system analysis, Fig. 13) along with the Mg/Ca ratio, resultant to instable saturation of thermal waters with respect to carbonates during a consistent contact with a source rock. The variation must than be an effect of even minor changes in temperature and pressure, possible only through intra-formation vertical movements.

In combination to base overheating implications, thermal waters partially enter separated and spatially limited, insulated convection cells, however, through the entire vertical profile of the body because of building up a density and p-T gradient between its top and a base, as the latter is heated unconformly.

Discussion on carbonates equilibria

The entire geothermal structure is predominantly hosted in carbonates. These are involved in the infiltration and transition zone, even variably impure, or as a compound of calcareous pelites and sandstones. In reservoir positions,

dolomites prevail over limestones, along with the occurrence of transient and impure or intercalated forms.

Application of the K-Mg-Ca geoinicator (K-Ca- $p\text{CO}_2$ geobarometer) baffles expectations of CO_2 regime by assigning the CO_2 fugacity to Ca-alumosilicates hydrolysis. This is not possible in local conditions. In other words, the notice points to saturation of thermal waters with calcite (Fig. 19) prior effective intake into reservoir (production zone) position. Then, because thermal waters become saturated with calcite, samples organize towards an equilibrium with a wallrock (Na-K-Mg-Ca model, Fig. 21) by preferential dissolution of dolomite, increasing the Mg/Ca ratio continuously. However, thermal waters become over-saturated with dolomites soon, maintaining a contact with a source (Fig. 20). Onset of induced convection causes unit variation in saturation of carbonates. Because of a stable contact with a source rock, the unit of carbonate which has precipitated due to change in temperature and pressure in the vertical upflow is subsequently replaced by a unit dissolved in a downflow. This produces an excess CO_2 towards reservoir base, controlling immaturity of thermal waters by restraining a principal, Na-K, K-Mg and Na-Mg thermodynamic subsystems equilibrium attainment.

Discussion on silica regime

A silica regime has not been subjected to detailed analysis in previous. At deep reservoir conditions, use of saturation stability analysis (Fig. 4) records unbalanced distribution of records between conductive quartz solubility, chalcedony and LT opal polymorphs. At a given tem-




Fig. 30. Deep reservoir mass movements controls.

perature range, the variation is rather assigned to uneven silica content due to instable contact with its source. At deep reservoir conditions, chalcedony becomes metastable and in control on reservoir aqueous silica content (Fig. 5), attaining equilibration with a median of $T_{SI=0} \approx 67$ °C. Application of silica geothermometers yielded reasonable results for chalcedony as well with median for conductive cooling models of $T \approx 75$ °C. We assume the difference is not only a record of various sensitivity of functions to the dissolved silica content, but due to the instable contact of samples with a silica source and reaction time as well.

If the chalcedony becomes saturated, it is possible that some dissociated H_4SiO_4 becomes available for onward alteration of silicates. In vertical filtration regime, only a minor part of chalcedony can be deposited by a drop in p-T conditions, rather more is incorporated into Na/K silicates alteration and chlorite-talc activation.

Contribution to deep reservoir hydrochemical regime

Reservoir hydrochemical regime is controlled usually by primary lithology, temperature, pressure and residential period of thermal waters in the structure. In carbonates-hosted environment, a fluid saturation and available CO_2 plays a crucial role. Thermal waters of the deep reservoir are deficient in absolute Na^+ and K^+ content, expectable under insufficient source conditions. Compared to infiltration zone, the Na/K, K/Ca and K/Mg ratios decrease dramatically by conservation of absolute alkali concentrations and increase in Ca^{2+} and Mg^{2+} .

Application of Na/K qualitative geothermometry records samples organized in the rock dissolution region (Fig. 10) away of any alteration equilibration (Fig. 9), causing quantitative estimates (Appendix E) overrated. Under high CO_2 content, the inequilibrium reflects modification of Na-silicates into K-silicates, restraining albite equilibration while increasing adularia, illite and muscovite activity. While complex equilibrium of albite (9/18 samples) is roughly of $T_{CEQ} = 30$ –46 °C, adularia equilibrates at $T_{CEQ} = 40$ –65 °C.

The immaturity of reservoir fluids ($MI = -0.1$ –0.4; Fig. 16) is promoted by easier activation of Na- and K- montmorillonites (Tabs. 5 and 6), reaching equilibration (even in a magnitude of equilibration activity) with the reservoir fluid. Excess CO_2 and muscovite activity (Fig. 15) allows LT alteration of impure siliceous reservoir dolomites not involving metamorphic ion exchange reactions, but with LT talc activation, secondary Mg-calcite, calcite and CO_2 production. The incoming CO_2 is available to contribute further on a unit-scaled calcite – dolomite – Mg-calcite dissolution/precipitation cycle during reservoir convection.

These processes are controlled by equilibration temperature and depth. The talc equilibrates at $T_{CEQ} = 65$ –70

°C and $T_{RMED} = 60$ °C preferentially (Tabs. 2 and 5). The chlorite records a balance in equilibration constant and thermodynamic activity at $T_{CEQ} = 70$ –75 °C and $T_{RMED} = 80$ °C with some preference. Then, the excess Mg^{2+} in the reservoir is assumed to persist up to 50–60 °C, which corresponds to upper reservoir zones, maintaining low K/Mg and Na/Mg ratios (Fig. 31).

Both carbonates tend to precipitate in evasion regime, simultaneously allowing some equilibration of alkalis with consumed CO_2 as a rate of Na-silicates modification decreases proportionally. Under given trend between DRTE and SRTW (1) at Na-K-Mg-Ca geoinicator (Fig. 21), Mg-calcite is lost more intensively as the fluid breaks a contact with a source rock.

Discussion on deep reservoir adiabatic boiling

A boiling as one of crucial reservoir dynamics has been studied simultaneously with qualitative geothermometry analysis. A basis for this study was a discussion of boiling and mixing models (Fričovský and Tometz, 2013; Fričovský et al., 2015). After carried qualitative geothermometry analysis, we conclude that the boiling is definitely disqualified as:

- samples plot deep in the non-boiling region (Fig. 7) at the silica–total carbonate mixing model (missing any trend organization towards a boiling line)
- enthalpy and silica fraction functions intercept (Fig. 8) implying rather polystadial mixing instead of steam-loss induced silica deposition
- saturation curves (Figs. 5, 15, 19, 20 and 24) of particular solids are conserved in shape, not varying saturation states transitions
- total saturation index curves (Figs. 25 and 26) plot as continuous in shape, not varying complex saturation states transitions

Shallow reservoir (accumulation/production zone) model

With analogy to deep reservoir, a shallow aquifer hosted in CHN Mid Triassic carbonates and upper zone of KNA Jr-Cr2 succession is compared to infiltration zone in CSM construction procedure. Reservoir geochemistry is represented by the samples of BEH-1 and FBe-1 wells of different depths, thus screening different zones of an aquiferous body.

Thermal waters are of secondary origin that is well documented on:

- facies difference in silica content, inverse to depth and temperature distribution in the reservoir body (Fig. 4)
- temperature overestimates on the silica-enthalpy mixing model (Fig. 6)




Fig. 31. Deep reservoir CSM. For its localization see Fig. 1.

- transition between rock dissolution and preferential alkali leaching observed at Na/K (Fig. 11) equilibrium plot (variable Na/K, K/Mg and Na/Mg ratio in a formation); and
- inconsistent plot of samples on Mg-SiO₂ alteration plot (Fig. 13) varying equilibrium temperatures between quartz and chalcedony stability at persistently low K/Mg ratio
- A reference is also given to use of mixing models in SRTW conditions (Fričovský and Tometz, 2013; Fričovský, 2014; Fričovský et al., 2015). Two reservoir facies are then recognized as:
- deep reservoir originated facies (vertically invading the shallow aquifer)
- shallow transition recharged facies (laterally recharging the shallow aquifer)

Deep reservoir originated facies

We expect the deep reservoir thermal waters to evade the bearing horizon at temperatures about 50–55 °C, initially saturated with chalcedony and oversaturated with carbonates. The vertical evasion realizes at open fault systems and quasi-vertical fissure swarms.

By the geochemical type, it is possible the evading waters preserve deep reservoir originated transient (Ca-Mg-HCO₃-SO₄ to Ca-Mg-SO₄-HCO₃) character.

Because of a loss of a contact with a source rock and change in p-T conditions, Mg-calcite is of higher tendency to precipitate spontaneously, consuming CO₂ and magnesium at various magnitude. Onset of calcite deposition is

expected to realize at late evasion stages. A drop in aqueous aggressive CO₂ stabilizes modification of Na-alkalis, slowly increasing the Na/K ratio, progressively fixing a portion of dissolved SiO₂ in increasing thermodynamic activity of Na-silicates with its minor precipitation only. This is documented on an increase in albite saturation inverse to temperature (Fig. 15).

Some loss in silica in deeper reservoir parts or prior intake is also documented on silica-enthalpy mixing model by overestimating deep reservoir temperatures considered of end-member quality. An inverse overview, provides an option of post-mixing silica release from the IWCP Borové Formation.

Shallow transition originated facies

The separate recharge towards the shallow reservoir is expected due to an open hydrogeological character of a system, aided by facies analysis commented above. At shallow transition, the hydrochemical character of intakes into aquiferous zone may vary between transient and carbonatogene type, controlled by early dilution prior submersion. Thermal waters sampled at FBe-1 well are facially affine to infiltration zone facies at the Na/K, conserving the relative ratio (Fig. 11) with simultaneous increase in alkali concentrations. Because of their position within preferential Na⁺ and K⁺ leaching regime we expect the sodium intake realizing by minor hydrolysis of sodium-silicates in the KNA Jr-Cr2 profile, altered with the slow, but progressive K⁺ dissociation from the CHN Mid Triassic dolomites and IWCP formations after the intake into production zone (Fig. 12).




Fig. 32. Shallow reservoir CSM. For its localization see Fig. 1.

Discussion on shallow reservoir mixing

Both facies mix in the shallow reservoir, in the TCM (thermal-cold-mixing) regime. Even sampled waters are clearly of thermal character ($T_{\text{res}} > 20^\circ\text{C}$), only deep reservoir facies can be recognized as thermal end-member. A hydrochemical type of associated thermal waters may vary by a magnitude of modifications between transient and carbonatogene character, controlled by a mixing rate (Fig. 32).

A mixing event off any iso-chemical and iso-thermal parameters destructs any optional equilibration attained. By mixing of both facies, the Mg^{2+} increases as residential waters are enriched by magnesium from shallow transition.

The silica content also increases as shallow intake waters dissociate silicates from the IWCP and dolomite impurities, as well as a minor contribution from a contact with KNA Jr-Cr2 bedrock formations can not be excluded.

Vyšné Sliache (secondary) discharge zone model

There is no questioning of primary discharge zone location nearby the town of Bešeňová, associated with open faults-bound vertical upwelling of geothermal waters from the deep and shallow reservoir respectively. The mixed type of springs in Bešeňová is consequent to a definite dilution of reservoir thermal waters by the groundwater from the IWCP and Quaternary cover respectively, yet conserving oversaturation with respect to carbonates, equilibrated by deposition of travertines forming dome structures.

However, the existence of dry CO_2 fluxes and thermal springs in the Vyšné Sliache area provided some early note on possible thermal waters occurrence in its nearby deep territory. A first hint on some facies relations has been provided by application of mixing models (Fričovský and Tometz, 2013; Fričovský, 2014).

Silica-enthalpy-carbonate variation

There are distinguished facies of the Vyšné Sliache territory. We assigned the VSH-1 well to represent thermal waters prior upwelling, while thermal springs represent a surface propagation of waters, with primary end-member in the deep reservoir at a first assumption.

Facies relation of VSH-1 and springs yields overestimated temperatures of well facies, contravening expected SiO_2 deficiency of losing-by-cooling (in precipitation or association of silicates), however, there is some silica lost between DRTE and VSH-1 facies, recorded by overestimate of deep reservoir temperatures (Fig. 6). The silica-enthalpy fraction model (Fig. 8) provides an amorphous silica check-test with excess resultant temperature and overestimate in the temperature by post-mixing silica dissolution. Two subfacies have been, however, identified on a silica – total carbonate model (Fig. 7) with deficient (D-LTC) and excess (D-HTC) total carbonate content, relative to DRTE and SRTW facies. Probably, the D-LTC records a “primary” subfacies evading deep reservoir with losing silica fixed in activation of sodium-silicates due to loss of CO_2 , mixing with shallow reservoir waters to give an origin of D-HTC, to form a resultant VSH-1 facies. Because of the




Fig. 33. Vyšné Sliache minor discharge zone CSM. For its localization see Fig. 1.

silica enthalpy results, the TTM regime must be realized in the Jurassic–Mid Cretaceous profile, to increase the SiO_2 content. Then, VSH-1 facies are most probably cooled and diluted rapidly by shallow groundwater (SCGW) to drop the silica and carbonate content prior upwelling as tempered springs. A dramatic cooling (by dilution and fast upflow at open faults and fissure swarms) along with rapid drop in pressure causes induced CO_2 separation of different preferential propagation towards the surface.

Discussion on equilibrium and alteration

A carbon dioxide saturation plays an essential role in evaluation of geochemical processes related to lateral runoff of deep and shallow reservoir thermal waters towards the Vyšné Sliache zone. An increase in Na/K ratio compared to DRTE is expected through the drop in available CO_2 by deposition of carbonates, mostly Mg-calcite (increase in MI inverse to Mg^{2+} ; Fig. 16) combined with the preferential leaching of Na-silicates from KNA Jr-Cr2 profile (Fig. 11). Such an increase in absolute sodium concentration owes to mixing with the shallow groundwaters (second stage) and shallow reservoir evading thermal waters at a persistent contact with a source rock, and, probably, filtration deceleration, prolonging a water-rock contact sufficient for preferential leaching. Yet the maturity index is still low (MI = 0.8–0.9).

By comparing the SRTW and VSH-1 facies, an onset of calcite deposition realizes along a leaking due to runoff velocity low enough to enable its loss. The resultant increase in K/Ca ratio (Fig. 18) is rather to drop in Ca^{2+} as the absolute K^+ level is almost stable. Meanwhile, the K/Ca ratio increases between DRTE and VSH-1 facies, as a combination of loss in Ca^{2+} (calcite deposition) and some increase in K^+ . Under relatively low K/Mg ratio, the

GKM3 (Giggenbach, 1988) geothermometer yields temperatures conform to upper deep reservoir zones ($T_{\text{K/Mg}} = 59\text{--}63^\circ\text{C}$ at GKM3). Probably a leaching of potassium is fixed by adularia (Fig. 15) and illite (Fig. 24) oversaturation, controlled by CO_2 reduction of precipitation of Mg-calcite. An absolute concentration of alkalis in VSH-1 may then be a consequence of mixing between SRTW and DRTE facies in the KNA Jurassic–Mid Cretaceous horizon.

We also inspected facial and maturity relations on the Na-K-Mg geoindicator (Fig. 17). Thermal waters of VSH-1 are different in Mg^{2+} compared to DRTE, however, relatively conform to SRTW facies. On the Na-K-Mg-Ca geoindicator (Fig. 21), the VSH-1 shows, however, no relation to reservoir facies. In coincidence to silica – enthalpy conclusions, an expected loss in Mg^{2+} by deposition of carbonates at low temperatures is partially replaced by mixing of VSH-1 waters with S-CGW hosted in IWCP prior they upwell as thermal springs in the Vyšné Sliache area (Fig. 33). The polystageous mixing explains overrated dilution ratio at silica-enthalpy fraction model (Fig. 8).

A contribution to lateral leaking mechanisms

There has not been, however, any debate on probable driving mechanisms of lateral leakings towards the Sliache area. Hydraulic models presented in Fendek et al. (1988); Remšík et al. (1993); Remšík et al. (1998) provided some idea on preferential piezometry gradients from the central elevated parts towards SW. This may then be an option for general runoff, not providing conditions for some residency of thermal waters in the structure (within the accumulation/transition zone).

In fact, the mechanism has not been subjected to study neither in this paper. However, there are two more options

to be added into a complex CSM. The lateral leaking of thermal waters may then be induced by combination of:

- piezometric gradient
- build-up of density gradient (deep reservoir)
- difference in reservoir pressure

CONCLUSIONS

Tens of decades have passed since the geothermometers were successfully applied in the reservoir temperature estimation (e.g. Fournier, 1979; Arnórsson et al., 1983; Giggenbach, 1988) or the conceptual site models construction and analysis of geochemical regime and origination of thermal waters (e.g. Truesdell et al., 1977; Kharaka et al., 1982; Antrodischia et al., 1985). Only a few notes on isotope (Remšík et al., 1998) and sulphate (Bodiš and Borosová, 1996) geothermometry application were discussed with reference to the Bešeňová elevation geothermal structure. In fact, many techniques failed in temperature estimation (quantitative application) in local conditions (e.g. Fendek et al., 1988; Remšík et al., 1993; Remšík et al., 1998), because of the immaturity, low temperature conditions or predominant carbonate-hosted reservoirs. Being based on thermodynamics and hypothetical fluid-mineral equilibrium, geothermometers, however, represent a powerful tool for conceptual site models construction once correlated with the water chemistry (boiling and mixing) models, geology, alteration mineralogy, hydrogeology, and geothermics.

The Bešeňová elevation may be concluded as a complex hydrogeothermal structure of open hydrogeological character. Besides expected lateral inflow from the Liptovská Mara depression (e.g. Fig. 30), realized in the quasi-isochemical and isothermal regime, the correlation of ionic ratios with mixing models (presented e.g. in Fričovský and Tometz, 2013) has identified a minor recharge zone towards the Kľačianka valley that correlates with the local hydraulics and piezometry. Infiltration originates by at least one mixing episode between shallow and deep filtration channels (Fig. 29) prior separation from upwelling as tempered (warm) springs or descent into transition-accumulation zone. Deep reservoir waters are recharged in quasi-isochemical regime. Thermal waters are saturated with the calcite prior intake (Fig. 18), whilst attaining saturation with dolomite just after an inlet. Increase in CO₂ content under oversaturation with carbonates (Figs. 19 and 20) modifies Na-silicates towards K-silicates, simultaneous to preferential K⁺ leaching from reservoir impurities (Fig. 31). Meanwhile, it is possible the LT alteration of impure dolomites realizes on a thermodynamic activity scale, altering a talc-chlorite-muscovite (illite) subsystem. Immaturity of thermal waters because of low Na/K ratio

and possible modification of preferential thermodynamic activities of alkali silicates is driven and proportional by vertical separate filtration in convection cells due to unconform base overheating. Possibility of silicates alteration is documented by saturation of waters with chalcedony (Figs. 4 and 5). In vertical evasion regime, thermal waters rise up along open faults towards the shallow reservoir, where they mix with recharges from a shallow filtration, prior upwelling as springs, depositing travertine according to secondary saturation with calcite or travertine (Fig. 32). In the Sliače springs area (Fig. 33), it is possible that shallow and deep reservoir leakages mix in a first stage under TTM regime, prior dilution by shallow groundwaters (TCM regime). Rapid ascent along faults from the deep reservoir causes drop in pCO₂ driving formation of dry carbon dioxide fluxes on a surface, supported by second degassing stage after dilution. The mixture propagates as warm springs. Besides, a part of thermal waters leaks laterally towards the Ivachnová depression, due to regional piezometry (Fig. 30). Yet available geothermometers have been applied quantitatively as well. A silica-chalcedony conductive models appear of a best match referring to reservoir temperature conditions instead of comparing results to the wellhead measurements (Fig. 9, Appendix A). More precise results have been, however, derived of saturation index temperature (T_{CEQ} for chalcedony; Tabs. 1 and 2). For the deep reservoir environment, cation geothermometers are definitely skewed by immaturity (Figs. 16 and 17), with an exception of Mg-corrected Na-K-Ca geothermometer (Fig. 22). Multicomponent equilibrium geothermometry pointed on some complex equilibrium attainment in deeper (descending) parts of the deep reservoir (Figs. 25 and 26). Other zones were analysed in details in the main text.

Because of complexity and immaturity of a deep reservoir (assumed as a most potential of the entire basin), we rather recommend continuing studies focused at definition of local ge indicators (such as chalcedony, gypsum) for onward exploitation monitoring and relation analysis of thermal waters to other structures associated with the basin.

If not for quantitative temperature estimation and correlation, geothermometry techniques are distinctively valid for qualitative analysis either in low enthalpy or carbonates-hosted systems, if correlated together and applied with caution and reference to their essential principles.

Acknowledgement. Authors would sincerely like to thank the Andrea Vranovská, PhD. (Water Research Institute, Slovakia), prof. Peter Szucs (University of Miskolc, Hungary) and Peter Bajtoš, PhD. (State geological institute of Dionýz Štúr, Slovakia) for their contribution to this paper.

References

- ANTRODICCHIA, E., CIONI, R., CHIODINI, G., GAGLIARDI, R. & MARINI, L., 1985: Geochemical temperatures of the thermal waters of Phlegraean Fields (Naples, Italy). *Geothermal Resources Council Transactions*, 9, 287–292.
- ARNÓRSSON, S., 1985: The use of mixing models and chemical geothermometers for estimating underground temperatures in geothermal systems. *J. Volcanol. Geotherm. Res.*, 23, 299–335.
- ARNÓRSSON, S., 2000a: Mineral saturation. In: Arnórsson, S. (ed.): *Isotopic and chemical techniques in geothermal exploration, development and use*, IAEA, Vienna, 152–200.
- ARNÓRSSON, S., 2000b: Assessment of reservoir fluid composition from wet steam well data. In: Arnórsson, S. (ed.): *Isotopic and chemical techniques in geothermal exploration, development and use*, IAEA, Vienna, 212–228.
- ARNÓRSSON, S. & GUNNLAUGSSON, E., 1985: New gas geothermometers for geothermal exploration – calibration and application. *Geochim. Cosmochim. Acta*, 49, 1 307–1 325.
- ARNÓRSSON, S., SIGURDSSON, S. & SVAVARSSON, H., 1982: The chemistry of geothermal waters in Iceland I. Calculation of aqueous speciation from 0° to 370 °C. *Geochim. Cosmochim. Acta*, 46, 1 513–1 532.
- ARNÓRSSON, S., GUNNLAUGSSON, E. & SVAVARSSON, H., 1983: The chemistry of geothermal waters in Iceland III. Chemical geothermometry in geothermal investigations. *Geochim. Cosmochim. Acta*, 47, 567–577.
- ARNÓRSSON, S., ANDRÉSDÓTTIR, A., GUNNARSSON, I. & STÉFANSSON, A., 1988: New calibration for the quartz and Na/K geothermometers – valid in the range 0–350 °C. In: *Proceedings Geoscience Society of Iceland Annual Meeting, Reykjavik, Iceland*, 42–43.
- ARNÓRSSON, S., FRIDRIKSSON, Th. & GUNNARSSON, I., 1998: Gas chemistry of the Krafla geothermal field, Iceland. In: *Proceedings International Symposium on Water-Rock Interaction, Auckland, New Zealand*, 613–66.
- AXELSSON, G., 2009: Conceptual models of geothermal systems – introduction. In: *Proceedings Short Course V on Conceptual Modelling of Geothermal Systems, Santa Tecla, El Salvador*, 1–12.
- BODIŠ, D. & BOROSOVÁ, S., 1996: Hydrogeochemical evaluation of the mineral and thermal waters of the Liptovská kotlina Basin: Application of the factor analysis. *Geol. Práce, Spr.*, 102, 65–70. (In Slovak with English summary.)
- BROWNE, 1978: Hydrothermal alteration in active geothermal fields. *Ann. Rev. Earth Planet. Sci.*, 6, 229–250.
- CANNON, C., WOOD, T., NEUPANE, G., MCLING, T., MATTSON, E., DOBSON, P. & CONRAD, M., 2014: Geochemistry sampling for traditional and multicomponent equilibrium geothermometry in Southeast Idaho. *Geothermal Resources Council Transactions*, 38, 425–431.
- CHIODINI, G., CIONI, R., FRULLANI, A., GUIDI, M., MARINI, L., PRATI, F. & RACO, B., 1996: Fluid geochemistry of Montserrat Island, West Indies. *Bull. Volcanology*, 58, 380–392.
- CIONI, R., FANELLI, G., GUIDI, M., KINYARIRO, J. K. & MARINI, L., 1992: Lake Bogoria hot springs (Kenya): geochemical features and geothermal implications. *J. Volcanol. Geotherm. Res.*, 50, 231–246.
- COOPER, D. C., PALMER, C. D., SMITH, R. W. & MCLING, T. L., 2013: Multicomponent equilibrium models for testing geothermometry approaches. In: *Proceedings 38th Workshop On Geothermal Reservoir Engineering, Stanford University, California, paper No. SGP-TR-198*, 1–10.
- CUMMING, W., 2009: Geothermal resource conceptual models using surface exploration data. In: *Proceedings the 34th Workshop on Geothermal Reservoir Engineering, Stanford University, CA, USA*, 6 pp.
- ČINCURA, J., & KÖHLER, E., 1995: Paleoalpine karstification: the longest paleokarst period in the Western Carpathians (Slovakia). *Geol. Carpath.*, 46, 5, 343–347.
- DAVRAZ, A., 2014: Application of hydrogeochemical techniques in geothermal systems; examples from eastern Mediterranean region. In: Baba, A., Bundschuh, J. & Chandrasekharan, D. (eds.): *Geothermal systems and energy resources: Turkey and Greece*. CRC press, London, UK, 336 p.
- D'AMORE, F. & ARNÓRSSON, S., 2000: Geothermometry. In: Arnórsson, S. (ed.): *Isotopic and chemical techniques in geothermal exploration, development and use*, IAEA, Vienna, 152–200.
- ELLIS, A. J., 1970: Quantitative interpretation of chemical characteristics of hydrothermal systems. *Geothermics*, 2, 516–528.
- ELLIS, A. J. & MAHON, W. A. J., 1977: Chemistry and geothermal systems. *Academic Press, New York*, 392 pp.
- FENDEK, M., BODIŠ, D., BIELY, A., KULLMANOVÁ, A., GAŠPÁRIKOVÁ, V., SNOPOKOVÁ, P., KRÁL, M. & JANČI, J., 1988: The report on the investigation geothermal borehole ZGL-1 in the Bešeňová locality – the verification of the prognostic sources of geothermal energy in the Liptovská kotlina Basin – West. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava*, 85 pp. (In Slovak.)
- FENDEK, M., FENDEKOVÁ, M., DANIEL, S. & KULTAN, V., 2013: Bešeňová – usable quantity of geothermal energy, supplementary hydrogeological survey. *HYDROFEN Bratislava*, 91 pp. (In Slovak.)
- FOUILLAC, R. & MICHARD, S., 1981: Sodium/lithium ratio in water applied to geothermometry of geothermal reservoirs. *Geothermics*, 10, 55–70.
- FOURNIER, R. O., 1973: Silica in thermal waters: laboratory and field investigations. *Proceedings International Symposium on Hydrogeochemistry and Biogeochemistry, Tokyo, 1970, vol. 1, Hydrochemistry, Washington, DC, 1973*, 122–139.
- FOURNIER, R. O., 1977: Chemical geothermometers and mixing models for geothermal systems. *Geothermics*, 5, 41–50.
- FOURNIER, R. O., 1979: A revised equation for the Na-K geothermometer. *Geothermal Resource Council Transactions*, 3, 221–224.
- FOURNIER, R. O., 1980a: Application of water geochemistry to geothermal and reservoir engineering. *Spanish Institute of Geology and Mineralogy*, 5–56.
- FOURNIER, R. O., 1980b: The interpretation of Na-K-Mg relations in geothermal waters. *Geothermal Resources Council Transactions*, 14, 2, 1 421–1 425.
- FOURNIER, R. O., 1981: Applications of water chemistry for geothermal exploration and geothermal engineering. In: Rybach, L. & Muffler, L. P. J. (eds.): *Geothermal Systems – Principles and Case Histories*, Wiley PH, New York, USA, 109–143.

- FOURNIER, R. O., 1982: Geochemistry in geothermal exploration and production. In: *Proceedings 5th New Zealand Geothermal Workshop, Auckland, New Zealand*, 209–215.
- FOURNIER, R. O., 1989: Lectures on geochemical interpretation of hydrothermal waters. United Nations Geothermal Training Programme, Reykjavik, Iceland, 77 pp.
- FOURNIER, R. O., 1991: Water geothermometers applied to geothermal energy. In: D'Amore, R. (ed.): *Application of geochemistry in Geothermal Reservoir Development*, UNITAR, Rome, 37–69.
- FOURNIER, R. O. & ROWE, J. J., 1966: Estimation of underground temperatures from the silica content of water from hot springs and wet-steam wells. *Amer. J. Sci.*, 264, 685–697.
- FOURNIER, R. O. & TRUESDELL, A. H., 1971: An empirical geothermometer based on Na, K and Ca dissolved in natural waters (abstract). *Geol. Soc. Amer. Abstr.*, 3, 570.
- FOURNIER, R. O. & TRUESDELL, A. H., 1973: An empirical Na-K-Ca geothermometer for natural waters. *Geochim. Cosmochim. Acta*, 37, 1 255–1 275.
- FOURNIER, R. O., WHITE, D. E. & TRUESDELL, A. H., 1974: Geochemical indicators of subsurface temperature – Part 1, basic assumptions. *J. Res. U. S. Geol. Survey*, 2, 259–261.
- FOURNIER, R. O. & POTTER, R. W. II., 1978: A magnesium correction for the Na-K-Ca chemical geothermometer. *U.S. Geological Survey Open-File Report*, 78–986, 24.
- FOURNIER, R. O. & POTTER, R. W. II., 1979: Magnesium correction to the Na-K-Ca chemical geothermometer. *Geochim. Cosmochim. Acta*, 43, 1 543–1 550.
- FOURNIER, R. O. & POTTER, R. W. II., 1982: A revised and expanded silica (quartz) geothermometer. *Geothermal Resources Council Bulletin*, 11, 3–12.
- FRANKO, O., 1984: Bešeňová – carbonated thermal waters. In: *Hydrogeologický sprievodca VIII celoštátnej hydrogeologickej konferencie. GÚDŠ, Bratislava. (In Slovak.)*
- FRANKO, O., 2002a: Recent views on hydrogeological structure of mineral waters in Lúčky locality. *Podzemná voda*, VIII, 2, 123–132. (In Slovak.)
- FRANKO, O., 2002b: New data for suggestion of protection zones of natural healing resources in Lúčky locality. *Manuscript – Final report, Kúpele Lúčky*, 39 pp. (In Slovak.)
- FRANKO, O., BIELY, A., HRICKO, J., SZALAIOVÁ, V. & MATEOVIČ, L., 1974: Evaluation of the Demänová area in the Liptovská kotlina Basin for searching for the thermal waters. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava*, 50 pp. (In Slovak.)
- FRANKO, O., BODIŠ, D., GAZDA, S. & MICHALÍČEK, M., 1979: Hydrogeological evaluation of the Liptovská kotlina Basin from the view of the mineral waters occurrence. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava*, 79 pp. (In Slovak.)
- FRIČOVSKÝ, B., 2014: Composite conceptual model and hydrogeothermal evaluation of the Bešeňová elevation hydrogeothermal structure, Liptov Basin, Slovakia. *Manuscript – Doctoral (Ph.D.) Thesis, Institute of Geosciences, Technical University of Košice*, 301 pp.
- FRIČOVSKÝ, B. & TOMETZ, L., 2013: Reservoir boiling and mixing models overview and application: case study on the Bešeňová elevation hydrogeothermal structure (northern Slovakia). In: *Proceedings on Hydrogeochemistry 2013 International Conference, XIV, Sosnowiec, Poland*, 1–12.
- FRIČOVSKÝ, B. & TOMETZ, L., 2014: Use of the multicomponent geothermometry at formation of the conceptual model of hydrogeothermal structures, Bešeňová Elevation, Slovakia. In: *Geochémia 2014 – zborník vedeckých príspevkov z konferencie. Št. Geol. Úst. D. Štúra, Bratislava*, 42–46. (In Slovak.)
- FRIČOVSKÝ, B., TOMETZ, L., VIZI, L. & ŠTIAKOVÁ, J., 2014: Update on stationary temperature model on carbonates dominated, stratified, low enthalpy hydrogeothermal system of the Bešeňová Elevation, Slovakia. In: *Proceedings 14th SGEM Conference, Vol. II – Hydrogeology, engineering geology and geotechnics, Albena, Bulgaria*, 1 003–1 011.
- FRIČOVSKÝ, B., TOMETZ, L., FENDEK, M. & GUMÁŇOVÁ, J., 2015: Update on composite geochemical conceptual model for the Bešeňová elevation hydrogeothermal structure, Liptov Basin, northern Slovakia. In: *Proceedings World Geothermal Congress 2015, Melbourne, Australia, paper No. 14 092*, 1–11.
- GIBBS, R. J., 1970: Mechanisms controlling world water chemistry. *Science*, 170, 1 088–1 090.
- GIGGENBACH, W. F., 1981: Geothermal mineral equilibria. *Geochim. Cosmochim. Acta*, 45, 393–410.
- GIGGENBACH, W. F., 1984: Mass transfer in hydrothermal alteration systems. *Geochim. Cosmochim. Acta*, 48, 2 963–2 711.
- GIGGENBACH, W. F., 1986: Graphical techniques for the evaluation of water/rock equilibration conditions by use of Na, K, Mg and Ca contents of discharge waters. In: *Proceedings 8th New Zealand Geothermal Workshop*, 37–43.
- GIGGENBACH, W. F., 1988: Geothermal solute equilibria. Derivation of Na-K-Mg-Ca geoindicators. *Geochim. Cosmochim. Acta*, 52, 2 749–2 756.
- GIGGENBACH, W. F., 1991: Chemical Techniques in Geothermal Exploration. In: D'Amore, F. (ed.): *Application of geochemistry in geothermal reservoir development. UNITAR/UNDP publication, Rome*, 119–142.
- GIGGENBACH, W. F., 1995: Geochemical exploration of a difficult geothermal system, Paraso, Vella Lavella, Solomon Islands. In: *Proceedings World Geothermal Congress 1995, Firenze, Italy*, 995–1 000.
- GIGGENBACH, W. F. & GOGUEL, R. L., 1989: Collection and analysis of geothermal and volcanic water and gas discharges. DSIR report CD 2401, 4th ed., Pentone, New Zealand.
- GIGGENBACH, W. F., GONFIANTINI, R., JANGI, B. L. & TRUESDELL, A. H., 1983: Isotopic and chemical composition of Parbati Valley geothermal discharges, NW Himalaya, India. *Geothermics*, 12, 199–222.
- GIGGENBACH, W. F., SHEPPARD, D. S., ROBINSON, B. W., STEWART, M. K. & LYON, G. L., 1994: Geochemical structure and position of the Waiotapu geothermal field, New Zealand. *Geothermics*, 23, 599–644.
- GIORDANO, T. H. & SWANBERG, C. A., 1978: Influence of reservoir rock-type on the Na-K-Ca geothermometer. *Geological Society of America Abstracts with Programs*, 2, 7, 433.
- GISLASON, S. R., HEANEY, P. J., OELKERS, E. H. & SCHOTT, J., 1997: Kinetic and thermodynamic properties of moganite, a novel silica polymorph. *Geochim. Cosmochim. Acta*, 61, 1 193–1 204.
- GROSS, P., BIELY, A., KÖHLER, E. & PAPŠOVÁ, J., 1979: Geology of the Inner Carpathian Paleogene in the western part of the Liptovská kotlina Basin. *Manuscript – Technical report Sta-*

- te geological institute of Dionýz Štúr, Bratislava, 32 pp. (In Slovak.)
- GROSS, P., KÖHLER, E., BIELY, A., FRANKO, O., HANZEL, V., HRICKO, J., KUPČO, G., PAPŠOVÁ, J., PRIECHOVSKÁ, Z., SZALAIOVÁ, V., SNOPOKOVÁ, P., STRÁNSKÁ, M., VAŠKOVSKÝ, I. & ZBOŘIL, L., 1980: Geology of the Liptovská kotlina Basin. *Št. Geol. Úst. D. Štúra, Bratislava*, 236 pp. (In Slovak with English summary.)
- GUNNARSSON, I. & ARNÓRSSON, S., 2000: Amorphous silica solubility and thermodynamic properties of H_4SiO_4 in the range 0–350 °C at Psat. *Geochim. Cosmochim. Acta*, 64, 2 295–307.
- HANZEL, V., KULLMAN, E. & VRANA, K., 2008: Lúčky – computation of quantities of mineral waters. Final report. *Manuscript. Archives of the State Geological Institute of Dionýz Štúr, Bratislava*, 220 pp. (In Slovak.)
- HORITA, J., 2005: Stable isotope thermometry: There is more to it than temperature. *Geochem. J.*, 39, 481–496.
- JOHNSON, J. W., OELKERS, E. H. & HELGESON, H. C., 1992: SUPCTR92: A software package for calculating the standard modal thermodynamic properties of minerals, gases, aqueous species and reactions from 1 to 5 000 bar and 0 to 1 000 °C. *Computers & Geosciences*, 7, 899–947.
- JOSEPH, E. P., FOURNIER, N., LINDSAY, J. M., ROBERTSON, R., DENISE, M. & BECKLES, D. M., 2013: Chemical and isotopic characteristics of geothermal fluids from Sulphur Springs, Saint Lucia. *J. Volcanol. Geotherm. Res.*, 254, 23–36.
- JUREWYCZ, E., 2005: Geodynamic evolution of the Tatra Mts. and the Pieniny Klippen Belt (Western Carpathians): problems and comments. *Acta geol. pol.*, 55, 3, 295–338.
- KARINGITHI, C. W., 2009: Chemical geothermometers for geothermal exploration. In: *Proceedings Short Course on Exploration of Geothermal Resources, Lake Naivasha, Kenya, paper No. 22*.
- KHARAKA, Y. K. & MARINER, R. H., 1989: Chemical geothermometers and their application to formation waters from sedimentary basins. In: *Naeser, N. D. & McCulloh, T. H. (eds.): Thermal History of Sedimentary Basins. Springer, New York*, 99–111.
- KHARAKA, Y. K., LICO, M. S. & LAX, L. M., 1982: Chemical geothermometers applied to formation waters, Gulf of Mexico and California basins. *Amer. Assoc. Petrol. Geol. Bull.*, 66, 588.
- KLAGO, M. & STUCHLIKOVÁ, B., 1990: Liptovský Ján, Pod Bieľym, hydrogeologický prieskum. Final report. *Št. Geol. Úst. D. Štúra, Bratislava*, 40 pp.
- LYON, G. L., 1974: Geothermal gasses. In: *Kaplan, I. R. (ed.): Natural Gases in Marine Sediments, Plenum Press, New York*.
- MAĐAR, D., 1997: Liptovská kotlina Basin – regional hydrogeothermal evaluation, geophysical survey. *Manuscript – Final report, ESPRIT, s. r. o., Banská Štiavnica*, 27 pp. (In Slovak.)
- MARINI, L., 2004: Geochemical techniques for the exploration and exploitation of geothermal energy. *Viareggio, Italy*, 106 p.
- MOREY, G. W., FOURNIER, R. O. & ROWE, J. J., 1962: The solubility of quartz in water in temperature interval from 25 to 300 °C. *Geochim. Cosmochim. Acta*, 26, 1 029–1 043.
- NĚMEC, F. & BARTKOVÁ, J., 1987: Paleogene of the Liptovská kotlina Basin and its underlier. *Manuscript – Final report, Naftoprojekt Bratislava, š. p.*, 181 pp.
- NEUPANE, G., MATTSON, E., MCLING, T., PALMER, D., SMITH, R. W. & WOOD, T. R., 2014: Deep geothermal reservoir temperatures in the Eastern Snake River Plain, Idaho, using multi-component geothermometry. In: *Proceedings 39th Workshop on Geothermal Reservoir Engineering, Stanford University, California, SGP-TR-202*, 1–12.
- NICHOLSON, K., 2012: Geothermal fluids: chemistry and exploration techniques. *Springer Science & Business Media*, 263 pp.
- NIEVA, D. & NIEVA, R., 1987: Developments in geothermal energy in Mexico, part 12: A cationic composition geothermometer for prospection of geothermal resources. *Heat Recovery Systems and CHP*, 7, 243–258.
- PAČES, T., 1975: A systematic deviation from Na-K-Ca geothermometer below 75 °C and above 10^{-4} atm P_{CO_2} . *Geochim. Cosmochim. Acta*, 39, 541–544.
- PALMER, C. D., OHLY, S. R., SMITH, R. W., NEUPANE, G., MCLING, T. & MATTSON, E., 2014: Mineral selection for multicomponent equilibrium geothermometry. *Geothermal Resources Council Transactions*, 38, 453–459.
- PANG, Z. H., 1988: Multiple fluid-mineral equilibrium calculations and their applications to geothermometry and hydrochemical processes in geothermal systems. *Manuscript – United Nations University Geothermal Training Programme, Reykjavik, Iceland, report No. 5*, 82 pp.
- PANG, Z. H., 2001: Isotope and chemical geothermometry and its application. *Sciences in China – series E, Technological Sciences*, 44, 1, 16–20.
- PANG, Z. H. & REED, M. H., 1998: Theoretical chemical thermometry on geothermal waters: problems and methods. *Geochim. Cosmochim. Acta*, 62, 1 083–1 091.
- PALANDRI, J. & REED, M. H., 2001: Reconstruction of in-situ composition of sedimentary waters. *Geochim. Cosmochim. Acta*, 65, 1 741–1 767.
- PEIFFER, L., WANNER, C., SPYCHER, N., SONNENTHAL, E. L. & KENNEDY, B. M., 2013: Multicomponent vs. classical geothermometry: An evaluation using reactive transport modeling. *Proc. Earth Planet. Sci.*, 7, 665–668.
- PEIFFER, L., WANNER, C., SPYCHER, N., SONNENTHAL, E. L., KENNEDY, B. M. & IOVENITTI, J., 2014: Optimized multicomponent vs. classical geothermometry: Insights from modeling studies at the Dixie Valley geothermal area. *Geothermics*, 15, 154–169.
- REED, M. H., 1982: Calculation of multicomponent chemical equilibria and reaction processes in systems involving minerals, gases and aqueous phase. *Geochim. Cosmochim. Acta*, 46, 513–528.
- REED, M. H. & SPYCHER, N., 1983: Calculated effects of boiling, cooling and oxidation on precipitation of ore and gangue minerals from hydrothermal solution. In: *Proceedings 4th International Symposium on Water Rock Interaction – Extended Abstracts, Misasa, Japan*, 405–408.
- REED, M. H. & SPYCHER, N., 1984: Calculation of pH and mineral equilibria in hydrothermal waters with application to geothermometry and studies of boiling and dilution. *Geochim. Cosmochim. Acta*, 48, 1 479–1 492.

- REMŠÍK, A., FENDEK, M., KRÁL, M., BODIŠ, D. & MICHALKO, J., 1993: Geothermal energy of the Liptovská kotlina Basin. *Manuscript – Technical report, State geological institute of Dionýz Štúr, Bratislava, 97 pp. (In Slovak.)*
- REMŠÍK, A., FENDEK, M., MELLO, J., KRÁL, M., BODIŠ, D., MICHALKO, J., MAĐAR, D. & VÍKA, K., 1998: Liptovská kotlina Basin – regional hydrogeothermal evaluation. *Manuscript – Final report, Ministry of the Environment of the Slovak Republic, Bratislava, 94 pp. (In Slovak.)*
- REMŠÍK, A., FENDEK, M. & MAĐAR, D., 2005: The occurrence and distribution of the geothermal waters in the Liptovská kotlina Basin. *Miner. Slov., 37, 123–130.*
- SANJUAN, B., 2010: Use of a new Sodium/Lithium (Na/Li) geothermometric relationship for High-Temperature (HT) geothermal fluids derived from seawater/basalt interaction processes: Application to the Djibouti case. *In: Proceedings 3rd Third East African Rift Geothermal Conference AR-GEO-C3-DJIBOUTI, Djibouti, 231–248.*
- SANJUAN, B., GIRARD, J. P., LANINI, S., BOURGUIGNON, A. & BROSE, E., 2003: Geochemical modelling of diagenetic illite and quartz cement formation in Brent sandstone reservoirs: example of the Hild Field, Norwegian North Sea. *In: Worden, R. H. & Morad, S. (eds.): Clay mineral cements in Sandstones. Intern. Assoc. Sedimentol. Spec. Publ., Blackwell Publishing, Oxford, 34, 425–452.*
- SANTOYO, E. & DÍAZ-GONZÁLEZ, L., 2010: A new improved proposal of the Na/K geothermometer to estimate deep equilibrium temperatures and their uncertainties in geothermal systems. *In: Proceedings World Geothermal Congress 2010, Bali, Indonesia, 1–9.*
- SHERIDAN, M. F., SATKIN, R. L. & WOHLITZ, K. W., 1980: Geothermal resource evaluation at Castle Hot Spring. *Manuscript – Final report, Arizona Geological Survey, Tuscon, AZ, 57 pp.*
- SHIKANOZO, N., 1976: Thermodynamic interpretation of Na-K-Ca geothermometer in the natural water system. *Geochem. J., 10, 47–50.*
- SIMMONS, S. F. & CHRISTENSON, B. W., 1993: Towards a unified theory on calcite formation in boiling geothermal systems. *In: Proceedings 15th New Zealand Geothermal Workshop, Auckland, New Zealand, 145–148.*
- SIMMONS, S. F., 2013: Fluid-mineral equilibria in great basin geothermal systems: implications for chemical geothermometry. *In: Proceedings, 38th Workshop on Geothermal Reservoir Engineering, Stanford University, Stanford, California, 1–8.*
- SMITH, R. W., PALMER, C. D. & COOPER, D. C., 2012: Approaches for Multicomponent Equilibrium Geothermometry as a Tool for Geothermal Resource Exploration. *In: Proceedings American Geophysical Union Fall Meeting 2012, 12, abstract No. V23F-08, 1 p.*
- SONNEY, R. & VUATAZ, F., 2010: Validation of chemical and isotopic geothermometers from low temperature deep fluids of Northern Switzerland. *In: Proceedings World Geothermal Congress 2010, Bali, Indonesia, paper No. 1 426.*
- SPYCHER, N., SONNENTHAL, E. L. & KENNEDY, B. M., 2011: Integrating multicomponent chemical geothermometry with parameter estimation computations for geothermal exploration. *Geothermal Resource Council Transactions, 35, 663–666.*
- SPYCHER, N., PEIFFER, L., SONNENTHAL, E. L., SALDI, G., REED, M. H. & KENNEDY, B. M., 2014: Integrated multicomponent solute geothermometry. *Geothermics, 51, 113–123.*
- TAYLOR, S. R., 1964: Abundance of chemical elements in the continental crust, a new table. *Geochim. Cosmochim. Acta, 28, 1 273–1 285.*
- TOLE, M. P., 1988: Geochemical sampling, laboratory analysis, and mineral equilibria calculations for fluids from selected areas in Iceland. United Nations University Geothermal Training Programme, Reykjavik, Iceland, report No. 7, 55.
- TOLE, M. P., ÁRMANNSSON, H., PANG, Z. H. & ARNÓRSSON, S., 1993: Fluid/mineral equilibrium calculations for geothermal fluids and chemical geothermometry. *Geothermics, 22, 17–37.*
- TONANI, F., 1980: Some remarks on the application of geochemical techniques in geothermal exploration. *In: Proceedings Advances in European Geothermal Research, 2nd Symposium, Strassbourg, France, 428–443.*
- TRUESDELL, A. H., 1975: Geochemical techniques in exploration. *In: Proceedings 2nd UN Symposium on Development and Use of Geothermal Resources 1, 53–86.*
- TRUESDELL, A. H., 1976: Summary of section III: geochemical techniques in exploration. *In: Proceedings 2nd UN Symposium on Development and Use of Geothermal Resources, San Francisco, USA, 1, 53–89.*
- TRUESDELL, A. H. & FOURNIER, R. O., 1975: Calculation of deep temperatures in geothermal systems from the chemistry of boiling spring waters of mixed origin. *In: Proceedings 2nd U.N. Symposium on Geothermal Resources, San Francisco, CA, 837–844.*
- TRUESDELL, A. H. & FOURNIER, R. O., 1977: Procedure for estimating the temperature of a hot water component in a mixed water using a plot of dissolved silica versus enthalpy. *J. Res. U.S. Geological Survey, 5, 1, 49–52.*
- TRUESDELL, A. H., NATHENSON, M. & TYE, R. O., 1977: The effects of sub-surface boiling and dilution on the isotopic compositions of Yellowstone thermal waters. *J. Geophys. Res., 82, 3 694–3 703.*
- VANDROVÁ, G. & MATEJČEKOVÁ, E., 1991: Liptovský Ján, Jánska dolina valley – thermal well “Rudolf”, hydrodynamic tests. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava. (In Slovak.)*
- VANDROVÁ, G., ŠTEFANKA, P. & ČERVEŇAN, M., 2009: Bešeňová – revision of exploitation conditions of the sources ZGL-1 and FB-1. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava, 101 pp. (In Slovak.)*
- VANDROVÁ, G., ŠTEFANKA, P., HÓK, J. & SÝKORA, M., 2011: Bešeňová – hydrogeothermal borehole FGTB-1, 1st and 2nd phase. *Manuscript – Technical report, State geological institute of Dionýz Štúr, Bratislava, 140 pp. (In Slovak.)*
- VERMA, S. P., 2000: Limitations in applying silica geothermometers for geothermal reservoir evaluation. *In: Proceedings 25th Workshop on Geothermal Reservoir Engineering, Stanford University, CA, 1–6.*
- VERMA, S. P., 2012: Cation exchange geothermometry: A critique. *Geothermics, 25, 1, 53–56.*
- VERMA, S. P. & SANTOYO, E., 1997: New improved equations for Na/K, Na/Li and SiO₂ geothermometers by outlier detection and rejection. *J. Volcanol. Geotherm. Res., 79, 1–2, 9–24.*

- WALTHER, J. V. & HELGESON, H. C., 1977: Calculation of thermodynamic properties of aqueous silica and the solubility of quartz and its polymorphs at high pressures and temperatures. *Amer. J. Sci.*, 277, 1 315–1 351.
- WISHART, DeBONNE, N., 2015: Comparison of silica and cation geothermometers of Bath Hot Springs, Jamaica WI. In: *Proceedings World Geothermal Congress 2015, Melbourne, Australia, Paper No. 14 130*.
- ZBOŘIL, Ľ., MÁJOVSKÝ, J., TAKÁČOVÁ, H. & HUSÁK, Ľ., 1972: The Liptovská kotlina Basin – geoelectric, gas flux and thermometric measurements within the period 1971–1972. *Manuscript – Technical report, State geological institute of Dionýz Štúr, Bratislava, 40 pp. (In Slovak.)*
- ZEMBIAK, O., BIELY, A., BODIŠ, D., FENDEK, M., FRANKO, O., GABAUER, G. & BÁLINT, T., 1986: Geothermal energy – investigation of the re-injection possibilities and evaluation of the SSR potential – investigation of the geothermal sources of the Liptovská kotlina Basin. *Manuscript – Final report, State geological institute of Dionýz Štúr, Bratislava, 51 pp. (In Slovak.)*

Využitie geotermometrie pri odhade rezervoárovej teploty a konštrukcii koncepčných modelov: Princípy, metódy a aplikácie v podmienkach hlbokého rezervoáru hydrogeotermálnej štruktúry bešeňovskej elevácie, Slovensko.

Aplikácia geotermometrov pre odhad teploty (napr. Fournier, 1979; Arnórsson et al., 1983; Giggenbach, 1988), objasnenie geochemického pôvodu, podmienok formovania termálnych vôd a tvorbu koncepčných modelov (hydro)geotermálnych štruktúr a polí (napr. Truesdell et al., 1977; Kharaka et al., 1982; Antrodischia et al., 1985) je celosvetovo uznávanou metódou na podporu realizovaných geologických, hydrogeologických a geofyzikálnych štúdií.

Využívanie metód geotermometrického výskumu a prieskumu bolo v podmienkach Západných Karpát dlhodobo zaznávané a obmedzené na niekoľko zmienok v lokálnych podmienkach štruktúry Bešeňovskej elevácie (Bodiš a Borosová, 1996; Remšík et al., 1998). Je pravdou, že mnohé z nich pravidelne zlyhávali pri odhade rezervoárovej teploty (napr. Fendek et al., 1988; Remšík et al., 1993; Remšík et al., 1998) termálnych vôd v dôsledku ich termodynamickej nevyzrelosti, nízkych teplôt a prevažne karbonátového prostredia. Avšak, nakoľko sú tieto metódy založené na striktných termodynamických zákonitostiach a predpoklade zachovania rovnovážneho stavu na rozhraní, napríklad, termálna voda – minerálna fáza, geotermometre predstavujú v skutočnosti dôležitý prostriedok pri konštrukcii koncepčných modelov hydrogeotermálnych a geotermálnych štruktúr, ak sú korelované s modelmi zmien makrochemického zloženia vody a súvisiacej alterácie hornín a alteračných asociácií, vyhodnotením geotermických parametrov, poznaním geologickej stavby a hydrogeologických pomerov.

Cieľom prezentovanej štúdie je poukázať na základné geotermometrické techniky celosvetovo využívané pri odhade rezervoárovej teploty a konštrukcii koncepčných modelov (hydro)geotermálnych štruktúr s ich následnou aplikáciou v lokálnych podmienkach bešeňovskej elevácie.

Hydrogeotermálna štruktúra elevácie Bešeňovej môže byť celkovo charakterizovaná ako zložitý, hydrogeologicky otvorený systém. Koreláciou iónových pomerov s makrochemickými modelmi zmiešavania (prezentované v práci Fričovský a Tometz, 2013) bola identifikovaná infiltračná zóna v okolí Kľačianskej doliny, čo je v súlade s lokálnymi hydraulickými podmienkami a distribúciou piezometrických napätí. Zároveň je stále opodstatnený predpoklad kvázi izochemického prítoku termálnych vôd od štruktúry depresie Liptovskej Mary na východe. V okolí Kľačianskej doliny dochádza minimálne k jednej epizóde zmiešavania medzi hlbokým a plytkým infiltračným kanálom, ktorá prebieha pred separáciou zostupných, dotačných od výstupných a výverových smerov (obr. 29). Termálne vody v hlbokom obehu dosahujú saturáciu voči kalcitu ešte pred vstupom do produkčnej časti rezervoáru (obr. 18) v centrálnej tranzitno-akumulačnej zóne. Zvyšovanie obsahu CO₂ podmieňuje modifikáciu Na-silikátov na K-silikáty, preferenčne disociovaných zo siliciklastických prímiesí rezervoárových karbonátov (obr. 31). Výsledky multikomponentovej geotermometrie (tab. 5 – 6; obr. 27 – 28), analýzy saturačných stavov (obr. 20

a 24) a alteračných systémov (obr. 10 – 13) poukazuje na možnú aktiváciu nízko-teplotnej alterácie rezervoárových dolomitov so siliciklastickými prímiesami a polohami, zvyšovaním reaktivity talku pri rozpúšťaní Mg-kalcitu, presýtení termálnych vôd voči dolomitu a produkcii CO_2 .

Celková minimálna vyzrelosť hlbokých termálnych vôd je následne indikovaná nízkym iónovým pomerom Na/K a modifikáciou termodynamických reaktivít alkalických silikátov v súvislosti s vertikálnou filtráciou v izolovaných konvekčných bunkách pod vplyvom nerovnomerného prehrievania bázy (obr. 30). Alterácia na úrovni termodynamickej aktivity, respektíve čiastočnej ionovýmeny je podporovaná celkovou saturáciou termálnych vôd voči chalcedónu. Termálne vody vystupujú zároveň po otvorených zlomových systémoch v smere k plytkému rezervoáru, kde dochádza k ich zmiešavaniu s termálnymi vodami plytšieho obehu. Časť zmiešaných vôd následne pokračuje po tektonickom výstupe k povrchu, kde vyvierajú v podobe termálnych prameňov a podieľajú sa na tvorbe travertínových štruktúr (obr. 32). V prostredí sekundárnej výverovej zóny Vyšných Sliačov predpokladáme, že k prvej etape zmiešavania dochádza medzi termálnymi vodami z oboch rezervoárov pred ich následným zmiešavaním a nariedením plytkými podzemnými vodami (obr. 33). Rýchly výstup predovšetkým z hlbokého rezervoáru po otvorených zlomových systémoch spôsobuje prudký pokles parciálneho tlaku CO_2 , čo podmieňuje vznik suchých výronov plynov dokumentovaných v minulosti, k čomu prispieva aj sekundárne odpľňovanie po prudkom ochladení nariedením. Kým časť termálnych vôd z hlbokého rezervoáru vyviera v podobe termálnych prameňov v okolí Vyšných Sliačov, časť

termálnych vôd pokračuje v hlbokom prúdení v smere k depresii Ivachnovej (obr. 30).

Aplikácia geotermometrov pri odhade rezervoárovej teploty, eventuálne teploty posledného vyváženia termálnym vodám voči jednotlivým termodynamickým systémom, bola ovplyvnená termodynamickou nevyzrelosťou termálnych vôd (obr. 2, 16 – 17). Presvedčivé výsledky (obr. 9) boli dosiahnuté pri geochemickom modelovaní nasýtenia termálnych vôd štruktúry voči chalcedónu (obr. 4 – 5).

V podmienkach hlbokého rezervoáru sa ako perspektívny geotermometer ukázala Mg-korekcia Na-K-Ca geotermometra (obr. 22). Multikomponentová geotermometria poukázala na možnosť krátkodobého, celkového vyváženia termálnych vôd alebo prítomnosť geochemicky vyzretých termálnych vôd v hlbších častiach rezervoáru.

Na základe zložitosti nového predkladaného modelu štruktúry termálnych vôd, zreteľne odlišnej od predchádzajúceho modelu jednoduchej infiltrácie a výveru (napr. Fendek et al., 1988) odporúčame pozornosť zamerať na určenie lokálnych geochemických geoindikátorov vychádzajúcich z chalcedónu, alebo sadrovca.

Okrem kvantitatívneho využitia pri dlhodobom monitoringu termálnych vôd a rezervoárovej odozvy je zároveň možné tieto geoindikátory s určitou istotou využiť pri upresňovaní koncepčného modelu, definovaní koncepčnej histórie termálnych vôd a interpretácii vzájomných, genetických vzťahov termálnych vôd s termálnymi vodami Liptovskej kotliny.

Využitie geotermometrov preukázalo svoje opodstatnenie aj v podmienkach nízkoentálnych a karbonátových štruktúr, predovšetkým pri tvorbe koncepčného modelu a ako podporná metóda geochemického modelovania.

APPENDIX A – SILICA GEOTHERMOMETERS

Phase	Geothermometer	Author	Model	Code
Amorphous silica	$[731/(4.52 - \log CS)] - 273.15$	Fournier, 1977	conductive	GSA1
	$121.6 + 0.2694CS - 1.8101 \cdot 10^{-4}CS^2 + 7.5521 \cdot 10^{-8}CS^3 + 55.184 \log CS$	Gunarsson and Arnórsson, 2000	conductive	GSA2
Moganite	$30.7 + 0.53113CS + 1.2578 \cdot 10^{-4}CS^2 - 5.9241 \cdot 10^{-7}CS^3 + 19.57 \log CS$	Gislason et al., 1997	conductive	CGM
Chalcedony	$[1\ 032/(4.69 - \log CS)] - 273.15$	Fournier, 1977	conductive	GSC1
	$[1\ 021/(4.69 - \log CS)] - 273.15$	Arnórsson et al., 1983	conductive	GSC2
	$[1\ 112/(4.91 - \log CS)] - 273.15$	Fournier, 1991	conductive	GSC3
	$[1\ 264/(5.31 - \log CS)] - 273.15$	Arnórsson et al., 1983	adiabatic	GSC4A
Quartz	$[1\ 309/(5.198 - \log CS)] - 273.15$	Fournier, 1977	conductive	GSQ1
	$[1\ 315/(5.205 - \log CS)] - 273.15$	Truesdell, 1975	conductive	GSQ2
	$[1\ 157.7/(4.88 - \log CS)] - 273.15$	Verma, 2011	conductive	GSQ3
	$42.2 + 0.28831CS - 3.6686 \cdot 10^{-4}CS^2 + 3.1665 \cdot 10^{-7}CS^3 + 88.390 \log CS$	Fournier and Potter, 1982	conductive	GSQ4
	$55.3 + 0.3659CS - 5.3954 \cdot 10^{-4}CS^2 + 5.5132 \cdot 10^{-7}CS^3 + 74.360 \log CS$	Arnórsson et al., 1988	conductive	GSQ5
	$44.119 + 0.24469mS - 1.7414 \cdot 10^{-4}mS^2 + 79.305 \log mS$	Verma and Santoyo, 1997	conductive	GSQ6
	$66.9 + 0.1378CS - 4.9727 \cdot 10^{-5}CS^2 + 1.0468 \cdot 10^{-8}CS^3 + 87.841 \log CS$	Arnórsson et al., 1988	adiabatic	GSQ7A
	$[1\ 522/(5.75 - \log CS)] - 273.15$	Fournier, 1973	adiabatic	GSQ8A
	$[1\ 164/(4.9 - \log CS)] - 273.15$	Arnórsson et al., 1983	adiabatic	GSQ9A
	$[1\ 264/(5.81 - \log CS)] - 273.15$	Arnórsson et al., 1983	adiabatic	GSQ10A
	$[1\ 498/(5.7 - \log CS)] - 273.15$	Arnórsson et al., 1983	adiabatic	GSQ11A
Opal	$[1\ 000/(4.84 - \log CS)] - 273.15$	Fournier, 1977	conductive	GSOL
	$[781/(4.51 - \log CS)] - 273.15$	Fournier, 1977	conductive	GSOH

Explanations: CS – concentration of SiO₂ in mg·kg⁻¹; mS – concentration of SiO₂ in mol·l⁻¹

APPENDIX B – CATION GEOTHERMOMETERS

Phase	Geothermometer	Author	Code
Na/K	$[777/(0.7 + \log \text{NK})] - 273.15$	Fournier and Truesdell, 1973	GNK1
	$[856/(0.857 + \log \text{NK})] - 273.15$	Truesdell, 1976	GNK2
	$[1\,217/(1.483 + \log \text{NK})] - 273.15$	Fournier, 1979	GNK3
	$[883/(0.78 + \log \text{NK})] - 273.15$	Tonani, 1980	GNK4
	$[933/(0.993 + \log \text{NK})] - 273.15$	Arnórsson et al., 1983	GNK5
	$[1\,319/(1.699 + \log \text{NK})] - 273.15$	Arnórsson et al., 1983	GNK6
	$[1\,178/(1.47 + \log \text{NK})] - 273.15$	Nieva and Nieva, 1987	GNK7
	$[1\,390/(1.75 + \log \text{NK})] - 273.15$	Giggenbach, 1988	GNK8
	$[1\,289/(1.615 + \log \text{NK})] - 273.15$	Verma and Santoyo, 1997	GNK9
	$733.6 - 770.55\log Y + 378.189\log Y^2 - 95.753\log Y^3 + 9.544\log Y^4$	Arnórsson et al., 1988	GNK10
	$[876/(0.877 + \log \text{NK})] - 273.15$	Santoyo and Díaz-González, 2010	GNK11
K ² /Mg	$[2\,330/(7.35 + \log \text{KM})] - 273.15$	Fournier, 1991*	GKM1
	$[1\,077/(4.03 + \log \text{KM})] - 273.15$	Fournier, 1991**	GKM2
	$[4\,410/(14 + \log \text{KM})] - 273.15$	Giggenbach, 1988	GKM3
Na/Li	$[1\,590/(0.770 + \log \text{NL})] - 273.15$	Kharaka et al., 1982	GNL1
	$[1\,000/(0.389 + \log \text{NL})] - 273.15$	Fouillac and Michard, 1981***	GNL2
	$[1\,195/(0.13 + \log \text{NL})] - 273.15$	Fouillac and Michard, 1981****	GNL3
	$[1\,049/(0.44 + \log \text{NL})] - 273.15$	Verma and Santoyo, 1997***	GNL4
	$[1\,276/(0.07 + \log \text{NL})] - 273.15$	Verma and Santoyo, 1997****	GNL5
	$[2\,002/(1.322 + \log \text{NL})] - 273.15$	Sanjuan, 2010	GNL6
Li/Mg ^{0.5}	$[2\,200/(5.47 - \log \text{LM})] - 273.15$	Kharaka and Mariner, 1989	GLM
Na/Ca ^{0.5}	$[1\,096.7/(3.08 - \log \text{NC})] - 273.15$	Tonani, 1980	GNC
K/Ca ^{0.5}	$[1\,930/(3.861 - \log \text{KC})] - 273.15$	Tonani, 1980	GKC
Na-K-Ca	$[1\,647/(\log \text{NK} + \beta \log \text{CN} + 2.24)] - 273.15$		
	$\beta = 1/3 \text{ for } T_{\text{samp}} < 100\,^{\circ}\text{C}$		
	$\beta = 4/3 \text{ for } T_{\text{samp}} > 100\,^{\circ}\text{C}/\log \text{Ca}^{0.5}/\text{Na} < 0$	Fournier and Truesdell, 1973	GNKC

Explanations: NK – Na/K in mg·kg⁻¹; KM – K²/Mg in mg·kg⁻¹; Y – Na/K in mol·l⁻¹; NL – Na/Li in mg·kg⁻¹; LM – Li/Mg^{0.5} in mg·kg⁻¹; NC – Na/Ca^{0.5} in mg·kg⁻¹; KC – K/Ca^{0.5} in mg·kg⁻¹; CN – Ca^{0.5}/Na in mg·kg⁻¹; * – for $\log (\text{K}^2/\text{Mg}) > 1.25$ in mg·kg⁻¹; ** – for $\log (\text{K}^2/\text{Mg}) < 1.25$ in mg·kg⁻¹; *** – for mCl < 0.3 mol·kg⁻¹ in mol·kg⁻¹; **** – for mCl > 0.3 mol·kg⁻¹ in mol·kg⁻¹

APPENDIX C – DESCRIPTION OF SAMPLES (part 1)

Well, spring	Sample code	T _{samp}	pH	Ca	Mg	Na	K	HCO ₃	Cl	SO ₄	Li	SiO ₂	fCO ₂	TDS	REF
		°C	–	all concentrations in mg·l ⁻¹											
ZGL-1	z/21/2/87	61	6.8	510	159	30	23	707	18	1 384	2.2	35	400	2840	1
ZGL-1	z/4/3/87a	62	6.6	510	190	26	16	647	14	1 558	2.2	42	410	2978	1
ZGL-1	z/4/3/87b	61	6.1	510	159	30	23	695	19	1 397	2.5	80	1 307	2 844	1
ZGL-1	z/5/3/87	62	6.1	510	159	30	23	695	12	1 401	2.4	60	690	2 840	1
ZGL-1	z/6/3/87a	61	6.3	510	139	30	23	683	16	1 342	2	50	870	2 753	1
ZGL-1	z/6/3/87b	62	6.3	500	145	30	23	683	15	1 327	2.2	60	786	2 731	1
ZGL-1	z/7/3/87	62	6	500	148	30	23	695	15	1 324	2.1	80	650	2 743	1
ZGL-1	z/22/3/87	62	6.6	506	153	36	23	695	12	1 377	2.1	45	452	2 820	1
ZGL-1	z/31/3/87	53	6.3	473	224	28	22	610	9	1 641	2.2	45	660	2 033	1
ZGL-1	z/2/4/87	61	6.4	495	160	30	22	683	14	1 408	2.1	45	645	2 838	1
ZGL-1	z/3/4/87	62	6.4	495	199	30	22	683	9	1 558	2.1	45	647	3 021	1
ZGL-1	z/20/2/92	51	7	548	145	16	12	653	4	1 433	1.4	50	484	2 878	2
ZGL-1	z/31/10/97	61	6.4	553	138	17	24	711	6	1 376	0.3	20	230	2 867	2
ZGL-1	z/27/11/97	61	6.4	546	150	18	25	787	4	1 370	0.3	34	420	2 959	2
ZGL-1	z/16/12/97	61	6.3	534	156	17	23	730	7	1 369	0.2	27	290	2 897	2
ZGL-1	z/21/8/91	61	6.3	597	130	18	23	760	6	1 371	0.3	24	400	2 920	2
ZGL-1	z/25/2/98	61	6.4	550	130	20	23	757	5	1 432	0.2	27	350	2 897	2
ZGL-1	z/27/3/98	61	6.3	567	141	20	27	766	4	1 360	0.3	27	320	2 905	2
ZGL-1	z/30/4/98	61	6.4	594	124	21	27	725	5	1 374	0.3	28	660	2 865	2
ZGL-1	z/17/5/98	61	6.4	594	142	20	28	725	5	1 374	0.3	23	660	2 860	2
ZGL-1	z/28/5/98	61	6.4	582	128	23	28	757	5	1 375	0.3	23	367	2 917	2
ZGL-1	z/11/6/98	61	6.3	547	134	19	27	720	3	1 355	0.3	29	558	2 823	2
ZGL-1	z/12/11/07	60	6.6	612	144	25	20	900	5	1 380	0.3	37	757	3 101	3
ZGL-1	z/17/12/07	61	6.6	609	153	28	25	915	3	1 470	0.3	45	751	3 225	4
ZGL-1	z/18/2/08	61	6.9	609	158	31	25	909	5	1 400	0.3	39	1 300	3152	4
ZGL-1	z/26/2/09	61	6.8	611	155	31	25	927	5	1 380	0.3	40	1 300	2 662	4
ZGL-1	z/30/3/11	61	6.9	611	153	33	26	988	5	1 420	0.3	44	751	3 253	5
ZGL-1	z/13/4/11	61	6.8	609	159	34	26	990	5	1 440	0.3	40	684	3 278	5
ZGL-1	z/7/3/87/1390	63	5.6	500	165	30	23	61	15	1 418	2.5	60	700	2 830	1
ZGL-1	z/19/3/87/1390	63	5.7	517	157	40	29	708	10	1 377	3.3	70	750	2 810	1
ZGL-1	z/2/4/89/1390	62	6	506	232	50	33	707	18	1 653	2.6	65	770	3 164	1
ZGL-1	z/4/3/87/1510	64	5.8	540	190	28	23	1 000	26	1 342	2.5	60	800	3 162	1
ZGL-1	z/21/3/87/1510	64	6.4	506	173	28	23	700	10	142	2	70	750	2 858	1
ZGL-1	z/23/3/87/1510	63	6.6	517	196	30	25	1 061	15	1 255	2.1	70	800	3 088	1
ZGL-1	z/1/4/87/1510	64	6.4	440	272	38	30	988	17	1 436	2	60	750	3 202	1
ZGL-1	z/6/3/87/1730	60	5.9	570	300	26	23	2 123	38	991	2	50	760	4 110	1
ZGL-1	z/22/3/87/1730	68	6.9	341	528	30	27	2 647	12	914	3	55	720	4 493	1
ZGL-1	z/1/4/87/1730	69	6.6	396	495	32	29	2 501	24	1 006	2.5	57	590	4 468	1
ZGL-1	z/6/3/87/1880	69	5.6	510	159	30	23	683	18	1 416	2.5	54	630	2 853	1
ZGL-1	z/23/3/87/1880	69	6.8	572	378	30	25	2 623	14	868	2.9	56	650	4 496	1
ZGL-1	z/1/4/87/1880	69	6.7	231	605	35	27	2 647	24	932	2.3	53	612	4 488	1
FGTB-1	f/15/3/11	61	6.5	568	136	15	44	695	34	1 480	0.2	55	768	3 021	5
FGTB-1	f/30/3/11	61	6.5	566	147	15	30	720	13	1 500	0.2	58	770	3 017	5
FGTB-1	f/13/4/11	61	6.6	563	156	15	26	714	8	1 480	0.2	54	628	3 008	5

Explanations: REF (References): 1 – Fendek et al. (1988); 2 – Remšík et al. (1998); 3 – Hanzel et al. (2008); 4 – Vandrová et al. (2009); 5 – Vandrová et al. (2011)

APPENDIX C – DESCRIPTION OF SAMPLES (part 2)

Well, spring	Sample code	T _{sample}	pH	Ca	Mg	Na	K	HCO ₃	Cl	SO ₄	Li	SiO ₂	fCO ₂	TDS	REF
		°C	–	all concentrations in mg·l ⁻¹											
BEH-1	B/8/7/74	34	7.2	625	150	22	14	1 159	7	1 238	0.4	17	1 476	3 236	6
BEH-1	B/13/7/74	33	7.1	625	145	30	14	1 135	7	1 258	0.4	15	1 729	3 233	6
BEH-1	B/18/7/74	33	6.9	615	141	64	13	1 135	11	1 284	0.5	15	1 654	3 278	6
BEH-1	B/23/10/74	34	6.5	601	150	61	33	1 208	11	1 230	0.6	10	1 757	3 305	6
BEH-1	B/18/5/74	32	6.6	600	132	117	38	1 336	21	1 180	0.4	15	1 826	3 430	6
BEH-1	B/30/8/74	33	6.4	600	128	131	36	1 342	21	1 186	0.3	16	1 823	3 450	6
BEH-1	B/12/9/74	34	6.5	650	154	73	33	1 445	37	1 168	0.4	3	1 580	3 568	6
Fbe-1	F/12/9/09	26	6.6	585	172	103	36	1 740	47	833	0.5	27	1 812	3 536	4
Fbe-1	F/26/2/09	26	6.4	593	170	102	37	1 763	47	830	0.5	27	1 621	3 564	4
Fbe-1	F/11309	27	6.6	593	167	101	39	1 793	47	843	0.5	30	1 398	3 606	4
Fbe-1	F/30/3/11	26	6.7	585	162	100	39	1 769	45	863	0.5	27	1 186	3 580	4
Fbe-1	F/13/4/11	25	6.8	588	167	100	39	1 757	44	873	0.5	25	1265	3 584	4
LM-45	Lm/15/7/87	22	6.3	670	284	51	16	2 136	26	844	0.4	20	1 367	3 909	7
LM-45	Lm/241085	20	6.6	682	197	53	17	2 172	24	824	0.4	37	1 781	4 009	7
LM-45	Lm/30/10/90	20	6.5	703	192	54	17	2 178	24	825	0.4	38	1 586	4 034	7
LM-140	Lm/27/8/5	28	6.1	668	197	47	17	2 111	25	818	0.4	23	1 523	3 024	7
LM-140	lm/15/7/85	30	6.1	649	186	47	16	2 025	24	808	0.4	22	1 254	3 280	7
LM-140	lm/9/10/90	29	6.6	644	195	51	17	2 056	23	979	0.4	40	1 736	3 827	7
RUDOLF	r/15/9/87	27	6.3	648	168	43	17	1 971	24	751	0.4	21	745	3 660	8
RUDOLF	r/28/10/87	28	6.3	621	181	37	11	1 990	23	851	0.2	15	350	3 783	8
RUDOLF	r/24/10/87	29	6.8	643	197	48	16	2 062	22	790	0.3	36	1 192	3 819	8
RUDOLF	r/30/10/90a	29	6.7	643	192	49	16	2 044	23	781	0.4	37	1 225	3 770	7
RUDOLF	r/30/10/90b	29	6.4	672	194	45	17	2 062	24	836	0.5	17	1 250	3 856	7
RUDOLF	r/30/10/90c	28	6.3	622	181	51	17	1 970	23	851	0.4	15	632	3 720	7
VSH-1	Vsh/1	31	7	397	153	171	33	1 756	70	485	1.3	30	479	3 066	4
VSH-1	Vsh/2	33	6	408	153	163	33	1 770	67	487	1.2	30	693	3 080	4
VSH-1	Vsh/3	30	7	407	153	152	33	1 745	63	484	1.2	30	782	3 042	4
VSH-1	Vsh/4	31	7	448	168	167	36	1 920	69	532	1.3	33	860.2	3 384	4
VSH-1	Vsh/5	33	6	429	165	185	36	1 896	76	524	1.4	32	517.3	3 356	4
VSH-1	Vsh/6	30	7	380	147	165	32	1 684	68	465	1.3	29	434.2	2 981	4
VSH-1	Vsh/7	31	7	395	153	172	33	1 751	70	484	1.3	30	451.5	3 100	4
VSH-1	Vsh/8	33	6	369	142	159	31	1 633	65	451	1.2	28	445.5	2 890	4
VSH-1	Vsh/9	30	7	403	156	174	34	1 785	71	493	1.3	30	484.5	3 158	4
Sliač	s/spring/3	31	7	426	156	135	34	1 768	57	493	1.1	31	1 125	3 105	9
Sliač	s/spring/4	33	6	392	143	120	31	1 616	51	451	1	28	1 086	2 837	9
Sliač	s/spring/5	30	7	388	145	142	31	1 656	59	460	1.1	29	788.4	2 918	9
Štiavnička	s/spring/6	9	7.1	133	31	22	4	391	50	133	0.1	0.2	22	766	9
Sliač	s/spring/7	12	7.1	77	23	13	1	317	10	42	0.1	0.5	11	484	9
Sliač	s/spring8	11	7	74	20	9	1.2	287	7	45	0.1	0.5	26	443	9
Sliač	s/spring9	16	7	75	11	9	0.4	269	6	11	0.1	0.1	14	410	9
Sliač	s/spring/1	20	7.1	475	142	97	35	1 770	56	470	0.7	5	1 580	3 052	9
Sliač	s/spring/2	16	7	603	180	53	19	1 598	27	1 040	0.8	7	1 360	3 530	9

Explanations: REF (References): 4 – Vandrová et al. (2009); 6 – Franko et al. (1979); 7 – Klagó and Stuchlíková (1990); 8 – Vandrová and Matejčková (1991); 9 – Zbořil et al. (1992)

APPENDIX D – SILICA GEOTHERMOMETRY RESULTS (part 1)

Well, spring	Sample code	Silica geothermometer temperature estimate (°C)																			
	GSA1	GSA2	GSM	GSOL	GSOH	GSC1	GSC2	GSC3	GSC4A	GSQ1	GSQ2	GSQ3	GSQ4	GSQ5	GSQ6	GSQ7A	GSQ8A	GSQ9A	GSQ10A	GSQ11A	
ZGL-1	z/21/2/87	–	–	18	28	–	55	51	57	62	85	86	108	104	72	11	73	89	74	62	87
ZGL-1	z/4/3/87a	–	–	24	35	–	63	60	65	70	93	94	118	113	80	0	81	96	82	70	94
ZGL-1	z/4/3/87b	6	4	50	64	26	97	93	97	98	124	125	159	147	112	–	111	122	115	98	12
ZGL-1	z/5/3/87	–	–	36	50	13	81	77	82	85	110	111	140	131	97	–	97	110	100	85	109
ZGL-1	z/6/3/87a	–	–	29	42	5	72	68	73	77	101	102	129	122	88	–	89	103	90	77	101
ZGL-1	z/6/3/87b	–	–	36	50	13	81	77	82	85	110	111	140	131	97	–	97	110	100	85	109
ZGL-1	z/7/3/87	6	4	50	64	26	97	93	97	98	124	125	159	147	112	–	111	122	115	98	121
ZGL-1	z/22/3/87	–	–	26	38	0	67	63	68	73	96	97	122	116	83	–	84	98	85	73	97
ZGL-1	z/31/3/87	–	–	26	38	0	67	63	68	73	96	97	122	116	83	–	84	98	85	73	97
ZGL-1	z/2/4/87	–	–	26	38	0	67	63	68	73	96	97	122	116	83	–	84	98	85	73	97
ZGL-1	z/3/4/87	–	–	26	38	0	67	63	68	73	96	97	122	116	83	–	84	98	85	73	97
ZGL-1	z/20/2/92	–	–	29	42	5	72	68	73	77	101	102	129	122	88	–	89	103	90	77	101
ZGL-1	z/12/11/07	–	–	26	38	0	67	63	68	73	96	97	122	116	83	–	84	98	85	73	97
ZGL-1	z/17/12/07	–	–	21	32	–	60	56	62	67	90	91	114	109	76	5	78	93	79	67	91
ZGL-1	z/18/2/08	–	–	22	33	–	61	57	63	68	91	92	116	110	78	3	79	94	80	68	92
ZGL-1	z/26/2/09	–	–	25	37	–	66	62	67	72	95	96	121	115	82	–	83	97	84	72	96
ZGL-1	z/30/3/11	–	–	22	33	–	61	57	63	68	91	92	116	110	78	3	79	94	80	68	92
FGTB-1	f/15/3/11	–	–	33	46	9	77	73	78	81	105	106	135	126	93	–	93	106	95	81	105
FGTB-1	f/30/3/11	–	–	35	49	11	79	76	80	83	108	109	138	129	95	–	96	109	98	83	107
FGTB-1	f/13/4/11	–	–	32	46	8	76	72	77	80	105	106	133	125	92	–	93	106	94	80	104
ZGL-1	z/7/3/87/1390	–	–	36	50	13	81	77	82	85	110	111	140	131	97	–	97	110	100	85	109
ZGL-1	z/19/3/87/1390	0	–	43	57	20	90	86	90	92	117	118	150	139	105	–	105	117	108	92	115
ZGL-1	z/2/4/89/1390	–	–	40	54	16	86	82	86	88	114	115	145	135	101	–	101	113	104	88	112
ZGL-1	z/4/3/87/1510	–	–	36	50	13	81	77	82	85	110	111	140	131	97	–	97	110	100	85	109
ZGL-1	z/21/3/87/1510	0	–	43	57	20	90	86	90	92	117	118	150	139	105	–	105	117	108	92	115
ZGL-1	z/23/3/87/1510	0	–	43	57	20	90	86	90	92	117	118	150	139	105	–	105	117	108	92	115
ZGL-1	z/1/4/87/1510	–	–	36	50	13	81	77	82	85	110	111	140	131	97	–	97	110	100	85	109
ZGL-1	z/6/3/87/1730	–	–	29	42	5	72	68	73	77	101	102	129	122	88	–	89	103	90	77	101
ZGL-1	z/22/3/87/1730	–	–	33	46	9	77	73	78	81	105	106	135	126	93	–	93	106	95	81	105
ZGL-1	z/1/4/87/1730	–	–	34	48	10	79	75	79	82	107	108	137	128	94	–	95	108	97	82	107
ZGL-1	z/6/3/87/1880	–	–	32	46	8	76	72	77	80	105	106	133	125	92	–	93	106	94	80	104
ZGL-1	z/23/3/87/1880	–	–	34	47	10	78	74	79	82	106	107	136	127	94	–	94	107	96	82	106
ZGL-1	z/1/4/87/1880	–	–	31	45	7	75	71	76	79	104	105	132	125	91	–	92	105	93	79	104

APPENDIX D – SILICA GEOTHERMOMETRY RESULTS (part 2)

Well, spring	Sample code	Silica geothermometer temperature estimate (°C)																			
	GSA1	GSA2	GSM	GSOL	GSOH	GSC1	GSC2	GSC3	GSC4A	GSQ1	GSQ2	GSQ3	GSQ4	GSQ5	GSQ6	GSQ7A	GSQ8A	GSQ9A	GSQ10A	GSQ11A	
BEH-1	B/8/7/74	–	–	2	2	–	25	22	29	37	57	58	72	71	42	29	44	64	44	37	62
BEH-1	B/13/7/74	–	–	–	–	–	21	17	25	33	52	53	67	66	38	29	38	60	39	33	58
BEH-1	B/18/7/74	–	–	–	–	–	21	17	25	33	52	53	67	66	38	29	38	60	39	33	58
BEH-1	B/23/10/74	–	–	–	–	–	7	4	11	20	39	40	50	49	23	27	22	47	25	20	46
BEH-1	B/18/5/74	–	–	–	–	–	21	17	25	33	52	53	67	66	38	29	38	60	39	33	58
BEH-1	B/30/8/74	–	–	1	–	–	23	20	27	35	55	56	70	69	40	29	41	62	42	35	60
BEH-1	B/12/9/74	–	–	–	–	–	15	12	20	28	47	48	61	60	32	29	33	55	34	28	53
Fbe-1	F/12/9/09	–	–	12	18	–	44	40	47	53	74	75	95	92	61	21	63	79	62	53	78
Fbe-1	F/26/2/09	–	–	12	18	–	44	40	47	53	74	75	95	92	61	21	63	79	62	53	78
Fbe-1	F/11309	–	–	14	22	–	48	45	51	57	79	80	100	97	65	17	67	83	67	57	82
Fbe-1	F/30/3/11	–	–	12	18	–	44	40	47	53	74	75	95	92	61	21	63	79	62	53	78
Fbe-1	F/13/4/11	–	–	10	15	–	40	37	43	50	71	72	91	88	57	23	59	77	59	50	75
LM-45	Lm/15/7/87	–	–	5	7	–	31	28	35	42	63	64	80	78	49	27	50	69	50	42	67
LM-45	Lm/241085	–	–	20	30	–	57	54	60	65	87	88	111	107	74	8	76	91	76	65	89
LM-45	Lm/30/10/90	–	–	21	31	–	59	55	61	66	89	90	113	108	75	6	77	92	77	66	90
LM-140	Lm/27/8/5	–	–	8	12	–	37	34	40	47	68	69	87	85	54	25	56	74	56	47	72
LM-140	lm/15/7/85	–	–	7	10	–	35	32	39	45	66	67	84	83	52	26	54	72	54	45	71
LM-140	lm/9/10/90	–	–	22	33	–	61	57	63	68	91	92	116	110	78	3	79	94	80	68	92
RUDOLF	r/15/9/87	–	–	6	9	–	33	30	37	44	65	66	82	81	50	27	52	71	52	44	69
RUDOLF	r/28/10/87	–	–	–	–	–	21	17	25	33	52	53	67	66	38	29	38	60	39	33	58
RUDOLF	r/24/10/87	–	–	19	29	–	56	53	58	64	86	87	110	105	73	9	75	90	75	64	88
RUDOLF	r/30/10/90a	–	–	20	30	–	57	54	60	65	87	88	111	107	74	8	76	91	76	65	89
RUDOLF	r/30/10/90b	–	–	2	2	–	25	22	29	37	57	58	72	71	42	29	44	64	44	37	62
RUDOLF	r/30/10/90c	–	–	–	–	–	21	17	25	33	52	53	67	66	38	29	38	60	39	33	58
VSH-1	Vsh/1	–	–	14	22	–	48	45	51	57	79	80	100	97	65	17	67	83	67	57	82
VSH-1	Vsh/2	–	–	14	22	–	48	45	51	57	79	80	100	97	65	17	67	83	67	57	82
VSH-1	Vsh/3	–	–	14	22	–	48	45	51	57	79	80	100	97	65	17	67	83	67	57	82
VSH-1	Vsh/4	–	–	17	25	–	52	49	55	60	83	84	105	101	69	13	71	87	71	60	85
VSH-1	Vsh/5	–	–	16	25	–	51	48	54	60	82	83	104	100	68	14	70	86	70	60	84
VSH-1	Vsh/6	–	–	13	20	–	46	43	49	55	77	78	98	95	63	19	65	82	65	55	80
VSH-1	Vsh/7	–	–	14	22	–	48	45	51	57	79	80	100	97	65	17	67	83	67	57	82
VSH-1	Vsh/8	–	–	12	19	–	45	42	48	54	76	77	96	93	62	20	64	80	64	54	79
VSH-1	Vsh/9	–	–	15	22	–	49	45	51	57	79	80	101	97	66	17	68	84	68	57	82
VSH-1	Vsh/10	–	–	15	22	–	49	45	52	57	79	80	101	98	66	17	68	84	68	57	82
VSH-1	Vsh/11	–	–	13	19	–	45	42	48	54	76	77	96	94	62	20	64	81	64	54	79
VSH-1	Vsh/12	–	–	13	20	–	46	42	49	55	77	78	97	94	63	19	65	81	65	55	80

APPENDIX E – CATION GEOTHERMOMETRY RESULTS (part 1)

Well, spring	Sample code	Cation geothermometry temperature estimate (°C)																							
	GN1	GN2	GN3	GN4	GN5	GN6	GN7	GN8	GN9	GN10	GN11	GKM1	GKM2	GKM3	GLM	GLN1	GLN2	GLN3	GLN4	GLN5	GNC	GKC	GNKC	GNKCm	
ZGL-1	z/21/2/87	>	>	>	>	>	>	>	>	>	>	>	24	–	36	80	>	–	–	–	–	98	>	>	55
ZGL-1	z/4/3/87a	>	>	>	>	>	>	>	>	>	>	>	38	–	40	78	>	–	–	–	–	90	>	>	39
ZGL-1	z/4/3/87b	>	>	>	>	>	>	>	>	>	>	>	23	–	36	83	>	–	–	–	–	98	>	>	57
ZGL-1	z/5/3/87	>	>	>	>	>	>	>	>	>	>	>	23	–	36	82	>	–	–	–	–	98	>	>	57
ZGL-1	z/6/3/87a	>	>	>	>	>	>	>	>	>	>	>	21	–	35	79	>	–	–	–	–	98	>	>	57
ZGL-1	z/6/3/87b	>	>	>	>	>	>	>	>	>	>	>	21	–	36	81	>	–	–	–	–	98	>	>	57
ZGL-1	z/7/3/87	>	>	>	>	>	>	>	>	>	>	>	22	–	36	80	>	–	–	–	–	98	>	>	57
ZGL-1	z/22/3/87	>	>	>	>	>	>	>	>	>	>	>	22	–	36	79	>	–	–	–	–	>	>	>	59
ZGL-1	z/31/3/87	>	>	>	>	>	>	>	>	>	>	>	30	–	38	76	>	–	–	–	–	96	>	>	38
ZGL-1	z/2/4/87	>	>	>	>	>	>	>	>	>	>	>	24	–	37	79	>	–	–	–	–	99	>	>	61
ZGL-1	z/3/4/87	>	>	>	>	>	>	>	>	>	>	>	28	–	38	76	>	–	–	–	–	99	>	>	54
ZGL-1	z/20/2/92	>	>	>	>	>	>	>	>	>	>	>	44	–	42	70	>	–	–	–	–	65	>	>	60
ZGL-1	z/12/11/07	>	>	>	>	>	>	>	>	>	>	>	20	–	35	37	>	–	–	–	–	89	>	>	69
ZGL-1	z/17/12/07	>	>	>	>	>	>	>	>	>	>	>	20	–	35	37	>	–	–	–	–	95	>	>	73
ZGL-1	z/18/2/08	>	>	>	>	>	>	>	>	>	>	>	20	–	35	37	>	–	–	–	–	95	>	>	69
ZGL-1	z/26/2/09	>	>	>	>	>	>	>	>	>	>	>	18	–	35	37	>	–	–	–	–	98	>	>	72
ZGL-1	z/30/3/11	>	>	>	>	>	>	>	>	>	>	>	19	–	35	37	>	–	–	–	–	>	>	>	69
FGTB-1	f/15/3/11	>	>	>	>	>	>	>	>	>	>	>	1	–	29	31	>	–	–	–	–	61	>	>	65
FGTB-1	f/30/3/11	>	>	>	>	>	>	>	>	>	>	>	13	–	33	30	>	–	–	–	–	61	>	>	94
FGTB-1	f/13/4/11	>	>	>	>	>	>	>	>	>	>	>	19	–	35	30	>	–	–	–	–	61	>	>	82
ZGL-1a	z/7/3/87/1390	>	>	>	>	>	>	>	>	>	>	>	23	–	36	83	>	–	–	–	–	98	>	>	52
ZGL-1a	z/19/3/87/1390	>	>	>	>	>	>	>	>	>	>	>	15	–	34	91	>	–	–	–	–	>	>	>	69
ZGL-1a	z/2/4/89/1390	>	>	>	>	>	>	>	>	>	>	>	17	–	34	80	>	–	–	–	–	>	>	>	35
ZGL-1a	z/4/3/87/1510	>	>	>	>	>	>	>	>	>	>	>	26	–	37	81	>	–	–	–	–	93	>	>	47
ZGL-1	z/21/3/87/1510	>	>	>	>	>	>	>	>	>	>	>	24	–	36	77	>	–	–	–	–	94	>	>	58
ZGL-1	z/23/3/87/1510	>	>	>	>	>	>	>	>	>	>	>	24	–	36	76	>	–	–	–	–	97	>	>	69
ZGL-1	z/1/4/87/1510	>	>	>	>	>	>	>	>	>	>	>	23	–	36	71	>	–	–	–	–	>	>	>	28
ZGL-1	z/6/3/87/1730	>	>	>	>	>	>	>	>	>	>	>	34	–	39	70	>	–	–	–	–	87	>	>	31
ZGL-1	z/22/3/87/1730	>	>	>	>	>	>	>	>	>	>	>	38	–	40	73	>	–	–	–	–	>	>	>	77
ZGL-1	z/1/4/87/1730	>	>	>	>	>	>	>	>	>	>	>	34	–	39	70	>	–	–	–	–	>	>	>	48
ZGL-1	z/6/3/87/1880	>	>	>	>	>	>	>	>	>	>	>	23	–	36	83	>	–	–	–	–	98	>	>	61
ZGL-1	z/23/3/87/1880	>	>	>	>	>	>	>	>	>	>	>	35	–	39	76	>	–	–	–	–	95	>	>	68
ZGL-1	z/1/4/87/1880	>	>	>	>	>	>	>	>	>	>	>	40	–	41	65	>	–	–	–	–	>	>	>	63

APPENDIX E – CATION GEOTHERMOMETRY RESULTS (part 2)

Well, spring	Sample code	Cation geothermometry temperature estimate (°C)																							
	GN1	GN2	GN3	GN4	GN5	GN6	GN7	GN8	GN9	GN10	GN11	GKM1	GKM2	GKM3	GLM	GLN1	GLN2	GLN3	GLN4	GLN5	GNC	GKC	GNKC	GNKCm	
BEH-1	B/8/7/74	>	>	>	>	>	>	>	>	>	>	>	39	–	41	43	>	–	–	–	–	77	>	>	63
BEH-1	B/13/7/74	>	>	>	>	>	>	>	>	>	>	>	38	–	40	43	>	–	–	–	–	92	>	>	63
BEH-1	B/18/7/74	>	>	>	>	>	>	>	>	>	>	>	40	–	41	48	>	–	–	–	–	>	>	>	65
BEH-1	B/23/10/74	>	>	>	>	>	>	>	>	>	>	>	11	–	32	51	>	–	–	–	–	>	>	>	70
BEH-1	B/18/5/74	>	>	>	>	>	>	>	>	>	>	>	5	–	31	44	>	–	–	–	–	>	>	>	79
BEH-1	B/30/8/74	>	>	>	>	>	>	>	>	>	>	>	6	–	31	39	>	–	–	–	–	>	>	>	79
BEH-1	B/12/9/74	>	>	>	>	>	>	>	>	>	>	>	11	–	33	43	>	–	–	–	–	>	>	>	66
Fbe-1	F/12/9/09	>	>	>	>	>	>	>	>	>	>	>	10	–	32	46	>	–	–	–	–	>	>	>	66
Fbe-1	F/26/2/09	>	>	>	>	>	>	>	>	>	>	>	9	–	32	46	>	–	–	–	–	>	>	>	62
Fbe-1	F/11/3/09	>	>	>	>	>	>	>	>	>	>	>	7	–	31	47	>	–	–	–	–	>	>	>	67
Fbe-1	F/30/3/11	>	>	>	>	>	>	>	>	>	>	>	7	–	31	47	>	–	–	–	–	>	>	>	66
Fbe-1	F/13/4/11	>	>	>	>	>	>	>	>	>	>	>	7	–	31	47	>	–	–	–	–	>	>	>	66
LM-45	Lm/15/7/87	>	>	>	>	>	>	>	>	>	>	>	46	–	42	37	>	–	–	–	–	>	>	>	33
LM-45	Lm/241085	>	>	>	>	>	>	>	>	>	>	>	37	–	40	40	>	–	–	–	–	>	>	>	51
LM-45	Lm/30/10/90	>	>	>	>	>	>	>	>	>	>	>	36	–	40	41	>	–	–	–	–	>	>	>	62
LM-140	Lm/27/8/5	>	>	>	>	>	>	>	>	>	>	>	37	–	40	40	>	–	–	–	–	>	>	>	51
LM-140	lm/15/7/85	>	>	>	>	>	>	>	>	>	>	>	38	–	40	41	>	–	–	–	–	>	>	>	49
LM-140	lm/9/10/90	>	>	>	>	>	>	>	>	>	>	>	37	–	40	41	>	–	–	–	–	>	>	>	46
RUDOLF	r/15/9/87	>	>	>	>	>	>	>	>	>	>	>	34	–	39	42	>	–	–	–	–	>	>	>	48
RUDOLF	r/28/10/87	>	>	>	>	>	>	>	>	>	>	>	52	–	44	28	>	–	–	–	–	>	>	>	46
RUDOLF	r/24/10/87	>	>	>	>	>	>	>	>	>	>	>	39	–	41	35	>	–	–	–	–	>	>	>	47
RUDOLF	r/30/10/90a	>	>	>	>	>	>	>	>	>	>	>	39	–	40	41	>	–	–	–	–	>	>	>	51
RUDOLF	r/30/10/90b	>	>	>	>	>	>	>	>	>	>	>	37	–	40	45	>	–	–	–	–	>	>	>	48
RUDOLF	r/30/10/90c	>	>	>	>	>	>	>	>	>	>	>	35	–	40	41	>	–	–	–	–	>	>	>	41
VSH-1	Vsh/1	>	>	>	>	>	>	>	>	>	>	>	11	–	33	68	>	–	–	–	–	>	>	>	49
VSH-1	Vsh/2	>	>	>	>	>	>	>	>	>	>	>	11	–	33	66	>	–	–	–	–	>	>	>	47
VSH-1	Vsh/3	>	>	>	>	>	>	>	>	>	>	>	11	–	33	66	>	–	–	–	–	>	>	>	44
VSH-1	Vsh/4	>	>	>	>	>	>	>	>	>	>	>	9	–	32	67	>	–	–	–	–	>	>	>	44
VSH-1	Vsh/5	>	>	>	>	>	>	>	>	>	>	>	10	–	32	69	>	–	–	–	–	>	>	>	49
VSH-1	Vsh/6	>	>	>	>	>	>	>	>	>	>	>	12	–	33	68	>	–	–	–	–	>	>	>	47
VSH-1	Vsh/7	>	>	>	>	>	>	>	>	>	>	>	11	–	33	68	>	–	–	–	–	>	>	>	48
VSH-1	Vsh/8	>	>	>	>	>	>	>	>	>	>	>	12	–	33	67	>	–	–	–	–	>	>	>	49
VSH-1	Vsh/9	>	>	>	>	>	>	>	>	>	>	>	11	–	32	68	>	–	–	–	–	>	>	>	43
VSH-1	Vsh/10	>	>	>	>	>	>	>	>	>	>	>	11	–	32	64	>	–	–	–	–	>	>	>	48
VSH-1	Vsh/11	>	>	>	>	>	>	>	>	>	>	>	12	–	33	63	>	–	–	–	–	>	>	>	46
VSH-1	Vsh/12	>	>	>	>	>	>	>	>	>	>	>	12	–	33	65	>	–	–	–	–	>	>	>	42

APPENDIX F – LIST OF ACRONYMS AND SYMBOLS IN TEXT

Symbol	Description	Unit (if any)
β	reaction stoichiometry coefficient	–
Δt_{Mg}	Mg-correction temperature	°C
a_k	activity coefficient of phase k	–
$a_{i,k}$	activity coefficient of species i in phase k	–
C	number of components in reaction	–
CSM	conceptual site model	–
CHN	Choč Nappe (system, succession, horizon)	–
cX	concentration of i-th component in water	ppm, mg·l ⁻¹ , mg·kg ⁻¹
D-HTC	discharge zone, high total carbon subfacies	–
D-LTC	discharge zone, low total carbon subfacies	–
DRTE	deep reservoir thermal endmember	–
F	number of independent variables in reactions	–
H	enthalpy (in general)	kJ·kg ⁻¹
HL	enthalpy of liquid	kJ·kg ⁻¹
H_{res}	reservoir enthalpy	kJ·kg ⁻¹
HT	high temperature (conditions)	–
IWCP	Inner Western Carpathian Paleogene (Sub-Tatric Group) succession	–
Jr/Cr-2	(Križna Nappe) Jurassic – Mid Cretaceous succession (duplexes)	–
K_{eq}	equilibrium constant	–
KNA	Križna Nappe (system, succession, horizon)	–
L_x	Logarithm of ionic concentration ratios	–
LT	low temperature (conditions)	–
M	number of equilibrated minerals in solution	–
MCG	multicomponent geothermometers	–
MEG	multicomponent equilibrium geothermometry	–
MI	maturity index	–
P	number of phases in reactions	–
pCO ₂	partial CO ₂ pressure	bar, MPa
Pg	Paleogene	–
p-T	(thermodynamic) pressure – temperature conditions	–
Q_k	activity coefficient for aqueous species of phase k	–
r_x	relative concentrations	–, %
RMED	clustered median (multicomponent geothermometry) approach	–
S-CGW	Sliache springs – cold groundwater facies	–
SI	saturation index	–
SRTW	shallow reservoir thermal waters facies	–
T	temperature (general)	°C
T_{bhm}	bottomhole temperature	°C
TCR	Tatricum Crystalline (bedrock)	–
TEU	Tatricum Envelope Unit (horizon)	–
Tr2	Mid Triassic (carbonates)	–
T_{est}	estimated temperature	°C
T_{res}	reservoir temperature	°C
T_{samp}	sampling temperature	°C
T_{wh}	wellhead temperature	°C
T_x	Geothermometer temperature	°C
TCM	thermal-cold mixing (regime)	–
TCLM	ternary chloride mixing model	–
TTM	thermal-thermal mixing (regime)	–
TSI	total saturation index	–
Tr3-Cr2	(Križna Nappe) Late Triassic – Mid Cretaceous profile	–
$v_{i,k}$	stoichiometric coefficient for species i in phase k	–
X_{CS}	silica fraction correction	–
X_{H}	enthalpy fraction correction	–

Terciárna mikrofauna z hlbokých vrtov v železovskej priehlbine (Dunajská panva)

ADRIENA ZLINSKÁ

Štátny geologický ústav Dionýza Štúra, Mlynská dolina 1, 817 04 Bratislava, Slovensko;
adriena.zlinska@geology.sk

Abstract: The results of the microfauna re-evaluation in the Cenozoic sediments of the Železovce depression of Danube Basin is presented in terms of the chrono- and lithostratigraphic units in problematic sections of two selected deep wells excavated in the 1960s by former Oil company ČND Hodonín. In the past, the oil boreholes were realized with intention to determine the stratigraphy and lithology of the Neogene sediments and to verify their oil and gas carrying capacity. Altogether, 22 samples have been analysed from the boreholes Modrany 1 and Nová Vieska 1.

The borehole Nová Vieska 1, according to our results terminated at a depth of 3 171 m in the latest Rupelian sediments, previously regarded as the Priabonian to Rupelian in age. The microfauna acquired from the well cores gave a proof of the younger Kiscellian to Egerian (Rupelian to Chattian) age. The overlying Miocene sequence was divided into the Badenian and Sarmatian part of the sedimentary record. Moreover, the deposits previously regarded as the Rupelian are of the Lower Badenian age.

Keywords: Danube Basin, Železovce depression, Foraminifera, Oligocene, Miocene, Badenian, Sarmatian, Kiscellian, Bajtava, Pozba and Vráble formations

Graphical abstract



Typical microfauna of the Middle Badenian
Spiroplectammina carinata biozone

Highlights

- 22 re-evaluated samples with 192 foraminifer taxa from two old oil wells in the Danube Basin revealed Badenian and Sarmatian age of Neogene sediments.
- Kiscellian to Egerian ages of Paleogene sediments represent the most striking new findings: Former Upper Eocene (Priabonian) sediments are reclassified to the Lower Oligocene (Kiscellian Fm.) and those of Lower Oligocene (Rupelian) into the Lower Badenian (Bajtava Fm.).
- In the Danube Basin, the Paleogene lithostratigraphic units of the Buda Paleogene (Gyalog and Császár, 1996) were used for the first time.

Úvod


Príspevok vznikol v rámci riešenia projektu geologickej úlohy *Výskum geologickej stavby a zostavenie geologických máp v problematických územiach Slovenskej republiky*.

Predmetom nášho revízneho štúdia boli foraminifery z paleogénnych a neogénnych súvrství v železovskej priehlbine Dunajskej panvy vo vrtach Modrany 1 a Nová Vieska 1 (obr. 1).

Vrt Modrany 1 (M-1) bol projektovaný a realizovaný ČND, závod Gbely. Projektovaná hĺbka vrtu bola 3 100 m

(reálne ukončený v hĺbke 2 294 m v jure). Vrtanie sa uskutočnilo 30. 8. 1968 – 7. 1. 1969. Účelom bolo zistiť litologický vývoj, stratigrafickú príslušnosť neogénnych súvrství a ich podložia a overiť nafto- a plynonosnosť jednotlivých súvrství na elevácii v poklesnutej kryhe.

Vrt Nová Vieska 1 (NV-1) bol projektovaný a realizovaný ČND Hodonín, závod Gbely. Projektovaná hĺbka vrtu bola 3 200 m (reálne ukončený v hĺbke 3 171 m, pôvodne v priabóne, po našej revízii v oligocéne). Vrtanie sa uskutočnilo 23. 10. 1954 – 10. 1. 1957. Účelom bolo zistiť prítomnosť naftonosných formácií hlavne v bádene, faciálny a litologický vývoj naftonosných formácií. Vrt




Obr. 1. Lokalizácia študovaných vrtov.
Fig. 1. Location of studied wells.

ani po prehĺbení projektovanej hĺbky nedosiahol predterciérne podložie a podľa pôvodných údajov sa skončil v sedimentoch priabónu.

Analýzou mikrofauny sme sa snažili prehodnotiť staršie biostratigrafické údaje v problematických úsekoch vrtov, vekovo zaradených do vrchného eocénu, oligocénu a bádenu, a na základe toho usadeniny zaradiť do súčasne platných chrono- a litostratigrafických jednotiek.

Regionálnogeologické členenie Dunajskej panvy a stručná geologická charakteristika územia

Študovaná oblasť je z hľadiska regionálnogeologického členenia (Vass et al., 1988) súčasťou Dunajskej panvy. Tá patrí medzi vnútrohorské panvy a prstovito zasahuje medzi jadrové pohoria Západných Karpát, ktoré vymedzujú jednotlivé čiastkové priehlbiny. Na západe je to *blatniarska priehlbina* vymedzená Malými Karpatmi a Považským Inovcom. *Rišňovská priehlbina* sa nachádza medzi Považským Inovcom a Tribečom, *komjatická priehlbina* je situovaná medzi Tribečom a stre-



Obr. 2. Regionálnogeologické členenie Dunajskej panvy.
Fig. 2. Regional-geological division of the Danube Basin.

doslovenskými vulkanitmi. Na juhu sa nachádza železovská priehlbina, ktorú z juhu vymedzuje koryto Dunaja, na východe dolina Ipl'a, vulkanity pohoria Burda a štúrovský paleogén (morské sedimenty lutétu až egeru). Dunajská panva sa takmer úplne kryje s regionálnou morfológickou jednotkou Podunajskou nížinou, ktorej centrálnu časť tvorí Gabčíkovská panva (obr. 2).

Hlavnú časť sedimentárnej výplne panvy tvoria morske a brakické súvrstvia stredného miocénu (báden – sarmat), ktoré sú prekryté menej hrubými brakickými a sladkovodnými jazernými sedimentmi vrchného miocénu až panónu a pliocénu.

Slovenská časť Dunajskej panvy je súčasťou rozsiahlej panónskej zaoblúkovej, resp. vnútrooblúkovej panvy, ktorú tvorí systém „menších“ čiastkových panví (Royden et al., 1983). Jednotlivé čiastkové panvy prešli počas miocénu a pliocénu čiastočne rozdielnou genézou. Dôsledkom toho sa vzájomne odlišujú nielen celkovou hrúbkou svojej neogénnej sedimentárnej výplne, ale aj chronostratigrafickým sledom súvrství v rámci jednotlivých sedimentárnych sekvencií, zastúpením, resp. absenciou niektorých súvrství, litologickým charakterom a hrúbkou (Kováč, 2000). Tvar slovenskej časti Dunajskej panvy je výsledkom zložitého tektonického a komplexného geologického vývoja v čase a priestore (Adam a Dlabač, 1961, 1969). Za dosiaľ najstaršie zistené neogénne horniny sa považujú vulkanity spodnobádenského veku nazvané šurianske vulkanity (Vass, 2002). Reprezentujú ich amfibolické andezity prechádzajúce do amfibolicko-pyroxenických andezitov (Pálfalvi, 1975). Sedimentácia na väčšine územia sa začína spodnobádenským špačinským súvrstvím (Rybár et al., 2015; NN-5 fide Martini, 1971). Ďalšie údaje o geologickej stavbe panvy sú známe z jej západného okraja, kde sú na kryštálických horninách podložia usadené až sedimenty vrchnobádenského veku, začlenené do báhoňského súvrstvia. Na báze súvrstvia sú svetlosivé a zelenkastosivé brekcie tvorené ostrohrannými úlomkami granitoidov. Na okraji panvy sedimentovali v tomto období biele a sivo-biele piesčité organodetritické vápence (Pálfalvi, l. c.). V ich nadloží sa vyskytujú svetlosivé drobné- až hrubozrnné pieskovce, ktoré smerom do nadložia prechádzajú do sivohnedých bridličnatých ílov. V uvedených íloch sa zistili spoločenstvá foraminifer bolivínovo-bulimínovej zóny morského vrchného báden. Pri západnom okraji panvy sa koncom vrchného báden usadili sladkovodné súvrstvia uhoľných ílov so slojčkami lignitu.

V nadloží bádenských sedimentov sa vyskytujú usadeniny vrábeľského súvrstvia sarmatského veku, deponované v brakickom, prevažne plytkomorskom prostredí. Bazálnu časť súvrstvia tvoria hrubozrnné piesky s vrstvami štrkov, ktorých obliaky sú tvorené granitoidmi a kryštálickými bridlicami. V nadloží štrkov sa nachádzajú zelenosivé piesčité íly, ktoré sú v spodnej časti silno škvŕnité. V okrajovej časti panvy ležia sarmatské sedimenty priamo na kryštá-

lickom podloží (Cílek, 1960). Panvový vývoj vrábeľského súvrstvia tvoria zelenosivé vápnité íly, prachovce a piesky dosahujúce hrúbku 300 – 600 m.

V spodnom až strednom panóne (zóny A – E panónu) prebiehala sedimentácia v Dunajskej panve v brakickom prostredí plytkého jazera, ktoré sa zaplňalo prevažne delťovými usadeninami transportovanými riekami z vyzdvihujúceho sa alpsko-karpatského orogénu. Usadzovali sa sedimenty ivanského súvrstvia, ktoré bolo v okrajovom vývoji reprezentované štrkami, pieskami a filmi s polohami lignitov. V panvovom vývoji boli v najväčšej miere zastúpené íly s polohami pieskov. Vo vrchnom panóne (zóny F – H) až ?ponte sa v prostredí plytkého jazera kaspibrakického charakteru usadzovali sedimenty beladického a volkovského súvrstvia. Reprezentujú ich prevažne sivozelené íly až prachy, v ktorých sa vyskytujú polohy so zuhoľnatenými zvyškami rastlín (vrstvy uhoľných ílov) až slojčkami lignitu ?pontského veku. Bazálne časti sedimentov zóny F volkovského súvrstvia sú miestami zastúpené štrkami, ktoré laterálne prechádzajú do pestrých ílov. Sedimenty zón G – H panónu pozostávajú zo zelenosivých, hrdzavožlté škvŕnitých, v menšej miere aj hnedo škvŕnitých vápnitých ílov, v ktorých sú polohy svetlozelenkavosivých a svetlosivých vápnitých pieskov až drobnozrnných štrkov s úlomkami suchozemských a sladkovodných gastropódov (Lunga, 1965).

V nadloží volkovského súvrstvia v priestore gabčíkovskej panvy sú usadené sladkovodné fluvialno-limnické až fluvialne štrkovo-piesčité až piesčito-ílovité pliocénne sedimenty kolárovskeho súvrstvia.

Kvartérne sedimenty majú na povrchu Dunajskej panvy dominantné postavenie. Vo veľkej hrúbke, ktorá je v centre gabčíkovskej depresie 500 a viac metrov, pokrývajú súvisle celé vymedzené územie. Kvartérna sedimentácia má kontinuálny priebeh a až do recentu vykazuje panvový superpozičný vývoj (Vaškovský a Vaškovská, 1977). Za najstaršie kvartérne sedimenty sa považujú bazálne vrstvy riečno-jazerných sedimentov, ktoré tvoria spodnú časť kvartérnej výplne centra gabčíkovskej depresie. Sedimenty bazálnych vrstiev kvartéru pozostávajú z cyklicky sa striedajúcich pestrých piesčito-štrkovitých vrstiev s častými, veľmi hrubými polohami ílov a siltov a tvoria pravdepodobne kontinuálny prechod z vrchnopliocénnych pieskov, drobných štrkov a pestrých ílov kolárovskeho súvrstvia, označovaných tiež ako gabčíkovské vrstvy. Fluvialne sedimenty staršieho obdobia stredného pleistocénu sa podieľajú na podstatnej časti stavby kvartérnej sedimentárnej výplne a tvoria spodnú časť tzv. stredného komplexu piesčito-štrkového stredného súvrstvia. V podstate ide o obrovský plochý vejár náplavového kužeľa Dunaja, zakrytý pokračujúcou fluvialnou sedimentáciou vrchného pleistocénu a holocénu. Toto stredné súvrstvie (stredný komplex) označil Janáček (1967) ako „dunajská štrková séria“. Leží na spodnom fluvialno-limnickom až fluvial-

nom palkovičovskom súvrství, od ktorého je oddelené hlinito-ílovitými polohami. Stredné súvrstvie tvoria prevažne hrubšie, dobre opracované štrky, piesčité štrky a piesky. Na rozdiel od staršieho spodného súvrstvia, zriedkavejšie sa v ňom vyskytujú hrubé polohy ílovitých, resp. hlinitých sedimentov. Hlinito-ílovité vrstvy sú po zrnitosti stránke rozmanité, od piesčitých a prachovitých hlin po ílovité hlíny a íly. V mladšej časti stredného pleistocénu kontinuálne pokračoval synsedimentárny pokles centrálnej gabčíkovskej priehlbiny. To sa dotýkalo aj jej menej intenzívne poklesávajúcich okrajových častí. Hrúbka strednej časti stredného súvrstvia nie je rovnaká.

Vrchnopleistocénne fluviálne akumulácie tvoria najvyššiu časť stredného súvrstvia, resp. stredného komplexu dunajskej štrkovej série a na okraji roviny budujú podložie riečnych nív Dunaja, Malého Dunaja, Čiernej vody, Dudváhu, Váhu a Nitry. Litologicky ich tvoria zvodnené, dobre vytriedené a opracované, prevažne hrubšie piesčité štrky a piesky, ktoré dominujú hlavne na povrchu. Štrkový materiál sa vyznačuje charakteristickým hrdzavohnedým až hnedožltým a sivým sfarbením. Fluviálne sedimenty stredného a vrchného pleistocénu v terasovom vývoji vystupujú na ľavobreží Dunaja v Bratislave a na okrajoch pahorkatín, kde sú pokryté hrubými vrstvami vrchnopleistocénnych spraší a sprašových hlin.

Sedimenty holocénu sú reprezentované najmä fluviálnymi akumuláciami litofaciálne pestrých, laterálne sa meniacich náplavov povrchového nivného krytu dnových akumulácií všetkých hlavných tokov. Nivy sú spestrené fluviálno-organickými sedimentmi vo výplniach mŕtvych ramien, prípadne organogénnymi sedimentmi – rašelinami slatinného typu – v iných lokálne podmáčaných depresiách (Hrubý Šúr, Pusté Úľany a i.).

Preskúmanosť študovaného územia

K poznaniu neogénnej stratigrafie významne prispel vrtný naftový prieskum, ktorý sa v tomto území začal v 40. rokoch minulého storočia v okolí Ivanky pri Dunaji. Stratigrafické začlenenie sedimentov na základe mikrofosílií (foraminifer, lastúrníček) urobil Pokorný (1946). Identifikoval bádenské, sarmatské a panónske sedimenty. Matějka (1949) v stručnosti opísal geologické a stratigrafické pomery Dunajskej panvy. V rokoch 1952 – 1954 bol vyhlásený prvý oporný vrt Báhoň-1 (Homola et al., 1955). Biostratigrafickým výskumom sedimentov uvedeného vrtu sa zaoberala Slavíková (1954). V 50. rokoch minulého storočia sa uskutočnilo mapovanie neogénnych sedimentov Podunajskej nížiny v rámci zostavovania generálnej mapy 1 : 200 000, list Bratislava (Buday, 1957). Mikrofaunu z povrchových lokalít, ako aj kopaných sond vyhodnotil Cicha (1957).

V 60. rokoch pokračoval geologický výskum v súvislosti s výstavbou vodného diela Dunaj, zameraný na objasnenie tektonickej stavby južnej oblasti slovenskej časti Dunajskej panvy (Janáček, 1966a, b, 1967).


V sedemdesiatych a osemdesiatych rokoch sa v rámci projektu výskumných geotermálnych vrtov realizovalo niekoľko hlbokých vrtov. Severozápadne od Galanty bol vyhlásený geotermálny vrt FGT-1 v Topoľníkoch (Franko, 1978), vrt FGGA-1 v Gabčíkove (Franko et al., 1984), vrt FGV-1 vo Vlčanoch (Remšík et al., 1984) a vrt GPB-1 v Boheľove (Fendek et al., 1984). V polovici 80. rokov minulého storočia sa na území centrálnej časti Bratislavy v rámci projektovanej rýchlodráhy realizoval vrtný prieskum (Vlasková et al., 1986). Z týchto vrtov boli spracované viaceré fosilné skupiny (mäkkýše, ostrakódy, palynomorfy, vápnitý nanoplanktón), čo prispelo k presnejšiemu rozčleneniu vrchnomiocénnych sedimentov okrajovej časti Dunajskej panvy (Fordinál, 1993a, b, 1995; Fordinál a Tuba, 1988, 1992; Fordinál et al., 1990, 1993; Nagy et al., 1995; Papšíková, 1989).

Začiatkom 21. storočia bola študovaná stratigrafia vrchnomiocénnych brakických a sladkovodných sedimentov slovenskej časti Dunajskej panvy (Kováč et al., 2010, 2011; Šujan et al., 2016).

Biostratigrafické vyhodnotenie vzoriek z regiónu Podunajská nížina-východ podáva Zlinská (1997). Paleoeologické trendy vrchnobádenských sedimentov Dunajskej panvy a Viedenskej panvy sú korelované v práci Zlinskej et al. (2002). Tektoniku, biostratigrafiu a paleoekológiu sarmatu a karpátu v severnej časti Dunajskej panvy riešia práce Hrušeckého et al. (2001), Zlinskej et al. (2005) a Fordinála et al. (2006a, b). Najnovšia syntéza biostratigrafie, sedimentológie a paleoprostredia Dunajskej panvy je zhrnutá v prácach Rybára et al. (2013, 2014, 2015), Kováča et al. (2015) a Hudáčkovej et al. (2015).

Študované oligocénne litostratigrafické jednotky Dunajskej panvy

Litostratigrafia paleogénnych sedimentov je prispôbená litostratigrafii severného Maďarska, tzv. budínskemu paleogénu (podľa Császára – ed., 1997, obr. 3), upravená v zmysle novších výskumov (Budai a Konrád, 2011). Najmladším členom je *séčenské súvrstvie* (*séčenský šlir*) veku eger – egenburg. Na označenie najmladších, kišcelských až spodnoegerských sedimentov sme použili termín *kišcelské súvrstvie* (v zmysle Tariho et al., 1993). Laterálnym ekvivalentom oboch spomenutých jednotiek sú *törökbálintské pieskovce* (*Törökbálint*). Pod kišcelským súvrstvom je usadené *tardské súvrstvie* (Báldi, 1984), vo vrchnej časti s laterálnym ekvivalentom *hárshégyského pieskovca* (*Hárshégvi*). Vek súvrstvia je po hranicu kišcel – priabón (~36 mil. rokov).



Obr. 3. Litostratigrafické jednotky terciéru Dunajskej panvy.
Fig. 3. Tertiary lithostratigraphic units of Danube Basin.

Kišcelské súvrstvie (kišcelské íly) – vrchný kišcel až spodný eger

Názov je zavedený v Maďarsku pre íly a slie ne rupel-ského veku s typovou lokalitou pri obci Kiscell. V oblasti Západných Karpát tento názov prvý raz použili Andrusov, Bystrická a Schaleková (1954) a Čílek (1954) pre oligocénne foraminiferové slie ne na jz. Slovensku.

Maďarský oligocén na základe foraminifer rozdelil Majzon (1960). Juhoslovenský oligocén s Majzonovými zónami koreloval Seneš (1964). Podľa výsledkov typová lokalita kišcelských ílov zodpovedá najspodnejšiemu horizontu oligocénnych morských slieňov (štvrtý Majzonov horizont), t. j. strednému „globigerínovo-kasidulinovému obzoru“. Tento „obzor“ v oblasti Dunajskej panvy zodpovedá spodnému oligocénu, spodným cyrenovým vrstvám, resp. pestrému uhľonosnému súvrstviu oligocénu, a v Maďarsku asi aj vrchným, už marinným „obzorom“ hárshegyských pieskovcov. Nie je vylúčená ich čiastočná ekvivalencia s vrchnou časťou budínskych slieňov (obr. 3). Vyššie horizonty kišcelských ílov tvoria rabdaminovo-cyklaminový piesčitý obzor (Majzonov obzor

č. 3), ktorý zodpovedá nášmu pieskovcovému obzoru, klastulínový a vrchný globigerínový obzor (Majzonov 1. a 2. obzor), ktorý zodpovedá našim foraminiferovým slieňom rupelu. K stratigraficky dôležitej mikrofaune patrí: *Tritaxia szabo* (HANTKEN), *Karrieriella hantkeniana* CUSHMAN, *Planularia kubinyi* (HANTKEN) atď.

Kišcelské súvrstvie predstavuje svetlosivý ílovitý a ílovito-slienitý aleurit a ílovitý slieňovec. Fácia je plytkobatyálna, s normálnou salinitou. Je všeobecne rozšírená na povrchu v Budínskych vrchoch a v predpolí Severného stredohoria. V severnej časti Dorogskej panvy sa prelína s mányským súvrstvom. Vo vrstvovom slede sa občas vyskytujú aj pieskovcové úseky, tie sú vyčlenené ako *budakesziský pieskovec*. Vyskytuje sa na povrchu v menšom množstve aj v okolí Recsku a v predpolí Bükku obsahuje aj mangánové ložiská. Hrúbka sa pohybuje obvykle v rozptí 30 – 500 m (max. 1 000 m).

Kišcelské súvrstvie sa vyskytuje v napr. v Budínskych vrchoch, Dorogskej panve, Cserháte, predpolí Bükku (Nagymarosy in Gyalog a Császár, 1996) a podľa našich terajších mikrofaunistických výskumov aj v železovskej priehlbine Dunajskej panvy.

Študované miocénne litostratigrafické jednotky Dunajskej panvy

Za základ litostratigrafie nám poslúžili litostratigrafické jednotky podľa Vassa (2002, obr. 3), ktorých stratigrafických rozsah a názvoslovie boli upravené v zmysle novších výskumov. Stupeň bádén sme ponechali ako trojdielny. Použitú časovú škálu Gradsteina a Ogga (2004) sme tiež upravili a prispôbili novším prácam, hlavne podľa Pillera et al. (2007) a Gradsteina et al. (2012).

Bajtavské súvrstvie, spodný bádén

Názov je odvodený od obce Bajtava sv. od Štúrova. Súvrstvie opísali ako bajtavskú sériu Seneš et al. (1971). Bazálnu a okrajovú časť súvrstvia tvoria transgresívne hrubé klastiká – zlepenca a andezitové vulkanoklastiká periférie stratovulkánu Börzöny vrátane andezitového epiklastického pieskovca, konglomerátu, brekcie, tufu s telesami riasového vápenca a amfistegínového pieskovca. Laterálne smerom do panvy tieto bazálne horniny prechádzajú do panvovej fácie tvorenej sivým vápnitým rozpadavým prachovcom a ílovcom s bridličnatým rozpadom. Z hľadiska objemu sedimenty panvovej fácie predstavujú hlavnú masu súvrstvia. Súvrstvie vzniklo v morskom prostredí. Do jeho pobrežnej zóny v oblasti Kováčovských kopcov (Burda) zasahovala periférna zóna vznikajúceho stratovulkánu Börzöny.

Typový profil je vrt ŠO-1 (hlbka 6,0 – 150 m) v. od obce Chľaba. Vrt a obzvlášť faunu a nanoflóru podrobne spracovali Brestenská, Lehotayová a Ondrejčíková (in Papp et al. – eds., 1978).

Severnejšie a severovýchodne od stratotypovej oblasti boli opísané *plášťovské vrstvy* (Vass, 1971), člen bajtavského súvrstvia. Hrúbka bajtavského súvrstvia je okolo 800 m. Vo fáciostrototypových vrtoch K-5 (Salka pri Štúrove) dosahuje súvrstvie hrúbku 867 m, vo vrte ŽI-2 (Lontov pri Želiezovciach) 389 m a vo vrte ŠO-1 (Chľaba pri Štúrove) 144 m (Fordinál et al., 2002). Bajtavské súvrstvie je rozšírené vo východnej časti Dunajskej panvy v komjatickej a železovskej priehlbine.

Spodnobádenský vek súvrstvia je doložený spoločenstvom foraminifer, v ktorom sú zastúpené *Lenticulina inornata* (ORB.), *Pappina bononiensis compressa* CUSH. a *Orbulina suturalis* BROENN., mäkkýšmi *Fusus* (APTYXIS) *lamellosus* BORSON, *Corbula* (VARICORBULA) *gibba* (OLIVI) a *Amussium cristatum badense* (FONT.) a vápnitým nanoplanktónom zóny NN-5 s indexovým druhmi *Sphenolithus heteromorphus* DEFL. (Lehotayová a Ondrejčíková, 1966; Lehotayová, Brestenská a Ondrejčíková in Papp et al., 1978).

Z novších výskumov spodnobádenské sedimenty bajtavského súvrstvia mikrofaunisticky potvrdila Zlinská (2004) vo vrte HGŽ-3 (Želiezovce) v hĺbke 770,5

– 854,6 m. Zastupujú ich bazálne zlepenca a hrubozrnné pieskovce s polohou ílov (836,0 – 841,0 m). Spoločenstvo foraminifer obsahovalo druhy: *Praeorbulina glomerosa* (BLOW), *Orbulina suturalis* BROENN., *Globigerinoides trilobus* (RSS.), *Pullenia bulloides* (ORB.), *Bulimina striata* ORB., *Uvigerina semiornata semiornata* ORB., *U. macrocarinata* PAPP et TURNOVSKY, *Cibicidoides pseudoungerianus* (CUSH.) a i. Foraminiferové asociácie patria k lagenidovej biozón (Grill, 1943).

K členom bajtavského súvrstvia patria: *turovské vrstvy* (spodný bádén), *plášťovské vrstvy* (spodný až stredný bádén), *šurianske vulkanity* (spodný bádén) a *vulkanity Burdy* (spodný bádén).

Pozbianske súvrstvie, stredný až vrchný bádén

Názov je odvodený od obce Pozba jz. od Levíc. Názov navrhol Vass (in Keith et al., 1989). Súvrstvie má na báze zlepenec, pieskovec a tuf. Smerom do panvy bazálne klastiká miznú. Panvový vývoj reprezentujú sivý vápnitý rozpadavý prachovec a ílovec. Bližšie k severnému okraju panvy v súvrství dominuje vápnitý, miestami piesčitý ílovec a prachovec (pri Zlatých Moravciach). V okolí Vrábľ a Mojmíroviec pieskovec a pelit sa striedajú v polohách hrubých 1 až 3 m (Gaža, 1963, 1965; fide Biela, 1978; Gaža a Beinhauerová, 1976, 1977). Súvrstvie vzniklo v morskom prostredí litorálnej zóny a na šelfe.

Typové profily sú vrt P-3 jv. od Pozby (od 950 do 1 020 m) a vrt Vr-1 sz. od mesta Vrábľa (od 2 300 do 2 460 m).

Súvrstvie leží zväčša skryto diskordantne na špačinskom súvrství, na severnom okraji panvy leží diskordantne na predneogénnom, zväčša mezozoickom podloží. Zakryté je vrábľským súvrstvom.

Maximálna hrúbka súvrstvia v komjatickej a železovskej priehlbine vo vrte ZM-1 pri Zlatých Moravciach je okolo 300 m (Gaža, 1970b).

Vek súvrstvia je stredný až vrchný bádén. Obsahuje rovnaké spoločenstvo morských foraminifer ako báhonské súvrstvie. Miestami dominuje spoločenstvo s brakickými formami (napr. vo vrte Vrábľa Vr-1; Jiříček in Gaža a Beinhauerová, 1976; Jiříček in Papp et al., 1978). Stratigrafický rozsah pozbianskeho súvrstvia v komjatickej depresii Hók et al. (1999) rozšírili aj na stredný bádén (obr. 3). Zdôvodňujú to tým, že strednobádenské a pozbianske sedimenty vo výplni depresie predstavujú jeden sedimentačný cyklus, a preto ich netreba členiť na dve súvrstvia, t. j. špačinské (stredný bádén) a pozbianske (vrchný bádén).

Strednobádenský vek súvrstvia je doložený foraminiferovými spoločenstvami zóny *Spiroplectamina carinata* (Grill, 1941). Mikrofaunisticky ho zistila Zlinská (2004) vo vrte HGŽ-3 v hĺbkovom intervale 550,2 – 770,2 m. Sedimenty tu tvoria sivé íly s premenlivým obsahom piesčitej a prachovej prímеси, pieskovce s polohami tu-

fitov a piesčitých tuftických ílov. Asociácie foraminifer obsahovali druhy: *Globigerinoides trilobus* (RSS.), *G. quadrilobatus* (ORB.), *Pullenia bulloides* (ORB.), *Globigerinella obesa* (BOLLI), *Spiroplectinella carinata* (ORB.), *S. acuta* (RSS.), *Uvigerina semiornata semiornata* ORB., *Valvulineria complanata* (ORB.), *Guttulina communis* ORB., *Cibicoides ungerianus* (ORB.), *Hansenisca soldanii* (ORB.), *Fursenkoina acuta* (ORB.) atď. (Zlinská, l. c.).

Sedimenty vrchného bádenu biostratigraficky preukázala Zlinská (2004) vo vrte HGŽ-3 v hĺbke 250,4–550,2 m. Predstavujú ich sivé íly s piesčitou a prachovou prímiesou, prachy a drobno- až strednozrnné piesky. Vyznačujú sa foraminiferami *bulimínovo-bolivínovej* zóny (Grill, 1941). Prítomné sú napr. druhy: *Bolivina dilatata dilatata* RSS., *B. hebes* MACF., *Bulimina elongata elongata* ORB., *Bulimina elongata longa* (VENGL.), *Globigerina bulloides* (ORB.), *G. druryi* AKERS, *Globigerinella obesa* (BOLLI), *Globigerinoides trilobus* (RSS.), *Lagena acuticosta* RSS., *Asterigerinata planorbis* (ORB.) atď.

V odkryve na ssz. okraji obce Malá nad Hronom mikrofaunu pozbianskeho súvrstvia určila Zlinská (1993).

Vrábeľské súvrstvie, sarmat

Názov je odvodený od mesta Vráble v komjatickej priehlbine Dunajskej panvy. Zaviedol ho Vass (1989). Súvrstvie má pestré litologické zloženie. V železovskej priehlbine sa na jeho stavbe podieľa zlepenec (hlavne na báze), piesčité, oolitické a lumachelové vápenec, pieskovec/piesok a kyslý biotitický tuf. Smerom do centra priehlbiny narastá podiel sivého vápnitého, miestami tuftického ílu. Vrchnú časť súvrstvia tvorí piesok s polohami ílu (Seneš in Seneš et al., 1962). V komjatickej depresii dominujú íl a piesok, na sv. a v. okraji sú uhoľné vrstvy (Gaža, 1968, 1970b; Gaža a Beinhauerová, 1976). V blatnianskej priehlbine dominujú vápnitý prach a íl, na západnom okraji priehlbiny s polohami štrku a organogénneho vápenca. Vrchnú časť súvrstvia tvorí piesok, pestrý a sivý vápnitý íl, uhoľný íl so šošovkami a slojčkami uhlia (Homola, 1955; Gaža, 1968, 1970b). Súvrstvie vznikalo v mori s nízkou salinitou, v jeho litorálnej zóne a na šelfe. Uhoľné vývoje svedčia najskôr o prostredí progradujúcej delty (deltová plošina; Vass in Keith et al., 1989).

Typový profil je vrt Vr-1 vyhlbený sz. od mesta Vráble (hlbkový interval 1 165 – 1 780 m). Súvrstvie leží so skrytou diskordanciou na pozbianskom, resp. na báhonskom súvrství (vrchný bádén) a je zakryté ivanským súvrstvom (panón). Hrúbka súvrstvia je 300 (železovská priehlbina), 500, resp. 700 m (blatnianska a komjatická priehlbina) a v rišňovskej priehlbine až 1 600 m. Súvrstvie je rozšírené v celej Dunajskej panve.

Sarmatský vek súvrstvia je doložený hlavne foraminiferovými spoločenstvami zóny veľkých elfidií, zóny s *Elphidium hauerinum* a zóny *Protelphidium subgranosum* (Grill, 1941). Mikrospoločenstvá z neho opísali Jiříček

(1972), Brestenská in Papp et al. (1974) a z novších výskumov Zlinská (2004) vo vrte HGŽ-3 (Železovce) v hĺbke 10 – 250 m. Faunu mäkkýšov opísali Seneš (1953), Seneš in Seneš et al. (1962) a Švagrovský (1971).

Vo vrte HGŽ-3 sarmatské sedimenty tvoria hlavne sivé až sivozelené piesčité íly s ojedinelými polohami ílov. Spodnosarmatskú mikroasociáciu s prevahou elfidií (zóna *Elphidium reginum*, Grill 1941), no bez typickej formy, zistila Zlinská (l. c.) v hĺbke 150,4 – 155,5 m. Prítomné sú napr. *Elphidium flexuosum flexuosum* (ORB.), *E. rugosum* (ORB.), *Bolivina sarmatica* DIDK., *Cibicides badenensis* (ORB.), *Lobatula lobatula* (W.-J.) a *Ammonia beccarii* (L.). Rod *Elphidium* sa vo všeobecnosti považuje za ukazovateľa malej hĺbky vody. Vo väčšom množstve sa nachádza len v hĺbke menšej než 80 – 100 m (Boltovskoy a Wright, 1976; Walton, 1964; Poag, 1981). Foraminifery sprevádzajú centrické, prevažne pyritizované rozsievky.

Z ostrakód bol prítomný druh *Cytheridea hungarica* ZÁLANYI, ktorý poukazuje na spodnosarmatskú biozónu *Cytheridea hungarica* – *Aurila mehesi* (Zlinská, l. c.).

Ekvivalentom vrábeľského súvrstvia vo Viedenskej panve sú holičské a skalické súvrstvie, vo Východoslovenskej panve stretavské a ptrukšianske súvrstvie.

K členom vrábeľského súvrstvia patria *vrstvy od Malej* (spodný sarmat).

Metodika

Po odbere vzorkového materiálu zo skladov Nafta, a. s., sme urobili jeho selekciu. Vo vrtoch sme sa zamerali hlavne na hraničné úseky jednotlivých stupňov. Vzorky po rozdrvení sa macerovali. Foraminifery sme získali z výplavu bežným laboratórnym postupom, plavením cez mlynársky hodváb a separáciou výplavu. V prípade negatívnych vzoriek sme odber zahustili a laboratórny postup zopakovali. Získaný materiál sme študovali pod binokulárnou lupou. Niektoré spoločenstvá boli pod ňou aj fotograficky zdokumentované a sú vyobrazené pri mikrofaunistickom vyhodnotení jednotlivých vrtov a vo fototabulkách. Prvé číslo za hlbkovým intervalom vrtu je číslo jadrovnice, druhé je číslo vzorkovnice. Na určenie foraminifer a stratigrafického rozšírenia taxónov bola použitá literatúra zameraná hlavne na centrálnu Paratetýdu (napr. Cicha et al., 1998). Profily vrtov sú prevzaté z práce Bielej (1978). Časová škála predstavuje modifikáciu prác Gradsteina a Ogga (2004), Pillera et al. (2007) a Gradsteina et al. (2012).

Výsledky


Modrany 1 (M-1; obr. 1, 4)

Na základe 106 taxónov foraminifer (tab. 1, fototab. 1 – 3, obr. 5) a ich zastúpenia sme vo vrte Modrany 1 determinovali nasledujúce litostratigrafické jednotky (obr. 3):



TAB. I. Spoločenstvo foraminifer z vrtu M-1: 1 – 5 – z hĺbky 1 098 – 1 103 m; 6 – 7 – z hĺbky 1 298 – 1 303 m; 8 – z hĺbky 1 990 až 1 995 m.

TAB. I. Intercommunity of foraminifera from borehole M-1: 1 – 5 – in depth 1 098 – 1 103 m ; 6 – 7 – in depth 1 298 – 1 303 m; 8 – in depth 1 990 – 1 995 m.



TAB. II. 1 – Spoločenstvo foraminifer z vrtu NV-1: 1 – z hĺbky 3 024 – 3 026,5 m; 2 – z hĺbky 3 057 – 3 062 m; 3 – 4 – *Elphidium josephinum* (ORB.), M-1 / 1 050 – 1 056 m; 5 – *Schackoinella imperatoria* (ORB.), M-1 / 1 050 – 1 056 m; 6 – *Orbulina suturalis* BROENN., NV-1 / 2 431 m; 7 – *Lenticulina* sp., NV-1 / 2 556 m; 8 – *Subbotina angiporoides* (HORNIBROOK), NV-1 / 3 024 m; 9 – *Globigerinoides primordius* BLOW et BANNER, NV-1 / 3 024 m; 10 – *Tenuitella munda* (JENKINS), NV-1 / 3 075 m; 11 – *Globigerina officinalis* SUBB., NV-1 / 3 075 m.

TAB. II. 1 – Intercommunity of foraminifera from borehole NV-1: 1 – in depth 3 024 – 3 026,5 m; 2 – in depth 3 057 – 3 062 m; 3 – 4 – *Elphidium josephinum* (ORB.), M-1 / 1 050 – 1 056 m; 5 – *Schackoinella imperatoria* (ORB.), M-1 / 1 050 – 1 056 m; 6 – *Orbulina suturalis* BROENN., NV-1 / 2 431 m; 7 – *Lenticulina* sp., NV-1 / 2 556 m; 8 – *Subbotina angiporoides* (HORNIBROOK), NV-1 / 3 024 m; 9 – *Globigerinoides primordius* BLOW et BANNER, NV-1 / 3 024 m; 10 – *Tenuitella munda* (JENKINS), NV-1 / 3 075 m, 11 – *Globigerina officinalis* SUBB., NV-1 / 3 075 m.



TAB. III. Jadrá z vrtu Modrany 1: 1 – z hlĺbky 1 050 – 1 056 m; 2 – z hlĺbky 1 402 – 1 406 m; 3 – z hlĺbky 1 698 – 1 702 m; 4 – z hlĺbky 1 755 – 1 760 m; 5 – z hlĺbky 1 990 – 1 995 m; 6 – 7 – z hlĺbky 2 102 – 2 105 m.

TAB. III. Centre cores from the well Modrany 1: 1 – from depth 1 050 – 1 056 m; 2 – from depth 1 402 – 1 406 m; 3 – from depth 1 698 – 1 702 m; 4 – from depth 1 755 – 1 760 m; 5 – from depth 1 990 – 1 995 m; 6 – 7 – from depth 2 102 – 2 105 m.



TAB. IV. Jadrá z vrtu Nová Vieska 1: 1 – z hĺbky 1 797 – 1 799 m; 2 – z hĺbky 2 556 – 2 562 m; 3 – z hĺbky 2 875 – 2 879 m; 4 – z hĺbky 3 024 – 3 026,5 m; 5 – z hĺbky 3 057 – 3 062 m; 6 – z hĺbky 3 089 – 3 093 m; 7 – z hĺbky 3 164 – 3 171 m.

TAB. IV. Centre cores from the well Nová Vieska 1: 1 – from depth 1 797 – 1 799 m; 2 – from depth 2 556 – 2 562 m; 3 – from depth 2 875 – 2 879 m; 4 – from depth 3 024 – 3 026,5 m; 5 – from depth 3 057 – 3 062 m; 6 – from depth 3 089 – 3 093 m; 7 – from depth 3 164 – 3 171 m.

Tab. 1

Prehľadná tabuľka zastúpenia foraminifer v jednotlivých intervaloch vrtu Modrany 1 (M-1).
Summary table of foraminifera representation in particular satin yardage of borehole Modrany 1 (M-1).

Taxon	1 050 – 1 056 m (12/1)	1 050 – 1 056 m (12/5)	1 098 – 1 103 m (13/1)	1 298 – 1 303 m (17/1)	1 402 – 1 406 m (19/1)	1 698 – 1 702 m (25/2)	1 755 – 1 760 m (26/2)	1 855 – 1 859 m (28/3)	1 990 – 1 995 m (31/3)
<i>Ammodiscus miocenicus</i> KARRER									x
<i>Ammonia viennensis</i> (ORB.)		x	x		x		x		
<i>Amphimorphina haueriana</i> NEUGEB.			x						
<i>Amphistegina mamilla</i> (F. et M.)			x						
<i>Angulogerina angulosa</i> (WILLIAMSON)								x	
<i>Anomalinoides badenensis</i> (ORB.)			x						
<i>Asterigerinata planorbis</i> (ORB.)			x	x	x				
<i>Asterigerinoides</i> sp.						x			
<i>Bathysiphon taurinensis</i> SACCO				x					x
<i>Bolivina dilatata</i> RSS.		x				x			
<i>Bolivina hebes</i> MACFAD.			x	x					
<i>Bolivina miocenica</i> MACFAD.					x				
<i>Bolivina moldavica granensis</i> C. et Z.	x	x							
<i>Bolivina pseudoplicata</i> HERON-ALLEN et EARLAND	x	x							
<i>Bulimina aculeata</i> ORB.			x						
<i>Bulimina</i> aff. <i>subulata</i> CUSH. et PARKER						x			
<i>Bulimina elegans</i> (ORB.)		x							
<i>Bulimina schischkinskayae</i> SAMOILOVA				x		x			
<i>Bulimina striata</i> ORB.			x						
<i>Catapsydrax</i> sp.									x
<i>Cibicidoides ungerianus</i> (ORB.)			x						
<i>Cyclammina karpatica</i> C. et Z.				x					
<i>Cyclammina</i> sp.				x					
<i>Cyclammina vulchoviensis</i> VENGL.			x						
<i>Cycloforina badenensis</i> (ORB.)			x						
<i>Cycloforina contorta</i> (ORB.)			x	x					
<i>Elphidium aculeatum</i> (ORB.)		x							
<i>Elphidium crispum</i> (L.)		x							
<i>Elphidium fichtellianum</i> (ORB.)		x	x		x				
<i>Elphidium hauerinum</i> (ORB.)	x	x							
<i>Elphidium josephinum</i> (ORB.)	x	x							
<i>Elphidium macellum</i> (F. et M.)			x						
<i>Elphidium reginum</i> (ORB.)	x								
<i>Elphidium rugosum</i> (ORB.)			x						
<i>Elphidium samueli</i> ZLINSKÁ		x							
<i>Elphidium</i> sp.							x		
<i>Eponides pygmeus</i> (HANTK.)									x
<i>Fursenkoina acuta</i> (ORB.)			x						
<i>Globigerina</i> aff. <i>postcretacea</i> MJATLJUK								x	
<i>Globigerina bulloides</i> ORB.			x	x	x				
<i>Globigerina concinna</i> RSS.			x						
<i>Globigerina diplostoma</i> RSS.			x						
<i>Globigerina officinalis</i> SUBB.								x	x
<i>Globigerina ouachitaensis</i> HOWE et WALLACE									x
<i>Globigerina praebulloides</i> BLOW			x		x				
<i>Globigerina praebulloides leroyi</i> BLOW								x	
<i>Globigerinella obesa</i> (BOLLI)			x	x					
<i>Globigerinoides quadrilobatus</i> (ORB.)			x	x					
<i>Globigerinoides trilobus</i> (RSS.)	x		x	x	x				
<i>Globoquadrina langhiana</i> CITA et GELATI				x					
<i>Globoturborotalita druryi</i> (AKERS)			x						
<i>Globoturborotalita woodi</i> (JENKINS)					x				

Pokračovanie tab. 1.

Taxon	1 050 – 1 056 m (12/1)	1 050 – 1 056 m (12/5)	1 098 – 1 103 m (13/1)	1 298 – 1 303 m (17/1)	1 402 – 1 406 m (19/1)	1 698 – 1 702 m (25/2)	1 755 – 1 760 m (26/2)	1 855 – 1 859 m (28/3)	1 990 – 1 995 m (31/3)
<i>Globulina gibba</i> ORB.						x			
<i>Guttulina communis</i> (ORB.)			x						
<i>Hansenisca soldanii</i> (ORB.)			x		x				
<i>Hanzawaia boueana</i> (ORB.)			x	x	x	x			
<i>Heterolepa dutemplei</i> (ORB.)			x		x				
<i>Chiloguembelina gracillima</i> (ANDREAE)								x	
<i>Chilostomella czizeki</i> RSS.									x
<i>Lenticulina cultrata</i> (MONT.)						x			
<i>Lenticulina inornata</i> (ORB.)				x		x			
<i>Lenticulina</i> sp.									x
<i>Lenticulina</i> sp. [cf. <i>L. melvilli</i> (CUSH. et RENZ)]					x				
<i>Lenticulina vortex</i> (F. et M.)				x					
<i>Lobatula lobatula</i> (W. et J.)		x	x		x			x	x
<i>Martinottiella communis</i> (ORB.)			x	x					
<i>Melonis pompilioides</i> (F. et M.)			x	x	x			x	
Miliolidné foraminifery	x					x			
<i>Nodosaria</i> aff. <i>acuminata</i> HANTKEN									x
<i>Nodosaria</i> sp.			x						
<i>Nonion bogdanowiczi</i> VOLOSH.	x	x							
<i>Nonion commune</i> (ORB.)			x		x	x	x	x	
<i>Nonion serenus</i> VENGL.	x								
<i>Orbulina suturalis</i> BROENNIMANN			x	x					
<i>Paragloborotalia?</i> <i>mayeri</i> (CUSH et ELL.)					x				
<i>Plectofrondicularia digitalis</i> (NEUGEB.)			x						
<i>Porosonion granosum</i> (ORB.)	x								
<i>Praeglobobulimina pupoides</i> (ORB.)			x	x					
<i>Praeglobobulimina pyrula</i> (ORB.)			x						
<i>Praeorbulina glomerosa</i> (BLOW)				x					
<i>Psammosiphonella cylindrica</i> (GLAESSNER)									x
<i>Pullenia bulloides</i> (ORB.)					x				
<i>Pygmaeosestron hispidum</i> RSS.			x						
<i>Quinqueloculina</i> sp.		x							
<i>Reophax nodulosus brevior</i> LOMNICKI				x					
<i>Reticulophragmium amplexans</i> (GRZYB.)									x
<i>Reussella spinulosa</i> (RSS.)					x				
<i>Semivulvulina deperdita</i> (ORB.)			x						
<i>Sphaeroidina bulloides</i> ORB.			x			x			
<i>Spiroplectinella carinata</i> (ORB.)			x	x					x
<i>Stilostomella advena</i> (CUSH. et LAIM.)			x						
<i>Subbotina linaperta</i> (FINLAY)									x
<i>Subbotina pseudovenezuelana</i> (BLOW et BANNER)									x
<i>Subbotina tapuriensis</i> BLOW et BANNER								x	
<i>Tenuitella danvilensis</i> (HOWE et WALLACE)								x	
<i>Tenuitella munda</i> (JENKINS)								x	
<i>Tenuitellinata angustumbilicata</i> (BOLLI)						x		x	
<i>Textularia gramen maxima</i> C. et Z.			x						
<i>Textularia mariae</i> ORB.			x						
<i>Textularia pala</i> CZJZ.						x			
<i>Triloculina</i> sp.		x							
<i>Turborotalia cerroazulensis</i> (COLE)									x
<i>Uvigerina aculeata</i> ORB.			x	x					
<i>Uvigerina pygmoides</i> P. et T.					x				
<i>Uvigerina semiornata</i> ORB.			x	x					
<i>Valvulineria complanata</i> (ORB.)			x	x					


Tab. 2

Prehľadná tabuľka zastúpenia foraminifer v jednotlivých intervaloch vrtu Nová Vieska 1 (NV-1).
Summary table of foraminifera representation in particular satin yardage of borehole Nová Vieska 1 (NV-1).

Taxón	1 797 – 1 799 m	2 431 – 2 435 m	2 556 – 2 562 m	2 875 – 2 879 m	3 024 – 3 026,5 m	3 057 – 3 062 m	3 075 – 3 078 m	3 089 – 3 093 m	3 103 – 3 108 m	3 156 – 3 161 m
<i>Almaena</i> sp.										x
<i>Ammodiscus glabratus</i> (ORB.)					x					
<i>Ammodiscus miocenicus</i> KARRER					x					
<i>Ammodiscus</i> sp.								x		
<i>Ammonia viennensis</i> (ORB.)		x	x							
<i>Angulogerina esuriensis</i> HORNIBROOK		x								
<i>Angulogerina muralis</i> (TERQUEM)								x		
<i>Asterigerinata planorbis</i> (ORB.)	x									
<i>Bathysiphon</i> sp.							x			
<i>Bathysiphon taurinense</i> SACCO									x	
<i>Bolivina</i> aff. <i>pokorny</i> C. et Z.		x								
<i>Bolivina</i> aff. <i>reticulata</i> HANTKEN									x	
<i>Bolivina</i> aff. <i>semistriata</i> HANTK.					x					
<i>Bolivina dilatata</i> Rss.		x								
<i>Bolivina scalprata muscosa</i> C. et Z.	x									
<i>Bolivina</i> sp.			x		x					
<i>Bolivina</i> sp. (cf. <i>B. dilatata</i> Rss.)	x									
<i>Bulimina coprolithoides</i> ANDREAE								x		
<i>Bulimina</i> sp.									x	
<i>Cibicidoides ornatus</i> (C. et Z.)		x								
<i>Cibicidoides pseudoungerianus</i> (CUSH.)								x	x	
<i>Cibicidoides ungerianus</i> (ORB.)							x			
<i>Elphidium</i> sp.	x									
<i>Epistomina</i> cf. <i>elegans</i> (ORB.)								x		
<i>Eponides pygmeus</i> (HANTK.)					x	x				
<i>Globigerina</i> aff. <i>ottnangiensis</i> ROEGL	x									
<i>Globigerina</i> aff. <i>anguliofficialis</i> BLOW							x			
<i>Globigerina officinalis</i> SUBB.						x	x	x		
<i>Globigerina ouachitaensis</i> HOWE et WALLACE								x		
<i>Globigerina praebulloidis</i> BLOW			x							
<i>Globigerina</i> sp.	x			x						
<i>Globigerinoides quadrilobatus</i> (ORB.)		x								
<i>Globigerinoides primordius</i> BLOW et BANNER					x					
<i>Globigerinoides trilobus</i> (Rss.)		x								
<i>Globorotalia bykovae</i> (AISENSTAT)		x								
<i>Globoturborotalita woodi</i> (JENKINS)	x									
<i>Hansenisca girardana</i> (Rss.)					x					
<i>Hansenisca soldanii</i> (ORB.)		x								
<i>Hanzawaia boueana</i> (ORB.)		x	x							

Pokračovanie tab. 2.

Taxón	1 797 – 1 799 m	2 431 – 2 435 m	2 556 – 2 562 m	2 875 – 2 879 m	3 024 – 3 026,5 m	3 057 – 3 062 m	3 075 – 3 078 m	3 089 – 3 093 m	3 103 – 3 108 m	3 156 – 3 161 m
<i>Heterolepa dutemplei</i> (ORB.)		x	x							
<i>Chiloguembelina gracillima</i> (ANDREAE)				x						
<i>Chilostomella ovoidea</i> RSS.		x								
<i>Lenticulina cultrata</i> (MONTFORT)		x								
<i>Lenticulina</i> sp.		x					x			
<i>Lobatula lobatula</i> (W. et J.)				x				x		
<i>Melonis pompilioides</i> (F. et M.)		x								
<i>Nonion commune</i> (ORB.)		x	x		x					
<i>Nothia excelsa</i> (GRZYB.)					x					
<i>Nothia latissima</i> (GRZYB.)						x	x			
<i>Orbulina suturalis</i> BROENN.		x								
<i>Paragloborotalia opima nana</i> (BOLLI)							x			
<i>Paragloborotalia</i> sp.					x					
<i>Praeorbulina sicana</i> (DE STEFANI)	x	x								
<i>Praeorbulina glomerata</i> (BLOW)		x								
<i>Pullenia bulloides</i> (ORB.)		x								
<i>Pullenia quinquelobata</i> (RSS.)					x					
<i>Recurviroides amplexans</i> (GRZYB.)						x				
<i>Repmanina serpens</i> (GRZYB.)							x			
<i>Rhabdammina cylindrica</i> GLAESSNER									x	
<i>Rhabdammina</i> sp.						x				
<i>Sigmoilopsis celata</i> (COSTA)		x								
<i>Subbotina</i> aff. <i>cryptomphala</i> (GLAESSNER)						x			x	
<i>Subbotina</i> aff. <i>galavisi</i> (BERMUDEZ)					x					
<i>Subbotina angiporoides</i> (HORNIBROOK)					x			x		
<i>Subbotina cryptomphala</i> (GLAESSNER)							x			
<i>Subbotina linaperta</i> (FINLAY)						x		x	x	
<i>Subbotina praeturritilina</i> (BLOW et BANNER)							x			
<i>Subbotina pseudoeocaena</i> (SUBB.)					x		x			
<i>Subbotina pseudovenezuelana</i> (BLOW et BANNER)					x		x			
<i>Subbotina</i> sp.										x
<i>Subbotina tapuriensis</i> BLOW et BANNER						x			x	
<i>Subbotina triangularis</i> (WHITE)					x					
<i>Tenuitella munda</i> (JENKINS)							x			
<i>Textularia gramen deltoidea</i> ORB.		x								
<i>Textularia gramen subangulata</i> ORB.		x								
<i>Textularia laevigata</i> ORB.		x								
<i>Trifarina bradyi</i> CUSH.			x							
<i>Tritaxilina pupa</i> (GUEMBEL)									x	
<i>Turborotalia cerroazulensis</i> (COLE)						x	x		x	
<i>Turborotalia</i> sp.										x
<i>Uvigerina acuminata</i> HOSIUS		x								
<i>Uvigerina multistriata</i> HANTKEN						x				
<i>Uvigerina pygmoides</i> P. et T.		x								
<i>Uvigerina</i> sp.			x							
<i>Valvulinaria complanata</i> (ORB.)		x								



Obr. 4. Litologický profil vrtov Modrany 1 a Nová Vieska 1 s odberom vzoriek a výsledkami revízie.

Fig. 4. Lithological profile of Modrany 1 and Nová Vieska 1 wells with collection of specimens and the revision results.

- **vrábeľské súvrstvie** sarmatského veku v hĺbke 1 050 – 1 056 m. Zastupuje ho typická elfidiovo-miliolidovo-nóniová asociácia foraminifer s druhmi *Elphidium josephinum* (ORB.), *E. reginum* (ORB.), *E. samueli* ZLINSKÁ, *Nonion bogdanowiczi* VOLOSH., *N. seren* V ENGL., *Bolivina moldavica granensis* C. et Z. atď. Mikrofaunisticky v zmysle Grilla (1941) zodpovedá biozóna *Elphidium reginum*.
- **pozbianske súvrstvie** (v zmysle Hóka et al., 1999) strednobádenského veku v hĺbke 1 098 – 1 303 m. V hĺbke 1 098 – 1 103 m je v ňom pomerne veľké zastúpenie aglutinovaných foraminifer rovnomennej zóny. Prítomné sú aj typické strednobádenské taxóny ako *Textularia gramen maxima* C. et Z., *Textularia mariae* ORB., *Semivulvulina deperdita* (ORB.), *Plectofrondicularia digitalis* (NEUGEB.) a planktonická forma *Globoturborotalita druryi* (AKERS). V hĺbke 1 298 – 1 303 m má výrazné zastúpenie *Orbulina suturalis* BROENN., ktorá sa v panvách centrálnej Paratetýdy objavuje vo vrchnej časti vrchnej lagenidovej zóny. Pomer aglutinovaných foriem s užším diapazónom však poklesol (obr. 5), vyskytuje sa tu len strednobádenský druh *Reophax nodulosus brevior* LOMNICKI a z vápnitého bentosu *Uvigerina aculeata* ORB.
- **bajtavské súvrstvie** v hĺbke 1 402 – 1 760 m. Charakterizuje ho spodnobádenská mikrofauna lagenidovej zóny, hlavne zástupcovia čeľade Vaginulinidae (rod *Lenticulina*). Z planktonickej zložky sú smerodajné *Globoturborotalita woodi* (JENKINS) a *Tenuitellinata angustumbilicata* (BOLLI).
- **kišcelské súvrstvie** oligocénneho až spodnoeager-ského veku v hĺbke 1 855 – 1 995 m. Charakterizuje ho veľmi drobné spoločenstvo planktonických foraminifer, ktoré objemovo tvoria asi 99 % v asociácii foraminifer. Prítomné sú napr. *Chiloguembelina gracillima* (ANDREAE), *Tenuitella munda* (JENKINS), *Globigerina* aff. *postcretacea* MJATLJUK, *Tenuitella danvilensis* (HOWE et WALLACE) a *Tenuitellinata angustumbilicata* (BOLLI). Vo vzorke z hĺbky 1 990 – 1 995 m narastá počet foriem prechádzajúcich z eocénu [napr. *Turborotalia cerroazulensis* (COLE), *Subbotina pseudovenezuelana* (BLOW et BANNER), *Subbotina linaperta* (FINALY) a *Reticulophragmium amplexans* (GRZYB.)]. V hĺbke 2 102 – 2 105 m boli len úlomky schránok mäkkýšov.

Nová Vieska 1 (NV-1; obr. 1, 4)

Na základe 86 taxónov foraminifer (tab. 2, fototab. 2 a 4, obr. 6) sme vo vrte Nová Vieska 1 determinovali nasledujúce litostratigrafické jednotky (obr. 3):


- **bajtavské súvrstvie** spodnobádenského veku v hĺbke 1 797 – 2 562 m. Charakterizuje ho mikrofauna lagenidovej zóny (rod *Lenticulina*) a indexové planktonické formy *Globigerinoides sicanus* DE STEFANI, *Praeorbulina glomerata* (BLOW) a *Orbulina suturalis* BROENN.
- **kišcelské súvrstvie** oligocénu až spodného egeru v hĺbke 2 875 – 3 108 m. V asociácii foraminifer má kvantitatívnu prevahu drobný planktón s charakteristickými druhmi ako: *Chiloguembelina gracillima* (ANDREAE), *Globigerinoides primordius*

Tab. 3

Prehľadná tabuľka litostratigrafického zaradenia vzoriek z jednotlivých vrtov.
Summary table of lithostratigraphical classification of samples from individual wells.


Modrany 1	Nová Vieska 1
1 050 – 1 056 m 12/1	1 797 – 1 799 m
1 050 – 1 056 m 12/5	2 431 – 2 435 m
1 098 – 1 103 m	2 556 – 2 562 m
1 298 – 1 303 m	2 800 – 2 805 m 85/2
1 402 – 1 406 m	2 875 – 2 879 m
1 698 – 1 702 m	3 024 – 3 026,5 m
1 755 – 1 760 m	3 057 – 3 062 m
1 855 – 1 859 m	3 075 – 3 078 m
1 990 – 1 995 m	3 089 – 3 093 m
2 102 – 2 105 m 33/3	3 103 – 3 108 m
2 102 – 2 105 m 33/4	3 156 – 3 161 m

vrábeľské súvrstvie
pozbianske súvrstvie
bajtavské súvrstvie
kišcelské súvrstvie
bez identifikácie



Obr. 5. Graf pomerného zastúpenia jednotlivých typov foraminifer vo vrte Modrany 1 (M-1): a – aglutinované, vb – vápnný bentos, p – planktón.

Fig. 5. Graph of proportional representation of different types of foraminifera in the well Modrany 1 (M-1): a – agglutinated, vb – calcareous benthos, p – plankton.



Obr. 6. Graf pomerného zastúpenia jednotlivých typov foraminifer vo vrte Nová Vieska 1 (NV-1): a – aglutinované, vb – vápnný bentos, p – planktón.

Fig. 6. Graph of proportional representation of different types of foraminifera in the well Nová Vieska 1 (NV-1): a – agglutinated, vb – calcareous benthos, p – plankton.

BLOW et BANNER, *Globigerina* aff. *anguliofficialis* BLOW, *Tenuitella munda* (JENKINS), *Globigerina ouachitaensis* HOWE et WALLACE atd'.

V hĺbke 2 800 – 2 805 m sa náhle zhoršili podmienky sedimentácie, čo sa odrazilo na absencii mikrofauny (obr. 6).

V hĺbke 2 875 – 3 026,5 m sme našli indexové fosílie *Chiloguembelina gracillima* (ANDREAE) s výskytom od spodného kišcelu až do spodnej časti vrchného kišcelu a *Globigerinoides primordius* BLOW et BANNER objavujúci sa vo vrchnom rupeli, resp. vrchnom oligocéne (chat, P22 – N4).

V hĺbke 3 057 – 3 093 m pribúda viac eocénnych taxónov, mnohé majú LAD v spodnom kišceli, ako napr.

Subbotina cryptomphala (GLAESSNER), *S. angiporoides* (HORNIBROOK), *S. pseudoeocaena* (SUBB.) a *Turborotalia cerroazulensis* (COLE), alebo sa priebežne vyskytujú od priabónu do egeru až egenburgu [*Subbotina praeturritilina* (BLOW et BANNER), *Globigerina officinalis* SUBB., *Paragloborotalia opima nana* (BOLLI)], ale zistili sme aj formy s FAD v kišceli, napr. *Tenuitella munda* (JENKINS) a *Globigerina* aff. *anguliofficialis* BLOW. Preto tieto sedimenty zaradujeme ešte k oligocénu.

V hĺbke 3 103 – 3 108 m sa nevyskytuje taxón s FAD v oligocéne, aj keď LAD niektorých sa končí v spodnom kišceli [*Subbotina tapuriensis* BLOW et BANNER a *Subbotina* aff. *cryptomphala* (GLAESSNER)].

V hĺbke 3 156 – 3 161 m sú rekryštalizované formy. Rod *Almaena* je známy z eocénu až oligocénu. Litologicky však ide pravdepodobne o hárshégyský pieskovec (obr. 3).

Revízia starších biostratigrafických údajov

Modrany 1

Na rozdiel od profilu Bielej (1978), ktorá z hĺbky 1 098 – 1 303 m uvádza sedimenty spodnobádenského veku, sme tento interval zaradili do stredného bádenu, zóny aglutinancií, s typickými strednobádenskými taxónmi. Túto zónu v hĺbke 1 098 – 1 103 m potvrdili aj pôvodné mikrofaunistické určenia Holzknechta a Jandovej v správe Gažu (1970a).

Spodný bádén (*bajtavské súvrstvie*) sme zistili až v najbližšej vzorke z hĺbky od 1 402 do 1 760 m, ale to už v hĺbke 1 670 m Biela (l. c.) udáva rupel (v dnešnom ponímaní kišcel). Ten sme ako *kišcelské súvrstvie* mikrofaunisticky preukázali hlbšie, až v intervale 1 855 – 1 995 m, kde Biela uvádza latorf a priabón (obr. 4).

V pôvodných mikrofaunistických určeníach (Holzknecht a Jandová) v správe Gažu (1970a) ani v geologickom opise vrtu sa o prítomnosti oligocénu vôbec neuvažuje. Pod sedimentmi spodného bádenu v hĺbke 1 665 – 1 786 m uvádza Gaža (l. c.) karpatskú formáciu a pod ňou vrchný eocén. Karpat ani typický vrchný eocén sme mikrofaunisticky nepotvrdili.

Nová Vieska 1

Na rozdiel od Bielej (1978), ktorá v hĺbke 2 533 – 2 746 m udáva sedimenty rupelského veku, sme našli spodný bádén (*bajtavské súvrstvie*) do hĺbky 2 746 m a namiesto latorfu (2 746 – 2 904 m) a priabónu (2 904 – 3 171 m) v hĺbke 2 750 – 3 171 m oligocén (obr. 4). Spodný bádén uvádzaný v Homolovej práci (1960) do hĺbky 2 533 m sme rozšírili až do hĺbky zhruba 2 750 m, teda pôvodný akvitán? v hĺbke 2 533 – 2 746 m po novom zodpovedá spodnému bádenu a vrchnoeocénny úsek (2 746 – 3 171 m) oligocénu (*kišcelské súvrstvie* a *hárshégyský pieskovec*; obr. 3).

Záver

Z dvoch vybraných hlbokých naftových vrtov (Modrany 1 a Nová Vieska 1; obr. 1) v Dunajskej panve, hĺbených v 60. rokoch minulého storočia bývalým podnikom ČND Hodonín, sme mikrofauisticky prehľadnotili terciérne sedimenty. Z celkovo zrevidovaných 22 vzoriek sme determinovali 192 taxónov foraminifer, na základe ktorých sme dosiahli tieto výsledky:

A. V neogénnych sedimentoch sme zistili príslušnosť k dvom stratigrafickým stupňom: bádenu a sarmatu:

- sarmatské sedimenty *vrábeľského súvrstvia* (foraminiferová zóna *Elphidium reginum*; fide Grill, 1941) sme zistili vo vrte Modrany 1 v intervale 1 050 – 1 056 m;
- **strednobádenské sedimenty** *pozbianskeho súvrstvia* (foraminiferová zóna *Spiroplecatmmi-na carinata*; fide Grill, 1941) sme zistili vo vrte Modrany 1 v intervale 1 098 – 1 303 m;
- **spodnobádenské sedimenty** *bajtavského súvrstvia* (mikrofauisticky prislúchajúce k lagenidovej biozón; fide Grill, 1943) sme zistili v oboch vrtoch, vo vrte Modrany 1 v intervale 1 402 – 1 760 m a vo vrte Nová Vieska 1 v intervale 1 797 – 2 562 m.

B. V paleogénnych sedimentoch sme zistili príslušnosť k stratigrafickým stupňom **kišcel až eger**. Ukázali sa tým najmarkantnejšie rozdiely oproti pôvodnému vyhodnoteniu mikrofauny a tým aj vekovému začleneniu sedimentov vo vrtach Modrany 1 a Nová Vieska 1. Sedimenty pôvodne zaradené do vrchného eocénu (priabón) sme preradili do spodného oligocénu (kišcelské súvrstvie) a spodnooligocénne (rupelské) do spodného bádenu (bajtavské súvrstvie). Korelácia výsledkov revízie je zobrazená na obr. 4.

Významným krokom bolo po prvýkrát použitie paleogénnych litostratigrafických jednotiek budínskeho paleogénu v Dunajskej panve (Gyalog a Császár, 1996). Získaná mikrofauna poukázala na príslušnosť sedimentov ku kišcelskému súvrstviu a doložila ich vek na kišcel až spodný eger. Pod kišcelským súvrstviem je vo vrte Nová Vieska 1 (zhruba od 3 148 m) *hárshegyský pieskovec*.

C. Z planktonických foraminifer smerodajných pre vek kišcel až spodný eger boli vo vrte Modrany 1 v intervale 1 855 – 1 859 m determinované napr. *Chiloguembelina gracillima* (ANDREAE), *Tenuitella munda* (JENKINS), *T. danvilensis* (HOWE et WALLACE), *Tenuitellinata angustiumbilitata* (BOLLI), *Angulogerina angulosa* (WILLIAMSON); vo vrte Nová Vieska 1 v intervale 2 875 – 2 879 m *Chiloguembelina gracillima* (ANDREAE), 3 024 – 3 026,5 m *Globigerinoides primordius*

BLOW et BANNER, 3 075 – 3 078 m *Globigerina aff. anguliofficialis* BLOW a *Tenuitella munda* (JENKINS).

D. Paleogénne asociácie foraminifer vykazujú podobné znaky ako foraminifery získané z oblasti Handlovskej kotliny (Zlinská, 2013; Zlinská a Gross, 2013). V oblasti železovskej priehlbiny je však mikrofauna viac diverzifikovaná. V Handlovskej kotline mala prevahu planktonická zložka, bentos sa vyskytoval len ojedinele. Spoločným znakom tanatocenóz sú vzrastové parametre foraminifer. Väčšinou sú veľmi drobného vzrastu, čo indikuje náhlu a rýchlu zmenu sedimentačných podmienok. V tomto prípade môže ísť o zmenu salinity a teploty.

Prehľad litostratigrafického zaradenia jednotlivých vzoriek zo študovaných vrtov je znázornený v tab. 3.

POĎAKOVANIE

Riešenie revízie sa realizovalo v rámci projektu MŽP 17 13. Poďakovanie patrí RNDR. I. Baráthovi, CSc., za litostratigrafický popis sedimentov v problematických úsekoch vrtov.

Literatúra

- ANDRUSOV, D., BYSTRICKÁ, H. a SCHALEKOVÁ, A., 1954: Správa o geologických výskumoch oblasti medzi Komárnom a Štúrovom. *Geol. Práce, Zpr. (Bratislava)*, 1, 83 – 84.
- ADAM, Z. a DLABAČ, M., 1961: Nové poznatky o tektonike Čs. časti Malé dunajské nížiny. *Věst. Ústř. Úst. geol. (Praha)*, 36, 3.
- ADAM, Z. a DLABAČ, M., 1969: Vysvětlivky k mapám mocností a litofaciálního vývoje Podunajské nížiny. *Zbor. geol. Vied, Západ. Karpaty, ZK (Bratislava)*, 11, 156 – 172.
- BÁLDI, T., 1984: The terminal Eocene and Early Oligocene events in Hungary and the separation of an anoxic, cold Paratethys. *Eclogae geol. Helv., Swiss J. Geosci. (Basel)*, 77, 1, 1 – 27.
- BIELA, A., 1978: Hlboké vrty v zakrytých oblastiach vnútorných Západných Karpát. *Region. geol. Západ. Karpát (Bratislava)*, 10, 224 s.
- BOLTOVSKOY, E. a WRIGHT, R., 1976: Recent Foraminifera, III – XVII. *Junk, Den Haag*, 1 – 515.
- BUDAI, T. a KONRÁD, G., 2011: Magyarország földtana. *Pécs, Fakulta Pécs University of Sciences*, 102 s.
- BUDAY, T., 1957: Zpráva o přehledném výzkumu neogenu pro generální mapu ČSR. Listy Žilina, Bratislava, Česká Třebová. *Manuskript. Bratislava, archiv Št. Geol. Úst. D. Štúra*.
- CICHA, I., 1957: Zpráva o mikrobiostatigrafických výzkumech neogénu na západním a východním úpatí Malých Karpat. In: Buday, T., 1957: Zpráva o přehledném výzkumu neogenu pro generální mapu ČSR. Listy Žilina, Bratislava, Česká Třebová. *Manuskript. Bratislava, archiv Št. Geol. Úst. D. Štúra*.
- CICHA, I., RÖGL, F., ČTYROKÁ, J., RUPP, CH., BAJRAKTAREVIC, Z., BÁLDI, T., BOBRINSKAYA, O. G., DARAKCHIEVA, ST., FUCHS, R., GAGIC, N., GRUZMAN, A. D., HALMAI, J., KRASHENINNIKOV, V. A., KALAC, K., KORECZ-LAKY, I., KRHOVSKY, J., LUCZKOWSKA,

- E., NAGY-GELLAI, A., OLSZEWSKA, B., POPESCU, G. H., REISER, H., SCHMID, M. E., SCHREIBER, O., SEROVA, M. Y., SZEGŐ, E., SZTRAKOS, K., VENGLINSKYI, I. V. a WENGER, W., 1998: Oligocene – Miocene Foraminifera of the Central Paratethys. *Abh. Senckenberg. naturforsch. Gesell. (Frankfurt a. Main)*, 549, 1 – 325, 61 Figs., 3 Tab., 79 Pls.
- CÍLEK, V., 1954: Geologické poměry oligocénu širšího okolí lázní Čízu na jihovýchodním Slovensku. *Rozpr. Čs. Akad. Věd, Ř. mat. přír. Věd (Praha)*, 64, 63 – 90.
- CÍLEK, V., 1960: Neogén v severovýchodním okolí Bratislavy. *Geol. Sbor. (Bratislava)*, 11, 2, 213 – 237.
- CSÁSZÁR, G. (ed.), 1997: Basic Lithostratigraphic Units of Hungary. Charts and Short Descriptions. *Magy. áll. földt. Intéz. (Budapest)*, 114.
- FENDEK, M., FRANKO, O., BRESTENSKÁ, E., KRÁL, M., PRIECHODSKÁ, Z. a VASS, D., 1984: Správa o výskumnom geotermickom pozorovacom vrte GPB-1 Boheľov. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- FORDINÁL, K., 1993a: Genus *Melanopsis* (Gastropoda) in Upper Miocene sediments in the Bratislava area. *Západ. Karpaty, Sér. Paleont. (Bratislava)*, 17, 57 – 69.
- FORDINÁL, K., 1993b: Representatives of genus *Parvidacna* (Bivalvia) in Pannonian sediments of the Bratislava area. *Západ. Karpaty, Sér. Paleont. (Bratislava)*, 17, 71 – 79.
- FORDINÁL, K., 1995: Bivalvia (Dreissenidae, Cardiidae) from Upper Miocene Sediments in Bratislava. *Geol. Práce, Spr. (Bratislava)*, 100, 27 – 36.
- FORDINÁL, K. a TUBA, L., 1988: Ekostratigrafia panónu a pontu veľkej Bratislavy na základe mäkkýšov a ostracód. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- FORDINÁL, K., MATANIOVÁ, D. a TUBA, L., 1990: Ekostratigrafia vrchného miocénu a pliocénu Bratislavy. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- FORDINÁL, K. a TUBA, L., 1992: Biostratigrafické a paleoekologické vyhodnotenie sedimentov územia centrálnej časti Bratislavy. *Geol. Práce, Spr. (Bratislava)*, 96, 63 – 68.
- FORDINÁL, K., NAGY, A. a RAKOVÁ, J., 1993: Nové poznatky o geologickej stavbe centrálnej časti Bratislavy. In: Hamršíd, B. (ed.): *Nové výsledky v terciéru Západných Karpat (Sborník referátů z 10. konference o mladším terciéru, Brno, 27. – 28. 4. 1992)*. *Knih. Zemn. Plyn Nafta (Hodonín)*, 15, 215 – 223.
- FORDINÁL, K., NAGY, A., ZLINSKÁ, A., SLAMKOVÁ, M., HALÁSOVÁ, E. a TÖRÖKOVÁ, I., 2002: New knowledge about stratigraphy of the eastern part of the Danube basin (Želiezovce Depression). *Slovak Geol. Mag. (Bratislava)*, 8, 3 – 4, 259 – 281.
- FORDINÁL, K., ZÁGORŠEK, K. a ZLINSKÁ, A., 2006a: Early Sarmatian biota in the northern part of the Danube Basin (Slovakia). *Geol. Carpath. (Bratislava)*, 57, 2, 123 – 130.
- FORDINÁL, K., ZLINSKÁ, A., VANĚKOVÁ, H., HALÁSOVÁ, E. a CHALUPOVÁ, B., 2006b: Fauna and flora of Sarmatian sediments in borehole TPM - 23B (Danube Basin, Slovakia). In: *Scripta Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis. Geology. 7th Paleontological Conference, October 19 – 20, 2006, Brno, Czech republic*, 30 – 32.
- FRANKO, O., 1978: Správa o výskumnom geotermálnom vrte FGT-1 Topoľníky. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- FRANKO, O., REMŠÍK, A., BODIŠ, D., BRESTENSKÁ, E., PRIECHODSKÁ, Z. a VASS, D., 1984: Správa o výskumnom geotermálnom vrte FGGA-1 Gabčíkovo. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- GAŽA, B., 1968: Záverečná vrtne-geologická správa o pionierskom vrte Vráble-1. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- GAŽA, B., 1970a: Záverečná geologická správa o pionierskom vrte Modrany-1. *Manuskript. Plavecký Štvrtok, archív Nafta*.
- GAŽA, B., 1970b: Záverečná vrtne-geologická správa o hlbokom štruktúrno-prieskumnom vrte Zlaté Moravce-1. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- GAŽA, B. a BEINHAEUEROVÁ, M., 1976: Príspevok ku geológii zlatomoraveckého zálivu. *Miner. slov. (Bratislava)*, 8, 221 – 240.
- GAŽA, B. a BEINHAEUEROVÁ, M., 1977: Tektonika neogénu jv. časti Podunajskej panvy. *Miner. slov. (Bratislava)*, 9, 4, 259 – 274.
- GRADSTEIN, F. a OGG, J., 2004: Geologic Time Scale 2004 – why, how, and where next! *Lethaia (Oslo)*, 37, 175 – 181.
- GRADSTEIN, F. M., OGG, J. G., SCHMITZ, M. D. a OGG, G. M., 2012: The Geologic Time Scale 2012. 2-Volume Set, 1st Ed. *Elsevier*, 1 176 s.
- GRILL, R., 1941: Stratigraphische Untersuchungen mit Hilfe von Mikrofaunen im Wiener Becken und den benachbarten Molasse-Anteilen. *Öl. u. Kohle (Berlin)*, 37, 595 – 602.
- GRILL, R., 1943: Über mikropaläontologische Gliederungsmöglichkeiten im Miozän des Wiener Becken. *Mitt. Reichsamts Bodenforsch. (Wien)*, 6, 33 – 44.
- GYALOG, L. a CSÁSZÁR, G., 1996: A rétegtani egységek rövid leírása. A negyedidőszaknál idősebb rétegtani egységek. In: Gyalog, L. (ed.): *A földtani térképek jelkulcsa és a rétegtani egységek rövid leírása. Mag. áll. földt. Intéz. Alk. Kiadv. (Budapest)*, 187, 65 – 150.
- HÓK, J., KOVÁČ, M., KOVÁČ, P., NAGY, A. a ŠUJAN, M., 1999: Geology and tectonic of the NE part of the Komjatice depression. *Slovak Geol. Mag. (Bratislava)*, 5, 3, 187 – 199.
- HOMOLA, V., 1955: Zpráva o výsledcích opěrné vrtby Báhoň-1, v sz. časti Malé dunajské nížiny. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- HOMOLA, V., 1960: Opěrná vrtba Nová Vieska-I v Malé dunajské nížině. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- HOMOLA, V., SLAVÍKOVÁ, K., SLAVÍK, J., BURKHARDT, R., JANÁK, J., ŠMERAL, J., UHMANN, J., HÜLKOVÁ, J., BURSA, O. a ZOUNKOVÁ, M., 1955: Zpráva o výsledcích opěrné vrtby Báhoň 1 v severozápadní časti Malé dunajské nížiny. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- HRUŠECKÝ, I., POSPIŠIL, L. a ZLINSKÁ, A., 2001 Northern (Slovak) part of the Danube Basin: tectonics, crustal and lithospheric dynamics. *Slovak Geol. Mag. (Bratislava)*, 7, 2, 215 – 216.
- HUDÁČKOVÁ, N., ZLINSKÁ, A. a HALÁSOVÁ, E., 2015: Dierkavce a nanoplanktón Dunajskej panvy z vrtov Nová Vieska-I a Modrany-1. *Spoločný kongres Slovenskej a Českej geologickej spoločnosti, 14. 10. až 17. 10. 2015 v Mikulove, s. 43, ISBN 978-80-87487-14-3*.
- JANÁČEK, J., 1966a: Tektonický výskum v oblasti vodného diela Dunaj. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- JANÁČEK, J., 1966b: Problémy tektonické stavby oblasti vodního díla Dunaj. *Geol. Práce, Zpr. (Bratislava)*, 38, 127 – 135.
- JANÁČEK, J., 1967: Tektonický výskum v oblasti vodného diela DUNAJ. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.

- JIRÍČEK, R., 1972: Problém hranice sarmat/panón ve Videnské, podunajské a východoslovenské pánvi. *Miner. slov. (Bratislava)*, 14, 4, 39 – 81.
- KEITH, J. F. JR., VASS, D., KANES, W. H., PERESZLÉNYI, M., KOVÁČ, M. a KRÁL, J., 1989: Sedimentary basins of Slovakia, Part II., Final report on the Hydrocarbon potential of Danube Lowland Basin, vol. 1. *Manuscript Univ. South Carolina, ESRI, Technical Report 89-0019*, 1 – 143.
- KOVÁČ, M., 2000: Geodynamický, paleogeografický a štruktúrny vývoj Karpatsko-panónskeho regiónu v miocéne: nový pohľad na neogénne panvy Slovenska. *Bratislava, Veda*, 5 – 202.
- KOVÁČ, M., SYNAK, R., FORDINÁL, K. a JONIAK, P., 2010: Významné eventy v paleogeografii severnej časti Dunajskej panvy – nástroj na upresnenie stratigrafie jej vrchnomiocénnej a pliocénnej výplne. *Acta Geol. Slov. (Bratislava)*, 2, 1, 23 – 35.
- KOVÁČ, M., SYNAK, R., FORDINÁL, K., JONIAK, P., TOTH, Cs., VOJTKO, R., NAGY, A., BARÁTH, I., MAGLAY, J. a MINÁR, J., 2011: Late Miocene and Pliocene history of the Danube Basin: inferred from development of depositional systems and timing of sedimentary facies changes. *Geol. Carpath. (Bratislava)*, 62, 6, 519 – 534.
- KOVÁČ, M., BARANYI, V., HALÁSOVÁ, E., HUDÁČKOVÁ, N., HÓK, J., KOVÁČOVÁ, M., RYBÁR, S., ŠARINOVÁ, K., ŠUJAN, M. a ZLINSKÁ, A., 2015: Cenozoic Sedimentary Record at the Central Western Carpathian and Northern Pannonian Domains Junction: Interpretation of a Complex Geodynamic Evolution. In: *Neogene of the Paratethyan Region. Workshop on the Neogene of Central and South-Eastern Europe an RCMNS Interim Colloquium. Programme, abstracts and field trip Guidebook. 31. May – 3. June 2015, Orfű, Hungary. Budapest, Hungarian Geological Society*, s. 49.
- LEHOTAYOVÁ, R. a ONDREJČKOVÁ, A., 1966: Fauna lanzendorfskej série od Bajtavy (jv. časť Podunajskej nížiny). *Geol. Práce, Zpr. (Bratislava)*, 40, 39 – 54.
- LUNGA, S., 1965: Geologická zpráva o štruktúrnym prieskume v oblasti Bučan za rok 1964. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- MAJON, L., 1960: Paleogene Foraminifera horizons of Hungary. *Földt. Köz. (Budapest)*, 60, 3, 355 – 362.
- MARTINI, E., 1971: Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: *Proceeding of 2nd planktonic conference, Roma 1970*, 739 – 785.
- MATĚJKA, A., 1949: Přehled geologických poměrů Malé Dunajské nížiny (Vysvětlivky k mapě 1 : 200 000. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- NAGY, A., FORDINÁL, K., BRZOBHATÝ, R., UHER, P. a RAKOVÁ, J., 1995: Vrchný miocén juhovýchodného okraja Malých Karpát (vrt Ma-1, Bratislava). *Miner. slov. (Bratislava)*, 27, 113 – 132.
- PÁLFAVI, F., 1975: Geotermálny vrt Bratislava. HGB-1. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- PAPP, A., CÍCHA, I., SENEŠ, J. a STEININGER, F. P. (eds.), 1978: Chronostratigraphie und Neostatotypen, Miozän M₄, Badenien. *Bratislava, Veda*, 1 – 594.
- PAPP, A., MARINESCU, F. a SENEŠ, J. (eds.), 1974: Chronostratigraphie und Neostatotypen, Miozän der Zentralen Paratethys, Bd. IV, M5 – Sarmatien (sensu Suess, E., 1866). *Bratislava*, 707 s.
- PAPŠÍKOVÁ, M., 1989: Biostratigrafické vyhodnotenie vrtov z územia Bratislavy (centrálna mestská zóna) na základe palynomorf. *Region. geol. Západ. Karpát (Bratislava)*, 25, 41 – 42.
- PILLER, W., HARZHAUSER, M. a MANDIC, O., 2007: Miocene Central Paratethys stratigraphy: Current status and future directions of stratigraphy. *Stratigraphy*, 4, 2, 3, 151 – 168.
- POAG, C. W., 1981: Ecologic Atlas of Benthic Foraminifera of the Gulf of Mexico, V – VIII. *Marine Science International, Woods Hole/Mass.*, 1 – 175.
- POKORNÝ, V., 1946: K mikrostratigrafii neogénu pannonské pánve v okolí Ivánky na Slovensku. *Věst. St. geol. Úst. Čs. Republ. (Praha)*, 21, 262 – 273.
- REMŠÍK, A., FRANKO, O., BODIŠ, D., BRESTENSKÁ, E., PRIECHODSKÁ, Z. a VASS, D., 1984: Správa o výskumnom geotermálnom vrte FGV-1 Vlčany. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- ROYDEN, L. H., HORVÁTH, F. a RUMPLER, J., 1983: Evolution of the Pannonian basin systém. I. Tectonics. *Tectonics*, 2, 61 – 90.
- RYBÁR, S., ŠUJAN, M. a KOVÁČ, M., 2013: Middle Miocene sedimentary facies and depositional environments of Blatné depression. *Miner. slov. (Bratislava)*, 4, *Geovestník*, 15.
- RYBÁR, S., ŠARINOVÁ, K., HALÁSOVÁ, E., HUDÁČKOVÁ, N., ZLINSKÁ, A., KOVÁČ, M. a ŠUJAN, M., 2014: Biostratigraphy, sedimentology and paleoenvironment of the middle miocene Želiezovce depression of the Danube basin. In: *9th ESSEWECA conference. Environmental, sedimentary and structural evolution of the western Carpathians. Abstract Book. November, 5 – 7, 2014, Smolenice, Slovakia. Bratislava, Geophysical Inst., Slovak Acad. Sci.*, 54 – 55.
- RYBÁR, S., HALÁSOVÁ, E., HUDÁČKOVÁ, N., KOVÁČ, M., KOVÁČOVÁ, M., ŠARINOVÁ, K. a ŠUJAN, M., 2015: Biostratigraphy, sedimentology and paleoenvironments of the northern Danube Basin: Ratkovce 1 well case study. *Geol. Carpath. (Bratislava)*, 66, 1, 51 – 67.
- SENEŠ, J., 1953: Spodnosarmatská fauna pri Malej nad Hronom. *Geol. Sbor. (Bratislava)*, 3, 3 – 4, 193 – 257.
- SENEŠ, J., 1964: Otázka latorfu, problém veku a korelácia juhoslovenského a severomaďarského oligocénu. *Geol. Práce, Zpr. (Bratislava)*, 33, 71 – 85.
- SENEŠ, J., FRANKO, O., KOŠTALIK, J. a PORUBSKÝ, A., 1962: Vysvětlivky k prehľadnej geologickej mape ČSSR 1 : 200 000, list Nové Zámky. *Bratislava, Geofond*, 1 – 106.
- SENEŠ, J. et al., 1971: Korrelation des Miozäns der zentralen Paratethys (Stand, 1970). *Geol. Zbor. Geol. carpath. (Bratislava)*, 22, 1, 3 – 9.
- SLAVÍKOVÁ, K., 1954: Mikrostratigrafie pannonské pánve. Mikro-biostratigrafie západní části Malodunajské nížiny (hlubinná vrtba Báhoň). *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- STEININGER, F. F., SENEŠ, J., KLEEMANN, K. a RÖGL, F. (eds.), 1985: Neogene of the mediterranean Tethys and Paratethys, vol. 1, 2, *University of Vienna*, 1: XIV + 189 s., 2: XXVI + 536 s.
- ŠUJAN, M., BRAUCHER, R., KOVÁČ, M., BOURLÈS, D. L., RYBÁR, S., GUILLOU, V. a HUDÁČKOVÁ, N., 2016: Application of the authigenic 10Be/9Be dating method to Late Miocene – Pliocene sequences in the northern Danube Basin (Pannonian Basin System): Confirmation of heterochronous evolution of sedimentary environments. *Glob. Planet. Change*, 137, 35 – 53.
- TARI, G., BÁLDI, T. a BÁLDI-BEKE, M., 1993: Paleogene retroarc flexural basin beneath the Neogene Pannonian Basin:

- a geodynamic model. *Tectonophysics (Amsterdam)*, 226, 433 – 455.
- ŠVAGROVSKÝ, J., 1971: Das Sarmat der Tschechoslowakei und seine Mollusken fauna. *Acta geol. geogr. Univ. Comen., Geol. (Bratislava)*, 20, 1 – 476.
- VASS, D., 1971: Sedimentological characterisation of the Plášťovce beds (Southern Slovakia). *Geol. Zbor. Geol. carpath. (Bratislava)*, 22, 1, 25 – 47.
- VASS, D., 1989: Alpine Molase basins as a mirror of genesis and development of block structure in the West Carpathians. *Z. geol. Wiss. (Berlin)*, 17, 9, 879 – 885.
- VASS, D., 2002: Litostratigrafia Západných Karpát: neogén a budínsky paleogén. *Bratislava, Št. Geol. Úst. D. Štúra*, 202 s.
- VASS, D., BEGAN, A., GROSS, P., KAHAN, Š., KRYSTEK, I., KÖHLER, E., LEXA, J., NEMČOK, J., RŮŽIČKA, M. a VAŠKOVSKÝ, I., 1988: Regionálne geologické členenie Západných Karpát a severných výbežkov Panónskej panvy na území ČSSR, mapa 1 : 500 000. *Bratislava, Geol. Úst. D. Štúra*.
- VAŠKOVSKÝ, I. a VAŠKOVSKÁ, E., 1977: Regionálny kvartérno-geologický výskum Žitného ostrova. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- VLASKOVÁ, D., ŠECHNY, J., MENZELOVÁ, O., KOZAKOVIČ, M. a MOTLIKOVÁ, M., 1986: Orientačný prieskum v CMČ Bratislavy. *Manuskript. Bratislava, archív IGHP*.
- WALTON, W. R., 1964: Recent Foraminiferal Ecology and Paleocology. In: *Imbrie, J. a Newell, N. (Hrsg.): Approaches to Paleocology, New York – London – Sidney, J. Wiley*, 151 – 237.
- ZLINSKÁ, A., 1993: Mikrofauna bádenu z okolia Malej nad Hronom (jv. časť podunajskej panvy). *Geol. Práce, Spr. (Bratislava)*, 97, 73 – 78.
- ZLINSKÁ, A., 1997: Biostratigrafické vyhodnotenie vzoriek z regiónu Podunajská nížina-východ. *Manuskript. Bratislava, archív Št. Geol. Úst. D. Štúra*.
- ZLINSKÁ, A., 2004: Zhodnotenie foraminiferových asociácií vo vrte HGŽ-3 (Želiezovce, Dunajská panva, južné Slovensko). *Geol. Práce, Spr. (Bratislava)*, 109, 101 – 106.
- ZLINSKÁ, A., 2013: Prvý nález mikrofauny z chrenoveckých vrstiev bielopotockého súvrstvia Handlovskej kotliny. *Acta Geol. Slov. (Bratislava)*, 5, 1, 69 – 82.
- ZLINSKÁ, A., HUDÁČKOVÁ, N., HALÁSOVÁ, E. a SLAMKOVÁ, M., 2002: Upper Badenian Sediments of the Danube and Vienna Basins (Central Paratethys Area) – Palaeoecological Trends. In: *The 3rd International Congress Environmental Micropaleontology, Microbiology and Meiobentology 2002, University of Vienna. Vienna, Inst. Palaeont.*, 209 – 210.
- ZLINSKÁ, A., FORDINÁL, K. a NAGY, A., 2005: Development of sediments of Karpathian age in the northern Part of the Danube Basin (Slovakia). In: *Patterns and Processes in the Neogene of the Mediterranean Region. 12th Congress R.C.M.N.S. 6. – 11. September 2005, Vienna*, 250 – 251.
- ZLINSKÁ, A. a GROSS, P., 2013: Vek a litologická charakteristika paleogénnych usadenín Handlovskej kotliny na základe reinterpretácie vrtu FGHn-1 Handlová. *Acta Geol. Slov. (Bratislava)*, 5, 2, 141 – 153.

Tertiary microfauna in deep wells of the Želiezovce depression (Danube Basin)

From two selected deep oil wells (Modrany 1 and Nová Vieska 1; Fig. 1) in the Danube Basin, we have microfaunistically re-evaluated Tertiary sediments. In the past, the oil boreholes were realized with the intention to determine the stratigraphy and lithology of Neogene sediments, and to verify their oil and gas carrying capacity. From totally 22 re-evaluated samples we have determined 192 foraminiferal taxons, on the basis of which we have detected in Neogene sediments the affiliation to the two stratigraphic stages: Badenian and Sarmatian. Sarmatian sediments of *Vráble Formation* (foraminiferal zone of *Elphidium reginum*; fide Grill, 1941) were detected in the depth interval 1 050 – 1 056 m in borehole Modrany 1. Middle Badenian sediments of the *Pozba Formation* (foraminiferal zone of *Spiroplectammia carinata*; fide zone Grill, 1941) were detected in depth interval 1 098 – 1 303 m in borehole Modrany 1 and the Lower Badenian sediments of *Bajtava Formation* (microfaunistically belonging to the *Lagenida* biozone; fide Grill, 1943) were detected in both boreho-

les. In borehole Modrany 1, they were detected in interval 1 402 – 1 760 m and in the borehole Nová Vieska in interval 1 797 – 2 562 m.

In Paleogene sediments we have found the affiliation to the stratigraphic stages Kiscellian to Egerian, whereby thus appeared the most striking differences in comparison to the primeval microfauna evaluation, and thereby also to the age incorporation of sediments in the wells Modrany 1 and Nová Vieska 1. Sediments originally incorporated to the Upper Eocene (Priabonian) we reclassified to the Lower Oligocene (*Kiscellian Formation*) and those of Lower Oligocene (Rupelian) into the Lower Badenian (*Bajtava Formation*).

An important step was that in the Danube Basin, there were for the first time used Paleogene lithostratigraphic units of the Buda Paleogene (Gyalog and Császár, 1996). The obtained microfauna thus pointed out and proved the affiliation of sediments to Kiscellian Formation. In a borehole Nová Vieska 1, there is developed under Kiscellian Formation a *Hárshegyi sandstone* (approx. from 3 148 m).

Chemické zloženie podzemnej vody a úmrtnosť na kardiovaskulárne ochorenia v Slovenskej republike

¹VERONIKA CVEČKOVÁ, ¹KATARÍNA FAJČÍKOVÁ, ²EUBOMÍR JURKOVIČ, ³BEÁTA STEHLÍKOVÁ
a ¹STANISLAV RAPANT

¹Štátny geologický ústav Dionýza Štúra, Mlynská dolina 1, 817 04 Bratislava, Slovenská republika


²Prirodovedecká fakulta, Univerzita Komenského, Mlynská dolina 16, 842 15 Bratislava,
Slovenská republika

³Fakulta ekonómie a podnikania, Paneurópska univerzita, Tematínska 10, 851 05 Bratislava,
Slovenská republika

Chemical composition of groundwater and mortality for cardiovascular diseases in the Slovak Republic

Abstract: This study deals with the analysis of relationship between chemical composition of the groundwater/drinking water and the data on mortality for cardiovascular diseases (MCVD) in the Slovak Republic. Primary data consist of the Slovak national database of groundwater analyses (20,339 chemical analyses, 34 chemical elements/compounds) and data on MCVD (8 health indicators) collected for the 10-year period (1994–2003). The chemical and health data were unified in the same form and expressed as the mean values for each of 2883 municipalities within the Slovak Republic. Pearson and Spearman correlation as well as artificial neural network (ANN) methods were used for analysis of the relationship between chemical composition of groundwater/drinking water and MCVD. The most significant chemical elements having influence on MCVD were identified together with their limit values (limit and optimal contents). Based on the results of calculations, made through the neural networks, the following chemical elements/parameters in the groundwater were defined as the most significant for MCVD: $\text{Ca}^{2+} + \text{Mg}^{2+}$ (mmol l^{-1}), Ca^{2+} , Mg^{2+} , TDS, Cl^{-} , HCO_3^{-} , SO_4^{2-} , NO_3^{-} , SiO_2 and PO_4^{3-} . The obtained results document the highest relationship between MCVD and the groundwater contents of $\text{Ca}^{2+} + \text{Mg}^{2+}$ (mmol l^{-1}), Ca^{2+} and Mg^{2+} . Following limit values were set for the most significant groundwater chemicals/parameters: $\text{Ca}^{2+} + \text{Mg}^{2+}$ 4.4 – 7.6 mmol l^{-1} , Ca^{2+} > 89.4 mg l^{-1} and Mg^{2+} 42–78.1 mg l^{-1} . At these concentration ranges the mortality for cardiovascular diseases in the Slovak Republic reaches the lowest levels. These limit values are about twice higher in comparison with the current Slovak valid guideline values for the drinking water.

Key words: Groundwater, drinking water quality, cardiovascular disease mortality, neural network, Slovak Republic



ÚVOD

Kardiovaskulárne ochorenia (CVD) sú na Slovensku dlhodobo najčastejšou príčinou úmrtí obyvateľstva. Predstavujú približne 50 % zo všetkých príčin smrti (NHIC 2012; OECD 2013). Za hlavné rizikové faktory CVD sa považujú stres, genetická predispozícia, nadváha, pravidelné fajčenie, nadmerný príjem alkoholu, nesprávne

stravovacie návyky a taktiež environmentálne faktory. K najdôležitejším environmentálnym faktorom patrí chemické zloženie, resp. úroveň kontaminácie podzemnej vody (najmä pitnej), pôdy a ovzdušia.

Predmetom článku je problematika vplyvu chemického zloženia podzemnej (pitnej) vody na rôzne indikátory úmrtnosti na CVD (relatívna a štandardizovaná úmrtnosť a potenciálne roky strateného života). Podzemná voda

predstavuje na Slovensku najvýznamnejší zdroj pitnej vody na zásobovanie populácie Slovenska, a to približne pre 90 % obyvateľov (Klinda a Lieskovská, 2010). Približne 10 % populácie Slovenska používa vodu z individuálnych studní na pitné účely a varenie. Okolo 50 % populácie je zásobovaných pitnou vodou z miestnych vodárenských zdrojov, spravovaných lokálnymi vodárenskými spoločnosťami, ktoré sa vyznačujú nízkou výdatnosťou (menej ako $10 \text{ l} \cdot \text{s}^{-1}$), zachytených a distribuovaných vodovodnými rozvodmi v blízkosti osídlených oblastí. Len v južnej časti Slovenska (v kvartérnych sedimentoch) je populácia zásobovaná z veľkých vodárenských zdrojov na vzdialenosť 50 – 100 km. V tejto práci považujeme podzemnú a pitnú vodu za jeden celok. Sme si vedomí istých nepresností, ktoré s tým súvisia a môžu limitovať naše výsledky. Veľkosť databázy (viac ako 20 000 chemických analýz, viac ako 30 chemických prvkov/parametrov) však do značnej miery redukuje možné neistoty. Podiel balenej vody v rámci pitného režimu ľudí sme neboli schopní zhodnotiť.

Úmrtnosť, resp. zvýšená incidencia na CVD sa mnohokrát spája s nadbytkom, resp. deficitom viacerých chemických prvkov v podzemnej vode využívané na pitné účely. Zrejme najčastejšie sa CVD spájajú s deficitným obsahom Ca^{2+} a Mg^{2+} , resp. nízkou „tvrdosťou“ vody (Dawson et al., 1978; Shaper et al., 1980; Rylander et al., 1991; Rahman a Husain, 2011). Existujú však práce, ktoré túto závislosť nepotvrdzujú, napr. Maheswaran et al. (1999). Mnohé ďalšie práce spájajú CVD so zvýšeným obsahom potenciálne toxických prvkov (PTE) v podzemnej (pitnej) vode, najmä As, Cd, Pb, Sb a Ba (Schroeder a Kraemer, 1974; Bhatnagar, 2006; Mitchell et al., 2011; Sturchio et al., 2013).

V článku sa zaoberáme hodnotením vzťahu medzi širokou škálou chemických prvkov bežne analyzovaných v podzemnej vode a CVD. Hlavným prínosom tejto práce je využitie inovatívnej a jedinečnej metódy – umelej inteligencie vo sfére štatistických metód – na analýzu vzťahu medzi geologickým prostredím a ľudským zdravím.

Hlavným cieľom práce je určiť, ktoré chemické prvky v podzemnej vode najviac vplyvajú na CVD v závislosti od geologickej stavby územia, a zároveň určiť ich limitnú koncentráciu (optimálnu, maximálne prípustnú a minimálne potrebnú), pri ktorej je úmrtnosť na CVD na Slovensku čo najnižšia.

MATERIÁL A METÓDY

Chemické zloženie podzemnej vody

Hlavným zdrojom informácií o chemickom zložení podzemnej vody boli údaje z národného environmentálno-geochemického mapovania, a to najmä z *Geochemického atlasu Slovenskej republiky*, časť *Podzemné vody*, a environmentálno-geochemických máp regiónov Slovenskej republiky (Vrana et al., 1997; Rapant et al., 1999). Tie boli doplnené najmä o údaje z národného monitoringu podzemnej vody, hydrogeochemických máp a ďalších regionálnych a lokálnych geochemických prác (SHMÚ – www.shmu.sk/en; Kordík et al., 2000). V našej databáze sú zahrnuté prakticky všetky zdroje podzemnej vody, ktoré sa používajú na hromadné zásobovania obyvateľstva pitnou vodou. Celkový počet zhromaždených chemických analýz podzemnej vody bol 20 339. Zahrnuté sú chemické analýzy vody od roku 1991, keď sa začalo moderné environmentálno-geochemické mapovanie Slovenskej republiky v rámci programu IGCP 360 *Geochemical Correlation Programme* (Darnley et al., 1995). Celkovo boli zhromaždené údaje za roky 1991 – 2010. V prípade viacerých analýz z rovnakého zdroja podzemnej vody sme použili najreprezentatívnejšiu chemickú analýzu. Hustota vzorkovania podzemnej vody bola približne 1 vzorka na $2,5 \text{ km}^2$. Obsah chemických prvkov/parametrov analyzovaných v životnom prostredí, v našom prípade v podzemnej vode, definujeme ako tzv. environmentálne indikátory – EI (Rapant et al., 2010).

Údaje o chemickom zložení vody sme upravili do takej podoby, aby sa dali prepojiť s údajmi o úmrtnosti


Tab. 1

Charakteristika chemického zloženia podzemnej vody Slovenska (priemerné hodnoty).
Characteristics of chemical composition of the groundwater in the Slovak Republic (mean values).

PODZEMNÁ VODA (n = 20 339)												
pH	MIN	CHSK _{Mn}	$\text{Ca}^{2+} + \text{Mg}^{2+}$	Li	Na^+	K^+	Ca^{2+}	Mg^{2+}	Sr	Fe	Mn	NH_4^-
7,33	629,75	2,18	3,5	0,019	20,34	11,10	93,56	28,29	0,36	0,17	0,12	0,10
F^-	Cl^-	SO_4^{2-}	NO_2^-	NO_3^-	PO_4^{3-}	HCO_3^-	SiO_2	Cr	Cu	Zn	As	Cd
0,13	32,96	79,32	0,11	38,76	0,20	303,85	18,21	0,001 3	0,002 6	0,267 3	0,001 9	0,001 0
Se	Pb	Hg	Ba	Al	Sb	^{222}Rn	^{226}Ra					
0,001 0	0,001 4	0,000 1	0,074 7	0,029 7	0,000 9	14,46	0,053					

Pozn.: MIN – celková mineralizácia, údaje okrem pH v $\text{mg} \cdot \text{l}^{-1}$, $\text{Ca}^{2+} + \text{Mg}^{2+}$ v $\text{mmol} \cdot \text{l}^{-1}$, ^{222}Rn a ^{226}Ra v $\text{Bq} \cdot \text{l}^{-1}$

Note: Data except of pH in $\text{mg} \cdot \text{l}^{-1}$, $\text{Ca}^{2+} + \text{Mg}^{2+}$ in $\text{mmol} \cdot \text{l}^{-1}$, ^{222}Rn and ^{226}Ra in $\text{Bq} \cdot \text{l}^{-1}$



Obr. 1. Distribúcia Ca^{2+} a Mg^{2+} ($\text{mmol} \cdot \text{l}^{-1}$) v podzemnej vode obcí Slovenskej republiky.

Fig. 1. Ca^{2+} and Mg^{2+} ($\text{mmol} \cdot \text{l}^{-1}$) distribution in the groundwater of the Slovak Republic at municipality level.

na CVD, ktoré sú vyjadrené ako hodnoty pre najmenšie územnosprávne jednotky SR – obce (spolu 2 883 obcí).

Na základe vstupných analytických údajov bola z celého územia SR spracovaná plošná distribúcia jednotlivých chemických prvkov/zlúčenín formou tzv. pixelových máp (1 pixel s plochou 1 km^2) prostredníctvom softvéru MapInfo Professional 9.0. Pre každý pixel bola vypočítaná priemerná koncentrácia prvku metódou inverzných vzdialeností od stredu pixela k 10 najbližším vzorkám. Priemerné hodnoty koncentrácie chemických prvkov/zlúčenín/parametrov v jednotlivých administratívnych celkoch SR (obce, okresy, celé územie SR) sa vypočítali ako aritmetický priemer všetkých pixelov spadajúcich do hraníc príslušných celkov. Pixely, ktoré len čiastočne spadali do administratívneho celku, boli zahrnuté do výpočtov.

Hodnoty priemernej koncentrácie hodnotených chemických prvkov/parametrov v podzemnej vode SR (EI) sú uvedené v tab. 1 (Rapant et al., 2014a). Príklad mapového vyjadrenia chemického zloženia podzemnej vody Slovenskej republiky v prípade parametrov $\text{Ca}^{2+} + \text{Mg}^{2+}$ ($\text{mmol} \cdot \text{l}^{-1}$) je uvedený na obr. 1. Ostatné environmentálne indikátory sú sprístupnené na internetovej stránke www.geology.sk/geohealth.

Úmrtnosť na kardiovaskulárne ochorenia

V zmysle medzinárodnej klasifikácie chorôb (MKCH, 10. revízia) (www.czisk.sk) kardiovaskulárne ochorenia zahŕňajú choroby obehovej sústavy, diagnózy I00 – I99. Ide najmä o hypertenziu, ischemické choroby srdca (in-

farkty), cievne choroby mozgu, choroby tepien, žíl a iné nešpecifikované choroby obehovej sústavy. Údaje hodnotiace incidenciu (výskyt) CVD neboli dostupné.

Údaje o CVD, ktoré hodnotíme v tejto práci, sú charakterizované ako zdravotné indikátory (ZI) pri vybraných diagnózach. Predstavujú priemerné hodnoty za roky 1994 – 2003, a teda priemerné hodnoty za jednotlivé obce Slovenskej republiky (2 883 obcí). Zdrojom údajov bola databáza Štatistického úradu Slovenskej republiky (www.statistics.sk).

Na charakteristiku úmrtnosti na CVD sme spracovali údaje týkajúce sa nasledujúcich skupinových diagnóz: I00 – I99 (všetky ochorenia obehového systému), I21 – I25 (ischemické choroby srdca) a I63 – I64 (mozgové porážky a infarkty). Úmrtnosť na jednotlivé diagnózy sme prepočítali do podoby zdravotných indikátorov.

Všetky zdravotné indikátory boli vypočítané formou kumulatívnej funkcie za roky 1994 až 2003, teda za 10-ročné obdobie, pričom všetky prípady úmrtí boli spočítané a všetky počty obyvateľov boli vyjadrené ako osobo-roky (počet obyvateľov k 31. decembru príslušného roku) za každý hodnotený územný celok. Metodika výpočtu a štandardizácie zdravotných indikátorov (REI00 – I99 až PYLL121 – 125; tab. 2) je v súlade s odporúčaniami Svetovej zdravotníckej organizácie (WHO) a iných autorov (Beaglehole et al., 1993; Jeníček, 1995; Last, 2001; Bencko et al., 2003a, b).

Jednotlivé diagnózy CVD sme prepočítali do podoby zdravotných indikátorov vyjadrujúcich relatívnu a štandardizovanú úmrtnosť na vybranú príčinu úmrtia a ako

potenciálne stratené roky života. V našej práci používame na hodnotenie zdravotné indikátory uvedené v tabuľke 2, kde je uvedený aj postup výpočtu jednotlivých indikátorov. V grafickej forme znázorňujeme relatívnu úmrtnosť na všetky druhy CVD (ReI00 – I99) v Slovenskej republike (obr. 2). Ostatné zdravotné indikátory sú dostupné na internetovej stránke www.geology.sk/geohealth.

Rozdelenie environmentálnych a zdravotných indikátorov podľa geologickej stavby

Geologická stavba SR je relatívne veľmi komplikovaná. Vyznačuje sa striedaním hornín rôznej genézy a veku, a teda aj rôzneho mineralogicko-petrografického charakteru a tým aj s rôznym geochemickým pozadím (Kohút et al., 1999). Výsledkom tejto skutočnosti je aj veľmi rozdielne chemické zloženie podzemnej/pitnej vody. Predpokladáme, že má aj rozdielny vplyv na výskyt CVD, resp. na ich úmrtnosť.

Geologická stavba SR bola v práci Rapanta et al. (2014a) rozčlenená na 8 hlavných celkov (obr. 3), a to:

- 1 – paleozoikum: hlavne metasedimenty a metavulkanity,
- 2 – kryštalinikum: hlavne granitoidy, ruly a migmatity,
- 3 – karbonatické mezozoikum a bazálny paleogén: hlavne vápence, dolomity a vápnité zlepenice,
- 4 – karbonaticko-silikátové mezozoikum a paleogén: hlavne sliene, slienité vápence, dolomity, pieskovce a bridlice,
- 5 – flyšový paleogén: hlavne pieskovce, bridlice a ílovce,

6 – neovulkanity: hlavne andezity, bazalty a ich vulkanoklastiká,

7 – neogén: hlavne íly, ílovce, zlepenice, piesky a štrky,

8 – kvartér: hlavne štrky, piesky, íly a úlomky hornín.

Poradie geologických celkov z hľadiska zdravotných indikátorov (od najpriaznivejšieho po najnepriaznivejšie) bolo takéto: flyšový paleogén – karbonaticko-silikátové mezozoikum a paleogén – karbonatické mezozoikum a bazálny paleogén – neogén – kvartér – kryštalinikum – paleozoikum – neovulkanity. Ako najpriaznivejšie geologické prostredie pre ľudské zdravie boli definované karbonatické horniny a najnepriaznivejšie silikátové horniny (Rapant et al., 2014a).

Podľa tohto rozčlenenia geologickej stavby sme následne rozčlenili chemické zloženie podzemnej vody (environmentálne indikátory – EI) a taktiež ukazovatele úmrtnosti na CVD (zdravotné indikátory – ZI), ktoré sme podrobili štatistickej analýze formou štatistických metód uvedených v nasledujúcom texte.

Štatistická analýza

Na štatistické spracovanie vzťahu údajov o EI a ZI sme použili klasické štatistické metódy, a to Pearsonov korelačný koeficient a Spearmanov korelačný koeficient. Štatistickú významnosť vypočítaných korelácií hodnotíme pomocou jeho hodnoty P takto: $P < 0,05$ – overená závislosť (+), $P < 0,01$ – vysoká závislosť (++), $P < 0,001$ – veľmi vysoká závislosť (+++).

Tab. 2
Charakteristika hodnotených zdravotných indikátorov pri úmrtnosti na kardiovaskulárne ochorenia.
Characteristics of evaluated health indicators for cardiovascular mortality.


Zdravotný indikátor	Popis indikátora	Spôsob výpočtu	Priemer SR*
Relatívna úmrtnosť na vybranú príčinu úmrtia			
ReI00 – I99	obehový systém	100 000 x [počet úmrtí na vybrané ochorenie/ počet obyvateľov]	531,05
ReI21 – I25	ischemické choroby srdca		269,82
ReI63 – I64	mozgové porážky a infarkty		63,57
Štandardizovaná úmrtnosť na vybranú príčinu úmrtia			
SMRI00 – I99	obehový systém	Nepriamo vekovo štandardizovaná úmrtnosť na slovenský štandard	
(19 vekových skupín)	100		
SMRI21 – I25	ischemické choroby srdca		100
SMRI63 – I64	mozgové porážky a infarkty		100
Potenciálne roky strateného života pri vybranej príčine úmrtia			
PYLLI00 – I99	obehový systém	100 000 x [súčet nedožítých rokov do veku 65 rokov (úmrtia vo veku 1 až 64 rokov) pri úmrtí na vybranú príčinu/počet obyvateľov]	866,19
PYLLI21 – I25	ischemické choroby srdca		396,32

Pozn.: zdravotné indikátory klasifikované v zmysle Medzinárodnej klasifikácie chorôb, MKCH, 10. revízia (www.czisk.sk),

* priemer pre Slovenskú republiku za obdobie 1994 – 2003


Note: Health indicators are classified according to International classification of diseases, ICD 10th revision (www.czisk.sk),

* mean for the Slovak Republic for the period 1994 – 2003



Obr. 2. Relatívna úmrtnosť na kardiovaskulárne ochorenia na Slovensku.
Fig. 2. Relative mortality for cardiovascular diseases in the Slovak Republic.

Rozčlenenie geologickej stavby Slovenskej republiky na hlavné geologické celky



Obr. 3. Rozčlenenie geologickej stavby Slovenskej republiky na hlavné geologické celky.
Fig. 3. Division of geological structure of the Slovak Republic in main geological units.

Neurónové siete

Sledovanie vzťahov medzi dvomi rôznymi premennými je doménou štatistiky. Výber vhodnej štatistickej metódy s cieľom spojenia dvoch databáz si však vyžaduje veľmi korektný prístup k získavaniu relevantných vzťahov závislosti. Na vyjadrenie intenzity stochastickej závislosti medzi dvomi premennými sa používajú korelačné koeficienty, ktoré vyjadrujú závislosť vzťahov medzi skúmanými atribútmi. Klasické Pearsonove korelačné koeficienty vyjadrujú mieru jednoduchšej lineárnej závislosti dvoch premenných. Spearmanove korelačné koeficienty sú mierou monotónnej závislosti. Naše údaje však nemajú

normálne rozdelenie, sú rozdelené nerovnomerne, často zaťažené chybou, neúplné a vykazujú vysokú variabilitu. Neistoty v prípade údajov o zdravotných indikátoroch, ktoré môžu modifikovať naše výsledky, súvisia najmä so zostavovaním štatistických hlásení. V prípade viacerých diagnóz sa ako príčina smrti uvádza väčšinou tá diagnóza, ktorá je v poradí uvedená ako prvá (napr. kardiovaskulárne ochorenie alebo zlyhanie obličiek). Táto diagnóza však nemusí byť vždy tou hlavnou, ktorá zapríčinila smrť pacienta.

Naše údaje majú všetky náležitosti bežného života, často sú neúplné, niekedy chybné. Bolo by preto nesprávne predpokladať existenciu funkčného vzťahu. Použitie

Tab. 3

Priemerné hodnoty zdravotných indikátorov kardiovaskulárnych ochorení vo vybraných oblastiach Slovenskej republiky.
Mean values for health indicators of cardiovascular diseases in selected areas of the Slovak Republic.

Geologický celok/okres	1	6	4	5	Krupina	Bardejov	SR
Zdravotný indikátor	n = 100	n = 303	n = 154	n = 727	n = 36	n = 86	
Rel100 – I99	569,73	638,78	505,07	463,32	889,20	492,82	531,05
Rel121 – I25	337,11	315,99	252,94	257,49	389,91	247,71	269,82
Rel163 – I64	51,95	93,45	77,04	41,58	133,66	56,00	63,57
SMRI100 – I99	111,73	108,50	98,86	100,03	131,06	100,71	100
SMRI121 – I25	128,21	101,39	97,54	109,94	119,09	98,92	100
SMRI163 – I64	84,78	125,53	121,31	72,61	168,12	97,26	100
PYLLI100 – I99	1 052,18	1 050,95	937,66	831,99	1 518,17	779,92	866,19
PYLLI121 – I25	548,98	488,65	395,63	372,17	683,12	310,97	396,32

Pozn.: 1 – paleozoikum, 4 – karbonatiko-silikátové mezozoikum a paleogén, 5 – flyšový paleogén, 6 – neovulkanity, SR – priemer v Slovenskej republike, n = počet obcí v hodnotenom geologickom celku/okrese

Note: 1 – Paleozoic, 4 – Carbonatic-silicate Mesozoic and Paleogene, 5 – Paleogene Flysch, 6 – Neovolcanic rocks, SR – mean for the Slovak Republic, n = number of municipalities in evaluated geological unit/district

Tab. 4

Priemerné hodnoty vybraných environmentálnych indikátorov vo vybraných oblastiach Slovenskej republiky.
Mean values for selected environmental indicators in selected areas of the Slovak Republic.

Geologický celok/okres	1	6	4	5	Krupina	Bardejov	SR
Environmentálny indikátor	n = 100	n = 303	n = 154	n = 727	n = 36	n = 86	
MIN	302,27	439,73	586,79	524,64	409,08	484,79	629,75
Ca ²⁺ + Mg ²⁺	1,68	2,11	3,45	3,02	1,80	2,75	3,50
Ca ²⁺	43,15	56,13	99,86	88,53	49,30	80,75	93,56
Mg ²⁺	14,70	17,14	23,27	19,67	13,95	17,98	28,29
Cl ⁻	13,18	21,66	21,24	17,14	21,45	13,77	32,96
SO ₄ ²⁺	45,65	49,70	65,38	62,72	34,08	44,96	79,32
NO ₃ ⁻	18,02	26,44	21,72	16,19	28,14	14,84	38,76
PO ₄ ³⁻	0,10	0,37	0,10	0,05	0,62	0,03	0,22
HCO ₃ ⁻	138,29	191,51	323,63	287,65	173,87	282,12	303,85
SiO ₂	13,72	41,72	12,34	11,26	48,79	11,30	18,21
As	0,008 63	0,002 41	0,001 35	0,000 79	0,001 63	0,001 14	0,001 92
Cd	0,000 50	0,001 27	0,000 52	0,000 62	0,002 96	0,000 25	0,001 02
Pb	0,001 42	0,001 34	0,001 21	0,001 25	0,001 09	0,000 94	0,001 36
Sb	0,007 20	0,000 25	0,000 34	0,000 62	0,000 16	0,000 13	0,000 94

Pozn.: 1 – paleozoikum, 4 – karbonatiko-silikátové mezozoikum a paleogén, 5 – flyšový paleogén, 6 – neovulkanity, SR – priemer v Slovenskej republike, n = počet obcí v hodnotenom geologickom celku/okrese, MIN – celková mineralizácia

Note: 1 – Paleozoic, 4 – Carbonatic-silicate Mesozoic and Paleogene, 5 – Paleogene Flysch, 6 – Neovolcanic rocks, SR – mean for the Slovak Republic, n = number of municipalities in evaluated geological unit/district

Tab. 5

Pearsonova a Spearmanova korelácia medzi EI a úmrtnosťou na kardiovaskulárne ochorenia v geologickom prostredí v celku.
 Pearson and Spearman correlation between EI and mortality for CVD for geological environment as a whole.

Parameter	Lineárna korelácia			Spearmanova korelácia		
	r	P	významnosť	R	P	významnosť
Ca ²⁺ + Mg ²⁺ & ReI	-0,0861	0,000	+++	-0,155	0,000	+++
Ca ²⁺ + Mg ²⁺ & SMRI	-0,020 8	0,396	-	-0,045	0,069	-
Ca ²⁺ + Mg ²⁺ & PYIII	-0,084 4	0,001	++	-0,121	0,000	+++
Ca ²⁺ & ReI	-0,081 8	0,001	++	-0,144	0,000	+++
Ca ²⁺ & SMRI	-0,030 6	0,211	-	-0,039	0,114	-
Ca ²⁺ & PYIII	-0,097 6	0,000	+++	-0,116	0,000	+++
Mg ²⁺ & ReI	-0,080 6	0,001	++	-0,174	0,000	+++
Mg ²⁺ & SMRI	0,002	0,934	-	-0,057	0,020	+
Mg ²⁺ & PYIII	-0,045 2	0,065	-	-0,121	0,000	+++
MIN & ReI	-0,067 1	0,006	++	-0,128	0,000	+++
MIN & SMRI	0,000 1	0,996	-	-0,033	0,177	-
MIN & PYIII	-0,070 2	0,004	++	-0,117	0,000	+++
NO ₃ ⁻ & ReI	-0,032 9	0,179	-	-0,095	0,000	+++
NO ₃ ⁻ & SMRI	0,077 7	0,002	++	0,053	0,030	+
NO ₃ ⁻ & PYIII	0,042 5	0,083	-	0,003	0,903	-
PO ₄ ³⁻ & ReI	0,063 4	0,01	+	0,099	0,000	+++
PO ₄ ³⁻ & SMRI	0,135 8	0,000	+++	0,099	0,000	+++
PO ₄ ³⁻ & PYIII	0,094 7	0,000	+++	0,077	0,002	++
As & ReI	-0,013 7	0,576	-	0,028	0,248	-
As & SMRI	-0,013 4	0,584	-	0,039	0,112	-
As & PYIII	-0,019 7	0,421	-	0,092	0,000	+++
Cd & ReI	-0,030 6	0,212	-	-0,013	0,599	-
Cd & SMRI	-0,014 2	0,561	-	0,038	0,124	-
Cd & PYIII	-0,029 1	0,234	-	0,054	0,028	+
Pb & ReI	-0,028 1	0,251	-	-0,017	0,498	-
Pb & SMRI	0,003 4	0,891	-	0,018	0,460	-
Pb & PYIII	-0,007 2	0,77	-	0,008	0,738	-
Sb & ReI	-0,032 3	0,187	-	-0,102	0,000	+++
Sb & SMRI	-0,035 6	0,146	-	0,018	0,466	-
Sb & PYIII	-0,010 6	0,665	-	0,028	0,258	-

Pozn.: r – Pearsonov korelačný koeficient, R – Spearmanov korelačný koeficient, P – hodnota: hladina významnosti = 0,05 – overená závislosť (+), P = 0,01 – vysoká závislosť (++) , P = 0,001 – veľmi vysoká závislosť (+++), MIN – celková mineralizácia

Note: r – Pearson correlation coefficient, R – Spearman correlation coefficient, P – value; level of significance = 0.05 – verified dependence (+), P = 0.01 – high dependence (++) , P = 0.001 – very high dependence (+++)

Tab. 6

Koeficient senzitivity a poradie vplyvu 10 najvplyvnejších prvkov v podzemnej vode na zdravotné indikátory úmrtnosti na kardiovaskulárne ochorenia podľa výpočtov ANN.

Coefficients of sensitivity and order of influence for 10 the most influential elements/parameters in groundwater in relation to health indicators of cardiovascular diseases according to calculations through ANN.

Indikátor	ReI00 – I99		ReI21 – I25		ReI63 – I64		SMRI00 – I99		SMRI21 – I25		SMRI63 – I64		PYLLI00 – I99		PYLLI21 – I25		xP
	s_r	P	s_r	P	s_r	P	s_r	P	s_r	P	s_r	P	s_r	P	s_r	P	
Ca ²⁺ + Mg ²⁺	1,370 0	1	1,153 9	1	1,077 8	1	1,677 1	1	1,159 3	1	1,043 9	2	1,046 2	1	1,102 8	1	1,1
Ca ²⁺	1,211 0	2	1,063 8	2	1,037 1	2	1,386 9	2	1,068 4	2	1,044 9	1	1,006 2	3	1,030 9	2	1,8
Mg ²⁺	1,150 0	3	1,054 6	3	1,019 1	6	1,291 0	3	1,067 8	3	1,040 4	3	1,002 4	6	1,020 8	3	3,2
MIN	1,053 0	4	1,028 0	4	1,032 1	3	1,018 4	4	1,002 8	9	1,019 7	4	1,002 1	7	1,015 5	4	4,1
HCO ₃ ⁻	1,026 0	5	1,027 0	5	1,026 0	4	1,005 5	5	1,011 1	4	1,004 1	9	1,010 0	2	1,008 4	5	4,1
Cl	1,027 0	6	1,018 8	6	1,021 4	5	1,001 0	11	1,002 1	11	1,014 4	5	1,001 7	8	1,000 7	10	6,4
SO ₄ ²⁻	1,009 0	7	1,008 5	7	1,008 3	8	1,001 1	10	1,003 0	8	1,002 1	11	1,002 5	5	1,001 0	8	6,6
NO ₃ ⁻	1,004 0	8	1,007 1	8	1,002 3	10	1,002 2	6	1,001 8	12	1,002 4	10	1,000 8	11	1,000 5	12	7,9
Na ⁺	1,002 1	9	1,001 4	13	1,001 5	12	1,000 7	13	1,001 3	14	1,001 9	12	1,002 7	4	1,001 0	7	8,6
SiO ₂	1,003 0	10	1,001 3	14	1,010 6	7	1,000 6	14	1,002 6	10	1,006 8	6	1,001 1	9	1,000 1	18	9,0
Ba	1,001 0	15	1,001 9	11	1,000 7	14	1,000 3	16	1,007 2	5	1,005 0	7	1,000 0	21	1,000 2	16	10,7
K ⁺	1,001 3	12	1,001 2	15	1,000 2	20	1,000 6	15	1,003 5	7	1,000 6	17	1,000 9	10	1,000 3	14	11,2
²²² Rn	1,000 4	18	1,000 1	26	1,004 9	9	1,000 1	18	1,001 6	13	1,004 5	8	1,000 0	23	1,001 1	6	12,3
pH	1,001 2	19	1,001 7	12	1,000 2	19	1,001	12	1,006 5	6	1,000 3	21	1,000 0	28	1,000 9	9	12,8
PO ₄ ³⁻	1,002 0	11	1,000 4	19	1,000 1	22	1,001 3	9	1,000 7	17	1,000 4	19	1,000 1	19	1,000 0	23	14,1
CHSK _{Mn}	1,000 0	21	1,000 0	31	1,000 3	18	1,001 6	8	1,000 5	20	1,000 5	18	1,000 1	18	1,000 3	13	14,9
Hg	1,000 0	20	1,002 7	9	1,000 0	27	1,000 1	19	1,000 1	24	1,000 1	25	1,000 1	16	1,000 1	20	16,2
Fe	1,000 2	23	1,001 9	10	1,000 2	21	1,000 1	20	1,000 6	19	1,000 0	30	1,000 1	17	1,000 0	29	17,1

Pozn.: s_r – koeficient citlivosti, P – poradie vplyvu, xP – aritmetický priemer poradia vplyvu všetkých hodnotených zdravotných indikátorov, MIN – celková mineralizácia

Note: s_r – coefficient of sensitivity, P – order of influence, xP – arithmetic mean of order of influence for all evaluated health indicators

Tab. 7


Výsledky výpočtov ANN a odvodené limitné hodnoty 10 najvplyvnejších prvkov v podzemnej vode Slovenskej republiky na ReI00 – I99.

Results of calculations of ANN and derived limit values for 10 the most influential chemical elements in groundwater of the Slovak Republic in relation to ReI00 – I99.

Poradie	Parameter	s_r	R ²	Limitný obsah		Optimálny obsah		Hodnotené funkcie závislosti	Obsah*	
				DH	HH	DH	HH		min	max
1.	Ca ²⁺ + Mg ²⁺	1,370	0,992	2,90	9,10	4,40	7,60	konvexná parabola	0,35	7,97
2.	Ca ²⁺	1,211	0,999	89,40	neexistuje	neexistuje	neexistuje	konvexná parabola	9,83	201,01
3.	Mg ²⁺	1,150	0,986	24,30	95,80	42,00	78,10	konvexná parabola	2,45	97,75
4.	MIN	1,053	0,960	553,10	1 263,20	629,40	1 186,80	konvexná parabola	87,30	1412,30
5.	Cl ⁻	1,027	0,988	31,80	neexistuje	neexistuje	neexistuje	konvexná parabola	1,23	143,74
6.	HCO ₃ ⁻	1,026	0,979	241,90	neexistuje	326,10	567,90	konvexná parabola	16,57	592,05
7.	SO ₄ ²⁻	1,009	0,961	neexistuje	73,30	neexistuje	neexistuje	priamka s negatívnou smernicou	9,38	319,50
8.	NO ₃ ⁻	1,004	0,939	neexistuje	37,60	neexistuje	neexistuje	priamka s negatívnou smernicou	1,33	227,09
9.	SiO ₂	1,003	0,999	18,20	neexistuje	neexistuje	neexistuje	konkávna parabola	< 0,000 1	0,20
10.	PO ₄ ³⁻	1,002	0,919	0,20	neexistuje	neexistuje	neexistuje	konkávna parabola	0,00	3,35

Pozn.: s_r – koeficient citlivosti, R² – koeficient determinácie, DH – dolná hranica, HH – horná hranica, *min. – max. obsah prvkov/parametrov v podzemnej vode SR (jednotky v mg . l⁻¹, Ca²⁺ + Mg²⁺ v mmol . l⁻¹), MIN – celková mineralizácia

Note: s_r – coefficient of sensitivity, R² – coefficient of determination, LL – lower limit, UL – upper limit, *minimum – maximum contents of chemical elements/parameters in groundwater of the Slovak Republic (units in mg l⁻¹, Ca + Mg in mmol l⁻¹)



Obr. 4. Vymedzenie limitných hodnôt vplyvných prvkov na Rel00 – I99.
Fig. 4. Definition of limit values for the influential elements on Rel00 – I99.

klasických metód regresnej analýzy nemusí zachytiť komplexnosť problematiky a mohlo by viesť k nesprávnym záverom. Komplexné situácie vyžadujú analytické prístupy riešenia. Preto na analýzu vzťahov medzi chemickým zložením podzemnej vody a úmrtnosťou na CVD (MCVD) používame umelú inteligenciu – umelé neurónové siete (ANN).

ANN sú jednou z najviac využívaných modelovacích techník v mnohých oblastiach výskumu. Podrobný prehľad histórie vzniku a rozvoja neurónových sietí podáva Kriesel (2007). Najvýznamnejšou vlastnosťou neurónových sietí je skutočnosť, že sú univerzálnym aproximátorom funkcie. Univerzálnosť neurónových sietí ako aproximátorov bola matematicky dokázaná (Hornik et al., 1989). Výhodou ANN je vystihnúť aj zložitých nelineárnych závislostí. Nevýhodou ANN je, že nepoznáme mechanizmus pôsobenia hodnôt jednotlivých faktorov na výstupný parameter. Absentuje ekvivalent regresnej rovnice, známej z regresnej analýzy.

Každá z natrénovaných sietí je unikátnym výsledkom, ktorý je jednoznačne definovaný len topológiou siete a vektorom synaptických váh. Globálna analýza citlivosti v neurónových sieťach však poskytuje poznatok o dôležitosti príslušnej vstupnej premennej v natrénovanej sieti. Kovalishyn et al. (1998) a Zurada et al. (1995) predložili niekoľko metód merania senzitivity vstupných premenných ANN. Gevrey et al. (2003) prezentovali a porovnali sedem metód merania dôležitosti vstupných premenných ANN. Pokiaľ je pomer citlivosti (s_r) pri danej vstupnej premennej menší ako jedna, dá sa predpokladať, že jej vyradenie nezniží, ale dokonca zvýši výkonnosť siete (StatSoft, 1999).

Kvalitu neurónovej siete môžeme hodnotiť pomocou viacerých metrík. Najčastejšie používanou je hodnota korelačného koeficientu R , ktorá určuje vzájomný vzťah medzi výstupmi a cieľmi, t. j. odhadnutými hodnotami a hodnotami výstupnej premennej. Hodnota 1 znamená úzku závislosť, hodnota 0 znamená žiadnu závislosť.

Pomocou neurónových sietí sa určovalo poradie vplyvu chemických prvkov v podzemnej vode na CVD spolu s limitnými hodnotami (maximálne prípustným, resp. minimálne potrebným a optimálnym obsahom) koncentrácie chemických prvkov vo vode.

Poradie vplyvu chemických prvkov vo vode na CVD bolo určené na základe hodnoty koeficientu citlivosti s_r . Vplyv na CVD majú tie chemické prvky, pri ktorých koeficient citlivosti je väčší ako 1. Na identifikáciu vplyvných prvkov z pohľadu chemického zloženia podzemnej vody bolo počítaných 100 sietí. Zvolený počet 100 sietí sa ukázal ako plne dostatočný, lebo pri ďalších sieťach už hodnota korelačného koeficientu nerástla, ale stagnovala alebo klesala. Napriek tomu, že výkonnosť (spoľahlivosť) siete je uspokojivá, vplyv jednotlivých chemických prvkov vo vode bol pomerne nízky a bol v prípade každej

siete rôzny. Preto sme na určenie poradia vplyvu chemických prvkov vo vode na ZI použili mediánové hodnoty s_r zo všetkých 100 vypočítaných sietí. Takýto postup použili napr. Opitz a Shavlik (1996), Han et al. (2011), Kourentzes et al. (2014) a Rapant et al. (2015).

Na základe mediánovej hodnoty koeficientu citlivosti (s_r) vypočítanej pre každý chemický prvok hodnotíme vplyv chemického prvku na MCVD. Vplyv je tým väčší, čím je väčšia hodnota s_r . Chemické prvky s hodnotou $s_r < 1$ považujeme za také, ktoré na CVD nevplyvajú. Štatistickú významnosť vypočítaných koeficientov s_r udáva koeficient determinácie R^2 . Čím je vyšší, tým väčšia je štatistická významnosť vypočítaných koeficientov senzitivity. Výsledky výpočtov ANN na získanie tvaru závislosti medzi EI a ZI sme overili metódou decilov, a to spôsobom, ktorý uvádzame. Rozpätie hodnôt obsahu skúmaného prvku vo vode sme rozdelili na decily. V ďalšom kroku sme našli ťažiská bodov, ktorých súradnica x prislúchala do jednotlivých decilov. Následne sme cez ťažiská 2. až 9. decilu preložili polynóm druhého stupňa, resp. priamku. Pri veľmi vplyvných prvkoch je zhoda vynikajúca. S poklesom vplyvu prvkov zhoda existuje, ale podobnosť klesá.

ANN nám umožňujú odvodiť aj limitné hodnoty chemických prvkov vo vode z hľadiska CVD, teda koncentráciu chemických prvkov, pri ktorých je úmrtnosť na CVD čo najnižšia. Vyčleňujeme dva druhy limitných hodnôt, a to limitné (kritické) hodnoty a optimálne limitné hodnoty. Limitné hodnoty predstavujú priesečník namodelovanej krivky obsahu chemických prvkov s priemernou hodnotou zdravotného indikátora. Optimálne hodnoty predstavujú priesečník priemernej hodnoty zdravotného indikátora $\pm$ štandardnú odchýlku zdravotného indikátora. V prípade, keď namodelovaná krivka obsahu chemických prvkov nepretínala priemernú hodnotu zdravotného indikátora, limitné hodnoty sa nedali určiť.

Ako priemernú hodnotu zdravotného indikátora používame bayesovský vyrovnaný priemer zdravotného indikátora, ktorý zohľadňuje počet obyvateľov v jednotlivých obciach. Takýto postup bol použitý napr. v prácach Chaikaw et al. (2009) a Chen et al. (2008).

VÝSLEDKY

Priemerné hodnoty zdravotných indikátorov úmrtnosti na CVD v dvoch najpriaznivejších geologických celkoch z hľadiska zdravotných indikátorov (flyšový paleogén, karbonatiko-silikátové mezozoikum a paleogén) a dvoch najnepriaznivejších geologických celkoch (neovulkanity a paleozoikum) z hľadiska zdravotných indikátorov sú uvedené v tab. 3 spolu s priemernými hodnotami za Slovenskú republiku a dva vybrané okresy SR, budované rôznym geologickým prostredím. Priemerný obsah 10 najvplyvnejších chemických prvkov na úmrtnosť na CVD podľa ANN a štyri klasické potenciálne toxické prvky

spájané so zvýšenou úmrtnosťou na CVD – arzén, kadmiu, olovo a antimón – v uvedených celkoch sú uvedené v tab. 4.

Vybrané výsledky lineárnej a Spearmanovej korelácie medzi EI a ZI z celého geologického prostredia sú uvedené v tab. 5.

V tabuľke 6 a 7 sú uvedené výsledky výpočtov ANN. V tabuľke 6 sú uvedené výsledky koeficientu senzitivity najvplyvnejších chemických prvkov v podzemnej vode v prípade hodnotených zdravotných indikátorov úmrtnosti na CVD spolu s poradím vplyvu jednotlivých prvkov. V tabuľke 7 sú zosumarizované výsledky výpočtov ANN týkajúce sa relatívnej úmrtnosti na CVD – ReI00 – I99 – spolu s odvodenými limitnými hodnotami najvplyvnejších chemických prvkov/zložiek/parametrov v podzemnej vode. Spôsob vymedzenia limitných hodnôt vplyvných prvkov je znázornený grafickou formou na obr. 4.

DISKUSIA

Na základe porovnania výsledkov úmrtnosti na CVD (tab. 3) v jednotlivých geologických celkoch a dvoch vybraných okresoch vyplýva, že existujú medzi nimi výrazné rozdiely. Nižšiu úmrtnosť na CVD dokumentujeme v karbonatických geologických celkoch (mezozoikum, paleogén), a to takmer vo všetkých hodnotených zdravotných indikátoroch. Napríklad sumárny ukazovateľ úmrtnosti na CVD ReI00 – I99 je v najpriaznivejšom geologickom celku (celok 5, flyšový paleogén = 463,32) takmer o 38 %

nižší ako v najnepriaznivejšom geologickom celku (celok 6, neovulkanity = 638,78). Podobný trend pozorujeme aj v prípade porovnania hodnôt ostatných zdravotných indikátorov (s výnimkou ukazovateľa ReI63 – I64, mozgové porážky). Výrazné rozdiely v ukazovateľoch úmrtnosti na kardiovaskulárne ochorenia pozorujeme v prípade dvoch porovnávaných okresov Krupina a Bardejov. Oba sú zasobované pitnou vodou len zo zdrojov podzemnej vody v rámci daného okresu. Oba okresy sa vyznačujú vidieckym typom osídlenia a majú podobný sociálno-ekonomický charakter (Rapant et al., 2014a). Odlišné v oboch okresoch je geologické prostredie. Okres Krupina je celý budovaný horninovým prostredím vulkanitov (vo vzťahu k zdravotným indikátorom je to najmenej priaznivé geologické prostredie). Okres Bardejov je celý budovaný horninovým prostredím flyšového paleogénu (vo vzťahu k zdravotným indikátorom je to najviac priaznivé geologické prostredie). Rozdiel medzi hodnotami ReI00 – I99 v týchto dvoch okresoch je viac ako 80 % a v prípade viacerých čiastkových zdravotných indikátorov (PYLLI00 – I99, PYLLI21 – I25, ReI63 – I64) je rozdiel viac ako 90 až 100 %.

Výraznejšie rozdiely v zdravotných indikátoroch kardiovaskulárnych ochorení v oboch okresoch v porovnaní s rozdielmi medzi jednotlivými geologickými celkami vysvetľujeme tým, že silikátové horniny (kryštalínikum, paleozoikum, vulkanity) sú menej zvodnené a obyvateľstvo je v nich často zasobované pitnou vodou zo vzdialenejších, väčšinou karbonatických hornín, teda s vyšším

Tab. 8

Limitné hodnoty slovenskej normy pre pitnú vodu (NV č. 496/2010 Z. z.)
v porovnaní s našimi odvodenými limitnými hodnotami.

Comparison of limit values of Slovak guideline for drinking water (NV č. 496/2010 Z. z.), with our derived limit values.

Parameter	Jednotka	Limitné hodnoty* NV č. 496/2010 Z. z.	Limitná hodnota		Optimálne hodnoty*	
			DH	HH	DH	HH
$\text{Ca}^{2+} + \text{Mg}^{2+}$	mmol . l ⁻¹	1,1 – 5,0 ^b	2,90	9,10	4,40	7,60
Ca^{2+}	mg . l ⁻¹	> 30 ^b	89,40	–	–	–
Mg^{2+}	mg . l ⁻¹	10 – 30 ^b	24,30	95,80	42,00	78,10
MIN	mg . l ⁻¹	1 000 ^a	553,10	1 263,20	629,40	1 186,80
Cl^-	mg . l ⁻¹	250 ^a	31,80	–	–	–
HCO_3^-	mg . l ⁻¹	–	241,90	–	326,10	567,90
SO_4^{2-}	mg . l ⁻¹	250 ^a	–	73,30	–	–
NO_3^-	mg . l ⁻¹	50 ^a	–	37,60	–	–
SiO_2	mg . l ⁻¹	–	18,20	–	–	–
PO_4^{3-}	mg . l ⁻¹	–	0,20	–	–	–

Pozn.: ^a medzná hodnota, ^b odporúčaná hodnota, DH – dolná hodnota, HH – horná hodnota, *nami odvodené limitné a optimálne hodnoty najvplyvnejších environmentálnych indikátorov vo vzťahu k ReI00 – I99, MIN – celková mineralizácia

Note: ^a limit value, ^b recommended value, DH – lower level, HH – upper level, *our derived limit and optimal contents for the most influential environmental indicators in relation to ReI00 – I99

obsahom Ca^{2+} a Mg^{2+} , ktoré sú na Slovensku oveľa viac zvodnené. Uvedené rozdiely v zdravotných indikátoroch medzi karbonatickými a silikátovými geologickými celkami sú podľa nás spôsobené najmä rozdielnym obsahom Ca^{2+} a Mg^{2+} a „tvrdosťou“ vody. Obsah týchto parametrov je v karbonatických geologických celkoch výrazne vyšší (tab. 4).

Na základe výsledkov Spearmanovej a lineárnej korelácie (tab. 5) nemožno vysloviť žiadne signifikantné závery. Naše premenné (EI a ZI) nemajú normálne rozdelenie a skúmané závislosti nie sú vo všeobecnosti lineárne a často ani monotónne, preto dosiahnuté výsledky nepovažujeme za preukazné. Korelačné koeficienty v oboch prípadoch sú veľmi nízke a vo viac než 90 % prípadov oscilujú medzi hodnotami $\pm < 0,1$. Dôležitá je však skutočnosť, že korelačné koeficienty medzi Ca^{2+} , Mg^{2+} a „tvrdosťou“ vody a zdravotnými indikátormi úmrtnosti na CVD vykazujú vo všetkých prípadoch (aj pri indikátoroch CVD neuvedených v tabuľke) záporné hodnoty, a to väčšinou pri štatisticky významnej korelácii. Táto skutočnosť naznačuje zvýšenú úmrtnosť na CVD pri nízkom (deficitnom) obsahu Ca^{2+} a Mg^{2+} v podzemnej/pitnej vode Slovenskej republiky a pri jej nízkej tvrdosti.

Z výsledkov výpočtov neurónových sietí (tab. 6) sa ako najvplyvnejšie prvky/zložky chemického zloženia podzemnej vody na zdravotné indikátory CVD prejavuje $\text{Ca}^{2+} + \text{Mg}^{2+}$, Ca^{2+} , Mg^{2+} , MIN a HCO_3^- . Týchto 5 environmentálnych indikátorov sa vyskytovalo vo všetkých zdravotných indikátoroch kardiovaskulárnych ochorení medzi desiatimi najvplyvnejšími EI. Ostatné EI uvedené v tab. 6 boli zastúpené medzi desiatimi najvplyvnejšími len pri niektorých ZI úmrtnosti na CVD. Z nich Cl^- , SO_4^{2-} , NO_3^- , Na^+ a SiO_2 majú veľmi nízky priemerný vplyv na hodnotené zdravotné indikátory kardiovaskulárnych ochorení ($xP < 10$).

Medzi najvplyvnejšími EI sa jasne vyčleňujú tri skupiny chemických prvkov/zložiek. Prvú skupinu predstavujú $\text{Ca}^{2+} + \text{Mg}^{2+}$, Ca^{2+} a Mg^{2+} . Tieto tri EI podľa nás majú najväčší vplyv na úmrtnosť na CVD. Vyznačujú sa aj najvyššími hodnotami koeficientu senzitivity s_r . Druhá skupina EI (MIN a HCO_3^-) má podľa nás len stochastický vplyv na úmrtnosť na CVD. Podmienené je to tým, že chemické zloženie podzemnej vody Slovenskej republiky je prevažne Ca–Mg– HCO_3 charakteru. MIN a HCO_3^- vnímate ako indikátory obsahu Ca^{2+} a Mg^{2+} v podzemnej vode. HCO_3^- predstavuje najviac zastúpený anión v podzemnej vode Slovenskej republiky a jeho obsah je spojený najmä s kationmi Ca^{2+} a Mg^{2+} (mineralizácia vody v dôsledku rozpúšťania karbonátov). Podobne je to aj v prípade mineralizácie vody (MIN), ktorej hodnoty v podmienkach SR sú podmienené najmä koncentráciou Ca^{2+} a Mg^{2+} vo vode (najzastúpenejšie kationy) a HCO_3^- (najzastúpenejší anión; Rapant et al., 1996). Tretiu skupinu vplyvných prvkov predstavujú Cl^- , SO_4^{2-} a NO_3^- . Tieto tri parametre sú kla-

sickým príkladom antropogénneho znečistenia podzemnej vody Slovenskej republiky. Ich vplyv na základe hodnôt koeficientu s_r je však výrazne nižší (väčšinou o 1 rád) než vplyv Ca^{2+} , Mg^{2+} a „tvrdosti“ vody. V prípade týchto troch parametrov dôležitú úlohu zohráva skutočnosť, že ich zvýšený obsah v podzemnej vode Slovenska vplyvom antropogénneho znečistenia je sprevádzaný väčšinou zvýšenou koncentráciou Ca^{2+} a Mg^{2+} , ktoré sa preukázali ako najvplyvnejšie z hľadiska úmrtnosti na CVD.

Výsledky výpočtov ANN v prípade ReI00 – I99 sú uvedené v tabuľke 7 spolu s vypočítanými limitnými hodnotami. Ako vplyvné prvky ($s_r > 1$) sa prejavili ešte: SiO_2 , Na^+ , PO_4^{3-} , F, K^+ , pH, Ba, Mn, Zn, ^{222}Rn , ^{226}Ra , Cu, Fe, NO_2^- a Sb. Ako nevplyvné prvky ($s_r \leq 1$) sa zistili: ChSK_{Mn} , NH_4^+ , As, Hg, Cr, Cd, Se, Pb a Al. Výsledky výpočtov ANN v prípade ostatných hodnotených zdravotných indikátorov kardiovaskulárnych ochorení sú dostupné na internetovej stránke www.geology.sk/geohealth. Zaujímavou skutočnosťou je zistenie, že prakticky všetky potenciálne toxické prvky sa prejavili ako nevplyvné, v prípade Sb so zanedbateľným vplyvom ($s_r < 1,000\ 2$). Toto zistenie je v súlade s doterajšími poznatkami o malom vplyve potenciálne toxických prvkov na zdravotný stav obyvateľstva v kontaminovaných územiach po banskej činnosti v Slovenskej republike (Rapant et al., 2014b).

Z dosiahnutých výsledkov je zrejmé, že z hodnotenej, relatívne širokej škály analyzovaných prvkov v podzemnej vode má určujúci vplyv na úmrtnosť na CVD obsah $\text{Ca}^{2+} + \text{Mg}^{2+}$, Ca^{2+} a Mg^{2+} . Viacerými prípadovými štúdiami sa preukázal ich vplyv (deficit) na CVD (Pocock et al., 1980; Sauvant a Pepin, 2000; Ferrándiz et al., 2004; Yang et al., 2006; Kousa et al., 2006; Leurs et al., 2010). Vápnik a horčík sú veľmi dôležité vnútrobunkové kationy a ich význam pre správnu činnosť srdca bol vo svetovej literatúre opísaný už viackrát (Bencko et al., 2011; Kožíšek, 2003, 2005; Rubenowitz-Lundin a Hiscock, 2005; Cotruvo a Bartram, 2009). Vo viacerých prácach sa v súvislosti s výskytom CVD často prikladá veľký význam optimálnej koncentracii Mg^{2+} , ktorý vplýva najmä na hypertenziu (Yang, 1998; Catling et al., 2008; Monarca et al., 2006; Rosanoff, 2013).

Pri vymedzení limitného obsahu sme na základe výpočtov ANN v prípade $\text{Ca}^{2+} + \text{Mg}^{2+}$, Mg^{2+} , MIN a HCO_3^- mohli vyčleniť limitný (minimálne potrebný, maximálne prípustný) aj optimálny obsah (tab. 7, obr. 4), pri ktorom bol výskyt ReI00 – I99 pod priemernou hodnotou EBrEl na Slovensku ($< 771,7$), resp. čo najnižší. V prípade HCO_3^- sa nepodarilo určiť hornú limitnú hodnotu (krivka nepretína priemernú hodnotu EBrEl). V prípade Ca^{2+} , Cl^- , SiO_2 a PO_4^{3-} sme dokázali určiť len dolnú limitnú hodnotu ich obsahu. V prípade SO_4^{2-} a NO_3^- sme dokázali určiť len hornú limitnú hranicu.

V tabuľke 8 uvádzame vypočítané limitné hodnoty najvplyvnejších prvkov v porovnaní s limitnými hodnotami slovenskej normy pre pitnú vodu. V prípade najvplyv-

nejších prvkov ($\text{Ca}^{2+} + \text{Mg}^{2+}$, Ca^{2+} , Mg^{2+}) sú naše limitné hodnoty výrazne vyššie (približne 2- až 3-krát), ako udáva slovenská norma pre pitnú vodu (NV SR č. 496/2010 Z. z.). Pri týchto „zvýšených“ hodnotách koncentrácie Ca^{2+} , Mg^{2+} , $\text{Ca}^{2+} + \text{Mg}^{2+}$ je úroveň úmrtnosti na CVD na Slovensku výrazne nižšia ako celoslovenský priemer.

Na základe výpočtov ANN ako limitné hodnoty v pitnej vode z hľadiska úmrtnosti na CVD môžeme teda navrhnúť v prípade $\text{Ca}^{2+} > 90 \text{ mg} \cdot \text{l}^{-1}$, pri $\text{Mg}^{2+} > 24 \text{ mg} \cdot \text{l}^{-1}$ a pri $\text{Ca}^{2+} + \text{Mg}^{2+} > 2,9 \text{ mmol} \cdot \text{l}^{-1}$.

Horné limity obsahu Ca^{2+} , Mg^{2+} , resp. „tvrdosti“ vody nepovažujeme za príliš dôležité. Podzemná/pitná voda prekračujúca horné hranice (či už limitované v slovenskej norme pre pitnú vodu, alebo nami odvodené limitné hodnoty) sa na úrovni Slovenskej republiky vyskytuje naozaj len ojedinele a takáto podzemná voda sa nepoužíva na zásobovanie obyvateľstva pitnou vodou.

ZÁVER

Z dosiahnutých výsledkov vyplýva, že úmrtnosť na CVD na území Slovenskej republiky je ovplyvňovaná chemickým zložením podzemnej vody, najmä obsahom Ca^{2+} , Mg^{2+} a ich sumou $\text{Ca}^{2+} + \text{Mg}^{2+}$ ($\text{mmol} \cdot \text{l}^{-1}$). Výrazne nižšia úmrtnosť na CVD v porovnaní s celoslovenským priemerom sa zistila pri obsahu týchto parametrov v podzemnej vode v nasledujúcich úrovniach: $\text{Ca}^{2+} > 90 \text{ mg} \cdot \text{l}^{-1}$, $\text{Mg}^{2+} > 24 \text{ mg} \cdot \text{l}^{-1}$ a $\text{Ca}^{2+} + \text{Mg}^{2+} > 2,9 \text{ mmol} \cdot \text{l}^{-1}$. Tieto naše hodnoty obsahu sú približne 2 – 3-krát vyššie ako limitný obsah slovenskej normy pre pitnú vodu (NV č. 496/2010 Z. z.). Odporúčame zvážiť zvýšenie súčasných odporúčaných limitných hodnôt týchto ukazovateľov pitnej vody. Definitívne limitné hodnoty budú určené až po komplexnom spracovaní chemického zloženia podzemnej/pitnej vody vo vzťahu k širokej škále zdravotných indikátorov (okrem CVD aj onkologické ochorenia, ochorenia tráviacej a dýchacej sústavy a zliaz s vnútorným vylučovaním).

Zvýšená úmrtnosť na CVD na území Slovenskej republiky je spätá s nízkym (deficitným) obsahom vápnika a horčíka v podzemnej/pitnej vode a nízkou „tvrdosťou“ vody.

Uvedomujeme si určité nepresnosti, ktoré limitujú naše výsledky. Najdôležitejšou skutočnosťou je to, že ľudia vždy nepijú presne takú istú vodu, akú uvádza naša priemerná analýza vody v obci. Naš model však zahŕňa celé územie Slovenskej republiky (približne 50 000 km^2), celú slovenskú populáciu (približne 5,5 mil. ľudí) rozdelenú do všetkých 2 883 obcí. Máme vyhodnotených viac ako 20 000 chemických analýz vody a 34 prvkov/zložiek a sledujeme úmrtnosť na kardiovaskulárne ochorenia prostredníctvom viacerých zdravotných indikátorov.

O dôležitosti Ca^{2+} a Mg^{2+} v prípade výskytu kardiovaskulárnych ochorení bolo publikovaných nespočetne veľa

prací (napr. Dawson et al., 1978; Shaper et al., 1980; Rylander et al., 1991; Rosborg – ed., 2015).

Kardiovaskulárne ochorenia spolu s onkologickými ochoreniami predstavujú v rozvinutých krajinách rozhodujúce príčiny úmrtí. Obsah Ca^{2+} a Mg^{2+} (resp. „tvrdosť“ vody) však nie sú limitované WHO v pitnej vode. Na základe dosiahnutých výsledkov týkajúcich sa kardiovaskulárnych ochorení odporúčame WHO prehodnotiť stanovenie medzinárodných štandardov obsahu Ca^{2+} a Mg^{2+} v pitnej vode.

Podakovanie

Tento výskum bol realizovaný v rámci projektov *Geohealth* (LIFE10 ENV/SK/000086) a *Life for Krupina* (LIFE12 ENV/SK/000094), ktoré sú podporované finančným nástrojom EÚ pre životné prostredie *Life+ program* a Ministerstvom životného prostredia Slovenskej republiky.



LITERATÚRA

- ANON, 2010: Nariadenie vlády č. 496/2010 Z. z., ktorým sa ustanovujú požiadavky na dosiahnutie dobrého stavu vôd.
- BEAGLEHOLE, R., BONITA R. a KJELLSTROM, T., 1993: Basic Epidemiology. Geneva, World Health Organization.
- BHATNAGAR, A., 2006: Environmental Cardiology: Studying Mechanistic Links Between Pollution and Heart Disease. Circulation research. *J. Amer. Heart Assoc.*, 99, 692 – 705.
- BENCKO, V., HRACH, K., MALÝ, H., PIKHART, J., REISSIGOVÁ, Š., SVAČINA, Š., TOMEČKOVÁ, M. a ZVÁROVÁ, J., 2003a: Biomedická statistika III, Statistické metódy v epidemiológii. Časť 1. Praha, Univerzita Karlova, 236 s.
- BENCKO, V., HRACH, K., MALÝ, H., PIKHART, J., REISSIGOVÁ, Š., SVAČINA, Š., TOMEČKOVÁ, M. a ZVÁROVÁ, J., 2003b: Biomedická statistika III, Statistické metódy v epidemiológii. Časť 2. Praha, Univerzita Karlova, 505 s.
- BENCKO, V., NOVÁK, J. a SUK, M., 2011: Zdraví a prírodné podmienky. (Medicína a geológia). Praha, DOLIN, 389 s.
- CATLING, L. A., ABUBAKAR, I., LAKE, I. R. et al., 2008: A systematic review of analytical observational studies investigating the association between cardiovascular disease and drinking water hardness. *J. Wat. Health* 2, 6 (4), 433 – 442.
- COTRUVO, J. a BARTRAM, J. (eds.), 2009: Calcium and Magnesium in Drinking-water: Public health significance. Geneva, World Health Organization, 180 s.
- DARNLEY, A. G., BJORKLUND, A. a BOLVIKEN, B. et al., 1995: A Global Geochemical Database for Environmental and Resource Management. *Earth Sciences (Paris)*, 19, UNESCO Publishing, 122 s.

- DAWSON, E. B., FREY, M. J., MOORE, T. D. a MCGANITY, J., 1978: Relationship of metal metabolism to vascular disease mortality rates in Texas. *Amer. J. Clinic. Nutrition*, 31, 1 188 – 1 197.
- FERRÁNDIZ, J., ABELLÁN, J. J., GÓMEZ-RUBIO, V., LÓPEZ-QUÍLEZ, A., SANMARTÍN, P., ABELLÁN, C., MARTÍNEZ-BENEITO, M. A., MELCHOR, I., VANACLOCHA, H., ZURRIAGA, O., BALLESTER, F., GIL, J. M., PÉREZ-HOYOS, S. a OCAÑA, R., 2004: Spatial Analysis of the Relationship between Mortality from Cardiovascular and Cerebrovascular Disease and Drinking Water Hardness. *Environ. Health Perspectives*, 112, 9, 1 037 – 1 044.
- GEVREY, M., DIMOPOULOS, I. a LEK, S., 2003: Review and comparison of methods to study the contribution of variables in artificial neural network models. *Ecol. Modelling*, 160, 249 – 264.
- HAN, S., LIU, Y. a YAN, J., 2011: Neural network ensemble method study for wind power prediction. In: *Power and Energy Engineering Conference (APPEEC), 2011 Asia-Pacific*, 1 – 4.
- HORNIK, K., STINCHCOMBE, M. a WHITE, H., 1989: Multilayer feedforward networks are universal approximators. *Neural Network*, 2, 359 – 366.
- CHAIKAEW, N., TRIPATHI, N. K. a SOURIS, M., 2009: International Journal of Health Geographics. *Int. J. Health Geograph.*, 8, 36.
- CHEN, J., ROTH, R. E., NAITO, A. T., LENGERICH, E. J. a MAC-EACHREN, A. M., 2008: Geovisual analytics to enhance spatial scan statistic interpretation: an analysis of US cervical cancer mortality. *Int. J. Health Geograph.*, 7, 1, 57.
- JENÍČEK, M., 1995: Epidemiology, The Logic of Modern Medicine. *Montreal, Epimed*, 322 s., ISBN 0-9698912-0-2.
- KLINDA, J. a LIESKOVSKÁ, Z., 2010: State of the environment report of the Slovak Republic. *Bratislava, Ministry of Environment of the Slovak Republic*, 192 s.
- KORDÍK, J., RAPANT, S., BODÍŠ, D. a SLANINKA, I., 2000: Hydrogeochemické mapy v mierke 1 : 50 000 – prezentácia výsledkov z vybraných regiónov Slovenska. *Podzemná voda (Bratislava)*, 6, 2, 130 – 137.
- KOHÚT, M., KOVACH, V. P., KOTOV, A. B., SALNIKOVA, E. B. a SAVATENKOV, V. M., 1999: Sr and Nd isotope geochemistry of Hercynian granitic rocks from the Western Carpathians – implications for granite genesis and crustal evolution. *Geol. Carpath. (Bratislava)*, 50, 6, 477 – 487.
- KOURENTZES, N., BARROW, D. K. a CRONE, S. F., 2014: Neural network ensemble operators for time series forecasting. *Expert Systems with Applications*, 41, 9, 4 235 – 4 244.
- KOUSA, A., HAVULINNA, A. S., MOLTCHANOVA, E., TASKINEN, O., NIKKARINEN, M., KARVONEN, J. a KARVONEN, M., 2006: Calcium:magnesium ratio in local groundwater and incidence of acute myocardial infarction among males in rural Finland. *Environ. Health Perspectives*, 114, 5, 730 – 734.
- KOVALISHYN, V. V., TETKO, I. V., LUIK, A. I., KHOLODOVYCH, V. V., VILLA, A. E. P. a LIVINGSTONE, D. J., 1998: Neural Network Studies. 3. Variable Selection in the Cascade-Correlation Learning Architecture. *J. Chem. Inf. Comput. Sci*, 38, 651 – 659.
- KOŽÍŠEK, F., 2003: Health significance of drinking water calcium and magnesium. *Prague, National Institute of Public Health*, 29 s.
- KOŽÍŠEK, F., 2005: Health risks from drinking demineralised water. World Health Organization, Geneva, 148 – 163. In: *WHO, 2005: Nutrients in drinking water. Water, Sanitation and Health Protection and the Human Environment. Geneva, World Health Organization*, 186 s.
- KRIESEL, D., 2007: Ein kleiner Überblick über Neuronale Netze. *Bonn, Rheinische Friedrich-Wilhelms Universität*, 238 s.
- LAST, J. M., 2001: A Dictionary of epidemiology. *Oxford University Press, ISBN 0-19-514169-5*.
- LEURS, L. J., SCHOUTEN, L. J., MONS, M. N., GOLDBOHN, R. A. a BRANDT, P. A., 2010: Relationship between Tap Water Hardness, Magnesium, and Calcium Concentration and Mortality due to Ischemic Heart Disease or Stroke in the Netherlands. *Environ. Health Perspectives*, 118, 3, 23 – 26.
- MAHESWARAN, R., MORRIS, S., FALCONER, S., GROSSINHO, A., PERRY, I., WAKEFIELD, J. a ELLIOTT, P., 1999: Magnesium in drinking water supplies and mortality from acute myocardial infarction in north west England. *Heart*, 82, 4, 455 – 460.
- MITCHELL, E., FRISBIE, S. a SARKAR, B., 2011: Exposure to multiple metals from groundwater – a global crisis: Geology, climate change, health effects, testing, and mitigation. *Metalomics, Roy. Soc. geochem.*, 3, 874 – 908.
- MONARCA, S., DONATO, F., ZERBINI, I., CALDERON, R. L. a CRAUN, G. F., 2006: Review of epidemiological studies on drinking water hardness and cardiovascular diseases. *Eur. J. Cardiovascular Prevention Rehabil.*, 13, 4, 495 – 506.
- Nariadenie vlády SR, č. 496/2010 Z. z., ktorým sa mení a dopĺňa nariadenie vlády Slovenskej republiky č. 354/2006 Z. z., ktorým sa ustanovujú požiadavky na vodu určenú na ľudskú spotrebu a kontrolu kvality vody určenej na ľudskú spotrebu.
- NHIC, 2012: Health statistics year book of the Slovak Republic 2011. *Bratislava, National Health Information center*, 257 s.
- OECD, 2013: Health at a Glance 2013: OECD Indicators. *OECD Publishing. http://dx.doi.org/10.1787/health_glance-2013-en*
- OPITZ, D. W. a SHAVLIK, J. W., 1996: Actively searching for an effective neural network ensemble. *Connection Sci.*, 8, 3 – 4, 337 – 354.
- POCOCK, S. J., SHAPER, A. G., COOK, D. G., PACKHAM, R. F., LACEY, R. F., POWELL, P. a RUSSELL, P. F., 1980: British Regional Heart Study: geographic variations in cardiovascular mortality, and the role of water quality. *British Medical J.*, 280, 6 226, 1 243 – 1 249.
- RAHMAN, A. a HUSAIN, T., 2011: Nutrient mineral deficiency in drinking water and its impact on human health. *Water Supply Systems: Demand, Distribution and Pollution*, 95 – 106.
- RAPANT, S., CVEČKOVÁ, V., DIETZOVÁ, Z., FAJČIKOVÁ, K., HILLER, E., FINKELMAN, R. B. a ŠKULTÉTYOVÁ, S., 2014a: The potential impact of geological environment on health status of residents of the Slovak Republic. *Environ. Geochem. Health.*, 36, 543 – 561.
- RAPANT, S., CVEČKOVÁ, V., FAJČIKOVÁ, K., KOHÚT, M. a SEDLÁKOVÁ, D., 2014b: Historical mining areas and their influence on human health. *Eur. J. Biomedical Inf.*, 10, 1, 24 – 34.
- RAPANT, S., FAJČIKOVÁ, K., CVEČKOVÁ, V., ĐURŽA, A., STEHLÍKOVÁ, B., SEDLÁKOVÁ, D. a ŽENISOVÁ, Z., 2015: Chemical composition of groundwater and relative mortality for cardiovascular diseases in the Slovak Republic. *Environ. Geochem. Health.*, 37, 4, 745 – 756.
- RAPANT, S., LETKOVIČOVÁ, M., CVEČKOVÁ, V., FAJČIKOVÁ, K., GALBAVÝ, J. a LETKOVIČ, M., 2010: Environmentálne a zdravotné indikátory Slovenskej republiky. Monografia.

- Bratislava, Št. Geol. Úst. D. Štúra, 279 s. (www.geology.sk/?pg=geois.ms_ezi_en).
- RAPANT, S., RAPOŠOVÁ, M., BODIŠ, D., MARSINA, K. a SLANINKA I., 1999: Environmental-geochemical mapping program in the Slovak Republic. *J. Geochem. Explor.*, 66, 2, 151 – 158.
- RAPANT, S., VRANA, K. a BODIŠ, D., 1996: Geochemický atlas Slovenskej republiky – časť I. Podzemné vody. Monografia. Bratislava, GS SR, 127 s.
- ROSANOFF, A., 2013: The high heart health value of drinking-water magnesium. *Medical Hypotheses*, 81, 6, 1 063 – 1 065.
- ROSBORG, I. (ed.), 2015: Drinking Water Minerals and Mineral Balance Importance, Health Significance, Safety Precautions. *Springer International Publishing Switzerland, Springer Verlag, ISBN 978-3-319-09592-9 (Print) 978-3-319-09593-6 (Online)*, 140 s.
- RUBENOWITZ-LUNDIN, E. a HISCOCK, K., 2005: Water hardness and health effects. Chapter 13. *Principles Medic Geol.*, 331 – 345.
- RYLANDER, R., BONEVIK, H. a RUBENOWITZ, E., 1991: Magnesium and calcium in drinking water and cardiovascular mortality. *Scandinav. J. Work, Environ. & Health*, 17, 91 – 94.
- SAUVANT, M. P. a PEPIN, D., 2000: Geografic variation of the mortality from cerebrovascular disease and drinking water in a french small area (Puy de Dome). *Environ. Res. Sec., A*, 84, 219 – 227.
- SHAPER, A. G., PACKHAM, R. F. a POCOCK, S. J., 1980: The British regional heart study: Cardiovascular mortality and water quality. *J. Environ. Pathol. Toxicol.*, 3, 89 – 111.
- SCHROEDER, H. A. a KRAEMER, L. A., 1974: Cardiovascular Mortality, Municipal Water, and Corrosion. *Archives Environ. Health: Int. J.*, 28, 6, 303 – 311.
- STATSOFT, 1999: Electronic Statistics Textbook. (On-line manual), <http://www.statsoft.com/textbook/statistics-glossary/s/button/s/>
- STURCHIO, E., ZANELATO, M., MINOIA, C. a BEMPORAD, E., 2013: Arsenic: Environmental contamination and exposure in Arsenic: Sources, Environmental Impact, Toxicity and Human Health – A Medical Geology Perspective. *Nova Sci. Publ.*, 3 – 38.
- VRANA, K., RAPANT, S., BODIŠ, D., MARSINA, K., LEXA, J., PRAMUKA, S., MAŇKOVSKÁ, B., ČURLÍK, J., ŠEČÍK, P., VOJTAS, J., DANIEL, J. a LUČIVIAŇSKY, L., 1997: Geochemical Atlas of Slovak Republic at a scale 1 : 1 000 000. *J. Geochem. Explor.*, 60, 7 – 37.
- YANG, CH. Y., CHANG, CH. CH., TSAI, S. S. a CHIU, H. F., 2006: Calcium and magnesium in drinking water and risk of death from acute myocardial infarction in Taiwan. *Environ. Res.*, 101, 407 – 411.
- YANG, CH. Y., 1998: Calcium and magnesium in drinking water and risk of death from cerebrovascular disease. *J. Amer. Heart Dis.*, 412 – 414.
- ZURADA, J. M., EBERHART, R. C. a CLOETE, I., 1995: Determining the Significance of Input Parameters Using Sensitivity Analysis. *Lecture Notes Computer Sci.*, 930, 382 – 388.
- KOHÚT, M., KOVACH, V. P., KOTOV, A. B., SALNIKOVA, E. B. a SAVATENKOV, V. M., 1999. Sr and Nd isotope geochemistry of Hercynian granitic rocks from the Western Carpathians – implications for granite genesis and crustal evolution. *Geol. Carpath. (Bratislava)*, 50, 6, 477 – 487.
- www.geology.sk/geohealth
www.statistics.sk
www.shmu.sk
www.czisk.sk
www.shmu.sk/en

Chemical composition of groundwater and mortality for cardiovascular diseases in the Slovak Republic

Cardiovascular diseases (CVD) have been the commonest cause of death in Slovakia for a long time, representing about 50 % of all causes of death (NHIC, 2012; OECD, 2013). The main risk factors for CVD include stress, genetic predisposition, obesity, regular smoking, excessive alcohol intake, unhealthy eating habits, as well as environmental factors, which include mainly the level of natural element concentrations and anthropogenic contamination of the groundwater (especially drinking water), soil and air.

The main objective of this work is to determine which chemical elements in the groundwater are most closely associated with CVD mortality, and, simultaneously, to suggest limit concentrations (optimum, maximum allowable and minimum required), for which CVD mortality is as low as possible.

Mortality and increased incidence of CVD are often associated with an excess or deficit of several chemical elements in the groundwater/drinking water, most often with a deficiency of Ca^{2+} and Mg^{2+} , and low water hardness (Dawson et al., 1978; Shaper et al., 1980; Rylander et al., 1991; Rahman & Husain, 2011). However, some studies do not confirm this dependence, e.g. Maheswaran et al. (1999). Other work associates CVD with increased concentrations of potentially toxic elements (PTE) in the groundwater/drinking water, mainly As, Cd, Pb, Sb and Ba (Schroeder & Kraemer, 1974; Bhatnagar, 2006; Mitchell et al., 2011; Sturchio et al., 2013).

Groundwater is the most important source of drinking water for most of population in Slovakia; accordingly, our work covers the source of drinking water for approximately 90 % of inhabitants (Klinda & Lieskovská, 2010).

Approximately 20 % of the Slovak population uses water from individual wells for drinking and cooking purposes. About 50 % of the population is supplied with drinking water from local water companies using local water resources with a low discharge (less than 5 l s^{-1}) captured and distributed to water supply pipes in the vicinity of settlements. Only in southern Slovakia (especially in the Quaternary sediments) the population is supplied from large water resources that are distributed across distances of 50 – 100 km.

In the present article we evaluate the association of a wide range of elements commonly analysed in the groundwater with data on mortality for cardiovascular diseases (Tab. 1). Two main datasets included data on chemical composition of groundwater defined as environmental indicators ($> 20,000$ analyses, 34 parameters) and data on health indicators of cardiovascular diseases, tab. 2 (ReI00-I99, ReI21-I25, ReI63-I64, SMRI00-I99, SMRI21-I25, SMRI63-I64, PYLLI00-I99, PYLLI21-I25) were analysed. Both datasets were divided according to geological structure of the Slovak Republic into 8 main geological units (1. Paleozoic, 2. Crystalline, 3. Carbonatic Mesozoic and basal Paleogene, 4. Carbonatic-silicate Mesozoic and Paleogene, 5. Paleogene Flysch, 6. Neovolcanic rocks, 7. Neogene, 8. Quaternary) and analysed through traditional statistical correlations (linear, Spearman) as well as through methods of artificial neural networks, both expressed in the form of mean values for the administrative-territorial units of the Slovak Republic (2,883 municipalities). The neural networks represent unique technique of statistical analysis and innovative method in the field of medical geochemistry used for analysis of relationship between geological environment and human health.

Based on achieved results we can conclude that mortality on cardiovascular diseases in the Slovak Republic is influenced by chemical composition of groundwater/drinking water, mainly Ca^{2+} , Mg^{2+} contents and $\text{Ca}^{2+} + \text{Mg}^{2+}$ levels (Tab. 6, 7). Mortality on CVD is significantly lower compared to Slovak average when groundwater content of these parameters are following: for $\text{Ca}^{2+} > 90 \text{ mg l}^{-1}$, for Mg^{2+} within range of $> 24 \text{ mg l}^{-1}$ and for $\text{Ca}^{2+} + \text{Mg}^{2+} > 2.9 \text{ mmol l}^{-1}$. Our derived limit values are about 2 – 3 times higher compared to limits defined within the Slovak guideline for drinking water and therefore we recommend increasing them. We are aware of certain uncertainties that limit our results. The most important fact is that people do not consume exactly the same water as it is defined by our average groundwater analysis for single municipalities. On the other hand, our model includes whole territory of the Slovak Republic (about 50,000 sqkm), whole Slovak population (about 5.5 millions of people) divided to all 2,883 municipalities. We have evaluated more than 20,000 chemical analyses for groundwater including 34 chemical elements/compounds/parameters and we have analysed mortality for cardiovascular diseases through several health indicators.

There are a number of works dealing with the Ca^{2+} and Mg^{2+} relevance to cardiovascular disease mortality (e.g. Dawson et al., 1978; Shaper et al., 1980; Rylander et al., 1991; Rosborg, 2015). Cardiovascular diseases represent in developed countries key causes of deaths for majority of population. However, Ca^{2+} and Mg^{2+} groundwater contents (or water hardness) are not revised by World Health Organization (WHO) as parameters influencing human health. Based on achieved results in this study we propose to WHO to give consideration to determine international drinking water standards for Ca^{2+} and Mg^{2+} content levels.

Inštrukcie autorom

Etika publikovania, záväzná pre publikovanie v časopise Mineralia Slovaca:

www.geology.sk/new/mineralia_etika_publikovania

1. Geovedný časopis Mineralia Slovaca publikuje scientometricky hodnotné recenzované pôvodné vedecké články s vysokým citačným potenciálom. V úvode príspevku musí autor jasne deklarovať, čím konkrétnym je jeho príspevok prínosný pre rozvoj geovied. Rešeršné štúdie sa publikujú len ojedinele.

2. Články na publikovanie (manuskripty) sa do redakcie zasielajú poštou (dva vytylačené exempláre a CD so všetkými súborami v editovateľnej podobe), alebo e-mailom (editovateľné súbory a kompletná verzia v PDF formáte).

3. Súčasne s článkom je potrebné redakcii poslať autorské vyhlásenie o originalite textu a obrázkov. Kópie obrázkov z iných publikácií musia byť legalizované získaním práva na publikovanie. Vyhlásenie musí obsahovať meno autora (autorov), akademický titul a trvalé bydlisko.

4. Rozsah manuskriptu na publikovanie je najviac 25 rukopisných strán (MS Word, Times New Roman, veľkosť písmen 12 bodov, riadkovanie 1,5) vrátane literatúry, obrázkov a vysvetliviek. V prípade veľkého odborného prínosu sú v ojedinelých prípadoch povolené aj dlhšie články.

5. Články sú publikované v angličtine alebo v slovenčine. Články v slovenčine musia obsahovať anglický preklad názvu, abstraktu, kľúčových slov, resumé a popisov k obrázkom a tabuľkám. Články písané v angličtine musia obsahovať slovenské resumé.

Text

1. Abstrakt stručne sumarizuje článok. Môže mať najviac 200 slov a nemá obsahovať citácie. Počet kľúčových slov je maximálne 6. Text má mať úvod, charakteristiku (stav) skúmaného problému, použitú metodiku, nové zistenia, ich interpretáciu, diskusiu, záver a zoznam literatúry. Vychádzajúce údaje musia byť zreteľne odlišné od interpretácií. V texte musia byť odvolávky na všetky použité obrázky a tabuľky.

2. Hierarchiu nadpisov v texte je potrebné vyznačiť ceruzkou na ľavom okraji strany manuskriptu: 1 – najvyššia, 2 – nižšia, 3 – najnižšia.

3. V texte sa uprednostňuje citácia v zátvorke, napr. (Dubčák, 1987; Hrubý et al., 1988), pred formou ... podľa Dubčáka (1987).

4. Pozícia obrázkov a tabuliek v texte sa označí. Nie je vhodné, aby text v editore MS Word obsahoval vložené obrázky, ale náhľadová verzia v pdf ich má obsahovať.

5. Grécke písmená treba identifikovať na ľavom okraji slovom (napr. sigma). Potrebné je odlišovať pomlčku od spojovníka. Symboly, matematické značky, názvy skamenelín a pod., ktoré sa majú vysádzať kurzívou, autor v rukopise podčiarkne vlnovkou.

Obrázky a tabuľky

1. Ilustrácie a tabuľky vysokej kvality bývajú publikované buď **na šírku stĺpca (81 mm), alebo strany (170 mm)**. Optimálna veľkosť písma a čísel v publikovaných obrázkoch je 2 mm. Články v slovenčine musia mať popisy v obrázkoch a tabuľkách v slovenčine, záhlavie tabuliek a texty pod obrázkami a tabuľkami sú v slovenčine a angličtine. Články v angličtine majú všetky texty v angličtine. **Maximálny rozmer ilustrácie a tabuľky vytylačený v časopise je 170 x 230 mm.** Väčšie (skladané) ilustrácie sú publikované len v ojedinelých prípadoch.

2. Pri počítačovej tvorbe obrázkov odporúčame používať programy s vektorovým zobrazením (Corel Draw, Adobe Illustrator, a pod.). Čiary tzv. **vlasovej hrúbky, softvérového alebo rastrová výplň ploch (napr. v Corel Draw) nie sú prípustné.** Výplne v obrázkoch musia pozostávať zo samostatne vysádzaných objektov.

3. Ilustrácie vrátane fotografií musia obsahovať **grafickú mierku** v centimetrovej či metrovej škále, prípadne sa rozmer zobrazených objektov vyjadri v popise obrázka. Mapy a profile musia mať aj **azimutálnu orientáciu** a jednotné vysvetlivky, ktoré sa uvedú pri prvom obrázku. Zoskupené obrázky, napr. fotografie a diagramy, sa uvádzajú ako jeden obrázok s jednotlivými časťami označenými písmenami (a, b, c, atď.).

4. Pri zasielaní fotografií vo forme počítačových súborov (formáty JPG alebo TIF) sa požaduje rozlíšenie minimálne 600 DPI. Publikovanie farebných ilustrácií môže byť spoplatnené.

Literatúra

1. **Minimálne 50 % citácií musí reprezentovať publikácie od roku 2000.** V zozname literatúry sa v abecednom poradí uvádza len literatúra citovaná v danom článku.

2. Spôsob uvádzania literatúry v zozname literatúry

Knižná publikácia: GAZDA, L. & ČECH, M., 1988: Paleozoikum medzevského príkrovu. *Alfa, Bratislava*, 155 s.

Časopis: VRBA, P., 1989: Strižné zóny v metapelitoch. *Miner. Slov.*, 21, 135 – 142.

Zborník: NÁVESNÝ, D., 1987: Vysokodraselné rhyolity. In: *Romanov, V. (ed.): Stratiformné ložiská gemerika. Spec. publ. Slov. geol. spol., Košice*, 203 – 215.

Manuskript: RADVANSKÝ, F., SLIVKA, B., VIKTOR, J. & SRNKA, T., 1985: Žilné ložiská jedloveckého príkrovu gemerika. Záverečná správa z úlohy SGR-geofyzika. *Manuskript. Archiv Št. Geol. Úst. D. Štúra, Spišská Nová Ves*, 28 s.

3. Pri článku viac ako dvoch autorov sa v texte cituje iba prvý autor s dodatkom et al., ale v zozname literatúry sa uvádzajú všetci.

Instructions to authors

Publication ethics, being obligatory for publishing in the journal Mineralia Slovaca:

www.geology.sk/new/mineralia_publikation_ethics

1. Geoscientific journal Mineralia Slovaca publishes scientometrically valuable original peer-reviewed scientific articles with a high citation potential. In the introduction of each article the author(s) must clearly declare, which innovative data the paper brings for the development of geosciences. The retrieval studies are published only exceptionally.

2. The articles for publishing (manuscripts) must be sent to Editorial Office by post (two printed copies and CD with editable files), or by e-mail (editable files plus complete preview version in PDF format).

3. Simultaneously with the article the Editorial Office must receive the author's proclamation that no part of the manuscript was already published and figures and tables are original as well. Copied illustrations from other publications must contain a copyright.

4. The extent of the manuscript for publishing is limited to 25 manuscript pages (MS Word, 12 points Times New Roman, line spacing 1.5) including figures, tables, explanations and references. In the case of contribution with a high scientific value, the longer manuscripts for publishing are exceptionally permitted.

5. Articles can be published in Slovak or English languages. The title, abstract, key words, shortened text (resumé), as well as description to figures and tables in Slovak articles are published also in English. Articles published in English contain Slovak resumé.

Text

1. Abstract briefly summarizing the article is limited to 200 words, no references are allowed. The maximum number of key words is 6. Text of the article has to contain the introduction, characterization (state) of investigated problem, applied methodology, obtained new data, discussion, conclusion and references. The obtained data must be distinctly separated from interpretations. All applied figures and tables must be referred in the text.

2. The hierarchy of headings in the manuscript must be indicated by a pencil note: 1 – highest level, 2 – lower, 3 – lowermost level.

3. The references in the text prefer parentheses, e.g. (Dubčák, 1987; Hrubý et al., 1988). The form "according to Dubčák (1987)" should be used only exceptionally.

4. Position of figures and tables must be indicated in the manuscript. Editable text of manuscript sent to editorial office must be without figures and tables, though the preview PDF has to contain them in a correct position.

5. Greek letter in the text must be identified at the left margin of the text (e.g. sigma). The text should strictly distinguish the dash from hyphen. Symbols, mathematic signs, names of fossils, etc., which should be printed in italics, must be underlined by the wavy line in the manuscript.

Figures and tables

1. The high quality figures and tables can be published either in **maximum width of column (81 mm) or page (170 mm)**. The optimum size of letters and numbers in the camera-ready figure is 2 mm. Articles published in Slovak contain the Slovak descriptions in figures and tables, the tables headings and descriptions beneath figures and tables are in Slovak and English. English articles contain all texts in English. **Maximum dimension of figures and tables in the journal is 170 x 230 mm. Larger (folded) illustrations are published only exceptionally.**

2. For figures drawing the editorial office recommends the vector graphics editors (Corel Draw, Adobe Illustrator, etc.). The very thin lines (hair lines), the pre-defined software or raster fillings of polygons (e.g. in Corel Draw) are not allowed. The filling must consist from separately set objects.

3. Each illustration including photographs must contain graphic (metric) scale, eventually the dimensions of visualized objects have to be stated in the describing text to figure. Maps and profiles must contain also the azimuth orientation, their detail explanations are stated at the first figure. Grouped figures, e.g. photographs and diagrams, are compiled as one figure with separate parts designated by letters (a, b, c, etc.).

4. The photographs sent as JPG or TIF files are required for having minimum 600 DPI resolution. Publishing of colour illustrations can be charged by a fee.

References

1. **Minimum 50 % of referred works must represent contemporary publications after 2000.** The references in alphanumeric order encompass only literature cited in the article.

2. Examples of referring:

Book: GAZDA, L. & ČECH, M., 1988: Paleozoic of the Medzev nappe. *Alfa, Bratislava*, 155 p.

Journal: VRBA, P., 1989: Shear zones in the metapelite complexes. *Miner. Slov.*, 21, 135–142.

Anniversary volume: NÁVESNÝ, D., 1987: High-potassium rhyolites. In: *Romanov, V. (ed.): Stratiform deposits of Gemericum. Spec. publ. Slov. geol. soc., Košice*, 203–215.

Manuscript: RADVANSKÝ, F., SLIVKA, B., VIKTOR, J. & SRNKA, T., 1985: Vein deposits of the Jedlovce nappe of Gemericum. Final report from the project SGR-geophysics. *Manuscript. Archive Št. Geol. Úst. D. Štúra, Spišská Nová Ves*, 28 p.

3. The article with more than two authors is referred by the name of the first author with the amendment et al., but the list of references contains names of all authors.

OBSAH – CONTENT

PŮVODNÉ ČLÁNKY – ORIGINAL PAPERS

Branislav Fričovský, Ladislav Tometz & Marián Fendek

**Geothermometry techniques in reservoir temperature estimation and conceptual site models construction:
Principles, methods and application for the Bešeňová elevation hydrogeothermal structure, Slovakia**

Využitie geotermometrie pri odhade rezervoárovej teploty a konštrukcii koncepčných modelov:
Princípy, metódy a aplikácie v podmienkach hlbokého rezervoáru hydrogeotermálnej štruktúry
bešeňovskej elevácie, Slovensko.

Zlinská, A.

Terciérna mikrofauna z hlbokých vrtov v železovskej priehlbine (Dunajská panva)

Tertiary microfauna in deep wells of the Železovce depression (Danube Basin)

Veronika Cvečková, Katarína Fajčíková, Ľubomír Jurkovič, Beáta Stehlíková & Stanislav Rapant

**Chemické zloženie podzemnej vody a úmrtnosť na kardiovaskulárne ochorenia
v Slovenskej republike**

Chemical composition of groundwater and mortality for cardiovascular diseases
in the Slovak Republic